

dsm version 2 – what's new?

David L. Miller (University of Rhode Island)

RUWPA seminar, University of St Andrews

18 December 2012

outline (disclaimer)

- ▶ nothing realy “new”

outline (disclaimer)

- ▶ nothing really “new”
- ▶ *not my work!* I just found things and added bugs

outline (disclaimer)

- ▶ nothing really “new”
- ▶ *not my work!* I just found things and added bugs
- ▶ (in fact mostly people just told me about cool stuff)

outline (disclaimer)

- ▶ nothing really “new”
- ▶ *not my work!* I just found things and added bugs
- ▶ (in fact mostly people just told me about cool stuff)
- ▶ dsm version 2


outline (disclaimer)

- ▶ nothing really “new”
- ▶ *not my work!* I just found things and added bugs
- ▶ (in fact mostly people just told me about cool stuff)
- ▶ dsm version 2
- ▶ variance estimation


outline (disclaimer)

- ▶ nothing really “new”
- ▶ *not my work!* I just found things and added bugs
- ▶ (in fact mostly people just told me about cool stuff)
- ▶ dsm version 2
- ▶ variance estimation
- ▶ other tricks


Collect spatially referenced data




Fit detection function



Aggregate to segments



Fit spatial model



dsm version 2

models look like:

```
dsm(N ~ s(x,y), ddf.obj,  
    segment.data, observation.data,  
    family=quasipoisson(), ...)
```

the ... is where the magic happens

mgcv-like syntax

dsm features

- ▶ once you've fitted the model you can treat it like a gam

dsm features

- ▶ once you've fitted the model you can treat it like a `gam`
- ▶ `gam.check`, `predict`, `summary` all do sensible things

dsm features

- ▶ once you've fitted the model you can treat it like a `gam`
- ▶ `gam.check`, `predict`, `summary` all do sensible things
- ▶ also: documentation *and* examples (on CRAN)

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM
 - ▶ detection function

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM
 - ▶ detection function
- ▶ standard approach:

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM
 - ▶ detection function
- ▶ standard approach:
 - ▶ moving block bootstrap for GAM

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM
 - ▶ detection function
- ▶ standard approach:
 - ▶ moving block bootstrap for GAM
 - ▶ likelihood-based intervals for detection function

uncertainty estimation

- ▶ want uncertainty in abundance (integral of the smooth)
- ▶ two sources of uncertainty:
 - ▶ GAM
 - ▶ detection function
- ▶ standard approach:
 - ▶ moving block bootstrap for GAM
 - ▶ likelihood-based intervals for detection function
 - ▶ combine with $CV^2(\hat{N}) = CV^2(\hat{p}) + CV^2(\hat{N}_{GAM})$

uncertainty estimation issues

- ▶ bootstrapping spatial data

uncertainty estimation issues

- ▶ bootstrapping spatial data
 - ▶ autocorrelation

uncertainty estimation issues


- ▶ bootstrapping spatial data
 - ▶ autocorrelation
 - ▶ is this a good idea?

uncertainty estimation issues

- ▶ bootstrapping spatial data
 - ▶ autocorrelation
 - ▶ is this a good idea?
- ▶ propagation of uncertainty from detection function


GAM confidence intervals

- ▶ let's stop using moving block bootstraps!




GAM confidence intervals

- ▶ let's stop using moving block bootstraps!
- ▶ using a bit of GAM theory we can find analytical results



GAM confidence intervals

- ▶ let's stop using moving block bootstraps!
- ▶ using a bit of GAM theory we can find analytical results
- ▶ not *more* assumptions but *different* ones



lpmatrix

- ▶ finding the variance is easy w. identity link.

lpmatrix

- ▶ finding the variance is easy w. identity link.
- ▶ use the “`lpmatrix`”, \mathbf{X}_p s.t. $\hat{\eta}_p = \mathbf{X}_p \hat{\beta}$

lpmatrix

- ▶ finding the variance is easy w. identity link.
- ▶ use the “`lpmatrix`”, \mathbf{X}_p s.t. $\hat{\eta}_p = \mathbf{X}_p \hat{\beta}$
- ▶ maps the model parameters to the linear predictor

lpmatrix

- ▶ finding the variance is easy w. identity link.
- ▶ use the “lpmatrix”, \mathbf{X}_p s.t. $\hat{\eta}_p = \mathbf{X}_p \hat{\beta}$
- ▶ maps the model parameters to the linear predictor
- ▶ so: $\mathbf{V}_{\hat{\eta}_p} = \mathbf{X}_p \mathbf{V}_{\hat{\beta}} \mathbf{X}_p^T$

lpmatrix

- ▶ finding the variance is easy w. identity link.
- ▶ use the “lpmatrix”, \mathbf{X}_p s.t. $\hat{\eta}_p = \mathbf{X}_p \hat{\beta}$
- ▶ maps the model parameters to the linear predictor
- ▶ so: $\mathbf{V}_{\hat{\eta}_p} = \mathbf{X}_p \mathbf{V}_{\hat{\beta}} \mathbf{X}_p^T$
- ▶ non-linear links more complicated...

solution by simulation

- $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:
 - a. Simulate from $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$, to obtain β_b .

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:
 - a. Simulate from $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$, to obtain β_b .
 - b. Calculate $\hat{\eta}_b = \exp(\mathbf{X}_p \beta_b)$ (e.g. if we are using the log-link)

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:
 - a. Simulate from $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$, to obtain β_b .
 - b. Calculate $\hat{\eta}_b = \exp(\mathbf{X}_p \beta_b)$ (e.g. if we are using the log-link)
 - c. Sum over the survey area

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:
 - a. Simulate from $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$, to obtain β_b .
 - b. Calculate $\hat{\eta}_b = \exp(\mathbf{X}_p \beta_b)$ (e.g. if we are using the log-link)
 - c. Sum over the survey area
 2. Calculate the appropriate summary statistics, e.g. median, 95% quantiles etc over b .

solution by simulation

- ▶ $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$
- ▶ could use the following algorithm:
 1. For $b = 1, \dots, N_b$ do the following:
 - a. Simulate from $\beta \sim N(\hat{\beta}, \mathbf{V}_{\hat{\beta}})$, to obtain β_b .
 - b. Calculate $\hat{\eta}_b = \exp(\mathbf{X}_p \beta_b)$ (e.g. if we are using the log-link)
 - c. Sum over the survey area
 2. Calculate the appropriate summary statistics, e.g. median, 95% quantiles etc over b .
- ▶ MVB, suggests that this is unnecessary: “*once you've got a CV so big that the delta-method doesn't work, then your estimate is officially Crap and there is not much point in expending extra effort to work out exactly how Crap!*”

analytic approximate solution

beef up the variance based (roughly) on the uncertainty in the linear predictor:

$$\left(\frac{\partial^2 \log_e \eta}{\partial \eta^2} \Big|_{\eta=\hat{\eta}} \circledast \mathbf{X}_p \right) \mathbf{V}_{\beta} \left(\frac{\partial^2 \log_e \eta}{\partial \eta^2} \Big|_{\eta=\hat{\eta}} \circledast \mathbf{X}_p \right)^T$$

where $\frac{\partial^2 \log_e \eta}{\partial \eta^2} \Big|_{\eta=\hat{\eta}}$ is the vector of second derivatives evaluated at the values of the linear predictor

and \circledast denotes R-style matrix-vector multiplication

step back

- ▶ this is nothing new

step back

- ▶ this is nothing new
- ▶ just an easier way of getting uncertainty estimates for GAMs

step back

- ▶ this is nothing new
- ▶ just an easier way of getting uncertainty estimates for GAMs
- ▶ all in Simon Wood's 2006 book

step back

- ▶ this is nothing new
- ▶ just an easier way of getting uncertainty estimates for GAMs
- ▶ all in Simon Wood's 2006 book
- ▶ we can now combine uncertainty between GAM and d.f. as usual: $CV^2(\hat{N}) = CV^2(\hat{p}) + CV^2(\hat{N}_{GAM})$

step back

- ▶ this is nothing new
- ▶ just an easier way of getting uncertainty estimates for GAMs
- ▶ all in Simon Wood's 2006 book
- ▶ we can now combine uncertainty between GAM and d.f. as usual: $CV^2(\hat{N}) = CV^2(\hat{p}) + CV^2(\hat{N}_{GAM})$
- ▶ in `dsm` as `dsm.var.gam`

variance propagation

Trick from MVB to get detection function uncertainty into the GAM

Our model is of the form:

$$\log(\mathbb{E}[N_j]) = \log(2l_j w \hat{p}(\theta)) + \sum_k f_k(z_k)$$

add in a derivative of $\hat{p}(\theta)$:

$$\log(\mathbb{E}[N_j]) = \log(2l_j w \hat{p}(\theta)) + \frac{d \log(\hat{p}(\theta))}{d\theta} \Big|_{\theta=\hat{\theta}} \cdot \gamma + \sum_k f_k(z_k)$$

should have no effect on $\mathbb{E}[N_j]$

variance propagation (2)

- ▶ Detection function uncertainty now included

variance propagation (2)

- ▶ Detection function uncertainty now included
- ▶ in dsm as `dsm.var.prop`

variance propagation (2)

- ▶ Detection function uncertainty now included
- ▶ in `dsm` as `dsm.var.prop`
- ▶ only works when there are no covariates in d.f.

summary – uncertainty estimation

- ▶ Wood (2006) and Augustin et al. (2009) for more on uncertainty in GAMs

summary – uncertainty estimation

- ▶ Wood (2006) and Augustin et al. (2009) for more on uncertainty in GAMs
- ▶ faster than MBB, in (limited) simulations marginally “better”

summary – uncertainty estimation

- ▶ Wood (2006) and Augustin et al. (2009) for more on uncertainty in GAMs
- ▶ faster than MBB, in (limited) simulations marginally “better”
- ▶ See Williams et al. (2010) for the detection function uncertainty trick (there is an error in the paper!)

select

```
option select=TRUE
```

extra shrinkage allows smooths to be removed from the model

effective degrees of freedom can be shrunk to zero

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value	
s(roughness)	1.401e+00	6	0.661	0.0531	.
s(gchl_winter)	5.999e-07	6	0.000	0.2947	
s(depthm)	8.322e-01	6	3.651	4.83e-08	***
s(x)	5.399e+00	6	15.571	< 2e-16	***
s(y)	3.909e+00	6	13.928	< 2e-16	***

here gchl_winter has been “removed” from the model

more info in ?gam

Tweedie

- ▶ sum of M gamma variables where M is Poisson distributed

Tweedie

- ▶ sum of M gamma variables where M is Poisson distributed
- ▶ one parameter $1 < p \leq 2$

Tweedie

- ▶ sum of M gamma variables where M is Poisson distributed
- ▶ one parameter $1 < p \leq 2$
- ▶ only sensitive to changes >0.1 , so line search feasible in most cases

Tweedie

- ▶ sum of M gamma variables where M is Poisson distributed
- ▶ one parameter $1 < p \leq 2$
- ▶ only sensitive to changes >0.1 , so line search feasible in most cases
- ▶ use GCV/REML score to determine “correct” p

Tweedie

- ▶ sum of M gamma variables where M is Poisson distributed
- ▶ one parameter $1 < p \leq 2$
- ▶ only sensitive to changes >0.1 , so line search feasible in most cases
- ▶ use GCV/REML score to determine “correct” p
- ▶ ?Tweedie or Candy (2004)

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.

```
family=negbin(theta=0.1)
```

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`
- ▶ optimisation is by outer iteration (default) or performance iteration (`optimizer="perf"`)

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`
- ▶ optimisation is by outer iteration (default) or performance iteration (`optimizer="perf"`)
- ▶ [[TALK ABOUT TECHNICAL STUFF]]

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`
- ▶ optimisation is by outer iteration (default) or performance iteration (`optimizer="perf"`)
- ▶ [[TALK ABOUT TECHNICAL STUFF]]
- ▶ default outer optimisation is slower but doesn't have convergence issues

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`
- ▶ optimisation is by outer iteration (default) or performance iteration (`optimizer="perf"`)
- ▶ [[TALK ABOUT TECHNICAL STUFF]]
- ▶ default outer optimisation is slower but doesn't have convergence issues
- ▶ you can find which theta was used with
`b$family$getTheta()`

finding theta for negbin()

- ▶ the theta parameter can be fixed e.g.
`family=negbin(theta=0.1)`
- ▶ or a range can be supplied e.g.
`family=negbin(theta=c(0.01,1))`
- ▶ optimisation is by outer iteration (default) or performance iteration (`optimizer="perf"`)
- ▶ [[TALK ABOUT TECHNICAL STUFF]]
- ▶ default outer optimisation is slower but doesn't have convergence issues
- ▶ you can find which theta was used with
`b$family$getTheta()`
- ▶ `?negbin` has more information

the future!

- ▶ MMPP

the future!

- ▶ MMPP
- ▶ we only think about one way variation detection function \Rightarrow GAM

the future!

- ▶ MMPP
- ▶ we only think about one way variation detection function \Rightarrow GAM
- ▶ temporal inference

the future!

- ▶ MMPP
- ▶ we only think about one way variation detection function \Rightarrow GAM
- ▶ temporal inference
- ▶ GAMMs? (technically possible but untested (by me at least))

the future!

- ▶ MMPP
- ▶ we only think about one way variation detection function ⇒ GAM
- ▶ temporal inference
- ▶ GAMMs? (technically possible but untested (by me at least))
- ▶ talk available on my website
<http://ninepointeighthone.net/about>

the future!

- ▶ MMPP
- ▶ we only think about one way variation detection function ⇒ GAM
- ▶ temporal inference
- ▶ GAMMs? (technically possible but untested (by me at least))
- ▶ talk available on my website
<http://ninepointeighthone.net/about>
- ▶ example analysis at <http://github.com/dill/dsm/wiki>

References

- Augustin, N., Musio, M., Wilpert, von, K., Kublin, E., Wood, S.N. & Schumacher, M. (2009) Modeling spatiotemporal forest health monitoring data. *Journal of the American Statistical Association*, 104, 899–911.
- Candy, S. (2004) Modelling catch and effort data using generalised linear models, the Tweedie distribution, random vessel effects and random stratum-by-year effects. *CCAMLR Science*, 11, 59–80.
- Williams, R., Hedley, S.L., Branch, T.A., Bravington, M.V., Zerbini, A.N. & Findlay, K.P. (2011) Chilean blue whales as a case study to illustrate methods to estimate abundance and evaluate conservation status of rare species. *Conservation Biology*, 25, 526–535.
- Wood, S.N. (2006) Generalized Additive Models: an Introduction with R. Chapman & Hall/CRC.