

Algorithms Lab HS23
Department of Computer Science
Prof. Dr. A. Steger, Prof. Dr. E. Welzl
cadmo.ethz.ch/education/lectures/HS23/algolab

## Exercise – Surveillance Photographs

Sherlock Holmes is solving a case with the aid of his homeless intelligence network. His secret agents located at various intersections in London have taken  $\ell$  surveillance photographs of members of Professor Moriarty's gang. These photographs will be used as crucial evidence during an upcoming trial. The problem is how to safely transport the photographs to a secure place: a safe in a police station.

Inspector Lestrade has k police stations at his disposal, which are scattered around London. In every station there is one policeman and one safe. Every policeman can collect at most one photograph. He can do so by driving from his police station to the photograph location and then back to some police station—not necessarily the one he started from. There the photograph is locked into the safe. Each safe can store at most one photograph.

To move between locations, the policemen use London's network of one-way streets. The streets must be used in the "correct" direction only. In order to avoid the risk of losing multiple photographs in case of an assault, every street can be used by at most one policeman who carries a photograph. In contrast, a street may be used by any number of policemen who do not carry a photograph (are on their way to collect one).

Given the map of the relevant part of London and the locations of police stations and photographs, help Dr. Watson to figure out how many photographs can be collected while preserving the security precautions. Hopefully it will be enough to convince the jury...

**Input** The first line of the input contains the number  $t \le 60$  of test cases. Each of the t test cases is described as follows.

- It starts with a line that contains four integers n m k l, separated by a space and such that  $1 \le n \le 500$ ,  $0 \le m \le 10^4$ , and  $1 \le k, \ell \le 500$ . Here n denotes the number of intersections, m the number of streets, k the number of police stations and  $\ell$  the number of photographs.
- The following line defines the locations of the police stations. Each of the k locations is described by an integer  $x \in \{0, ..., n-1\}$ , denoting the intersection at which the station is situated.
- In the same way, the following line defines the ℓ locations of the photographs.
- Each of the following m lines contains two integers x y, separated by a space and such that  $x, y \in \{0, ..., n-1\}$  and  $x \neq y$ . It indicates that there is a (one-way) street that leads from intersection x to intersection y.

Note that there may be multiple police stations and photographs at a single intersection. Furthermore, there may be multiple parallel streets connecting the same pair of intersections.

**Output** For each test case output the maximum number of photographs that can be collected subject to the constraints in the description.

**Points** There are two groups of test sets, worth 100 points in total.

- 1. For the first group of test sets, worth 40 points, and a corresponding hidden test set worth 10 points, you may assume that the map is symmetric, that is, for any pair of intersections x and y the number of streets from x to y is equal to the number of streets from y to x.
- 2. For the second group of test sets, worth 40 points, and a corresponding hidden testset worth 10 points, there are no additional assumptions.


Corresponding sample test sets are contained in testi.in/out, for  $i \in \{1, 2\}$ .

## Sample Input

## Sample Output

2

3 1



Sample Explanation The figure presents one of the possible ways to collect two photographs. The police stations are denoted  $S_0$  to  $S_3$ , the photographs  $P_0$  to  $P_2$ .

One policeman leaves  $S_0$ , collects  $P_0$  and deposits it in the safe in  $S_1$  (he could go back to  $S_0$  as well). A policeman from  $S_1$  can collect  $P_1$ , but he cannot go with the photograph to  $S_0$  since the street (2,3) was already used.

Another policeman from  $S_2$  collects  $P_2$  and deposits it in  $S_2$  without using any streets (he could deposit it in  $S_3$  as well).