



# Detection of Brno loanvirus (*Loanvirus brunaense*) in common noctule bats (*Nyctalus noctula*) in Southern Russia

Olesia V. Ohlopkova<sup>1,2</sup> · Kristina A. Stolbunova<sup>1,2</sup> · Ilia V. Popov<sup>3</sup> · Igor V. Popov<sup>3,4</sup> · Emmanuel Kabwe<sup>5</sup> · Yuriy N. Davidyuk<sup>5</sup> · Marina A. Stepanyuk<sup>1,2</sup> · Alexey D. Moshkin<sup>1,2</sup> · Yulia V. Kononova<sup>1</sup> · Ekaterina A. Lukbanova<sup>3</sup> · Alexey M. Ermakov<sup>3</sup> · Michael L. Chikindas<sup>3,6,7</sup> · Ivan A. Sobolev<sup>1</sup> · Svetlana F. Khaiboullina<sup>5</sup> · Aleksandr M. Shestopalov<sup>1</sup>

Received: 28 August 2024 / Accepted: 4 December 2024

© The Author(s) under exclusive licence to Sociedade Brasileira de Microbiologia 2024

## Abstract

Hantaviruses that infect humans are rodent-derived viruses with zoonotic potential. Several studies show that before emerging in rodents hantaviruses could emerge in bats, which makes it important to study bat-derived hantaviruses. In this study, we performed PCR screening of hantaviruses in samples from common noctules (*Nyctalus noctula* [182 fecal and 81 blood serum samples]), parti-coloured bats (*Vespertilio murinus* [41 fecal samples]), Kuhl's pipistrelles (*Pipistrellus kuhlii* [15 fecal samples]), and serotine bats (*Eptesicus serotinus* [8 fecal samples]) from Rostov Bat Rehabilitation Center (Rostov-on-Don, Russia) and phylogenetic analysis of detected viruses. As a result, hantaviruses were detected in samples from *N. noctula* bats with an overall prevalence of 4.94% (4/81, 95% CI 0.22–9.66%) in blood serum samples and 1.1% (2/182, 95% CI 0–2.61%) in fecal samples. Phylogenetic analysis revealed that detected hantaviruses are highly homologous to Brno loanviruses (*Loanvirus brunaense*) previously discovered in *N. noctula* bats from Central Europe, which brings some evidence that these are the same bat-derived viruses. This study shows that *Loanvirus brunaense* could be species-specific to the host and has a wide area of habitat: from Central Europe to Southern Russia. These are the first findings of this virus in Southern Russia and Ciscaucasus/Fore-Caucasus. Further studies with wider screening and genomic assays of *Loanvirus brunaense* in bats could reveal trends in the molecular evolution of hantaviruses and provide valuable data for the control of potential spillovers.

**Keywords** Hantaviruses · Bats · Phylogenetics · PCR · Sequencing

Communicated by Luis Augusto Nero.

✉ Olesia V. Ohlopkova  
ohlopkova\_ov@vector.nsc.ru

✉ Igor V. Popov  
ipopov@donstu.ru

<sup>1</sup> Research Institute of Virology, Federal State Budgetary Scientific Institution “Federal Research Center for Fundamental and Translational Medicine”, Novosibirsk 630117, Russian Federation

<sup>2</sup> State Research Center of Virology and Biotechnology “Vector”, Rospotrebnadzor, Koltsovo 630559, Russian Federation

<sup>3</sup> Faculty “Bioengineering and Veterinary Medicine” and Center for Agrobiotechnology, Don State Technical University, Rostov-on-Don 344000, Russian Federation

<sup>4</sup> Division of Immunobiology and Biomedicine, Center of Genetics and Life Sciences, Sirius University of Science and Technology, Federal Territory Sirius 354340, Russian Federation

<sup>5</sup> OpenLab “Gene and Cell Technologies”, Institute of Fundamental Medicine and Biology, Kazan Federal University, Kazan 420008, Russian Federation

<sup>6</sup> Health Promoting Naturals Laboratory, School of Environmental and Biological Sciences, Rutgers State University, New Brunswick, NJ 08901, USA

<sup>7</sup> Department of General Hygiene, I.M. Sechenov First Moscow State Medical University, Moscow 119435, Russia

## Introduction

Bats are recognized as natural hosts for multiple emerging viruses, such as Hendra, Nipah, Ebola, and Marburg viruses, as well as SARS-CoV, MERS-CoV, and SARS-CoV-2 [1–3]. Unique physiological features, such as limited inflammatory response within antiviral immune reactions, ability to fly, and high longevity make bats ideal “bioreactors” for emerging viruses and their unpredictable spillovers all over the globe [4–6]. This emphasizes the need for constant screening of potential bats-derived zoonotic viruses, especially because the order Chiroptera is the second most diverse mammalian order after Rodentia with more than 1400 species in it and the virome of most of these species is yet to be investigated [7]. Even though bats harbor abundant viral diversity, the vast majority of bat viruses have not emerged to cause disease in other animals and humans [8]. This also highlights the importance of studying bats’ viruses, as in combination with the fact that bats relatively peacefully co-exist with viruses due to their unique immune system this data could promote our understanding of viral evolution and its future directions [9, 10].

Hantaviruses, members of the family *Hantaviridae*, pose a huge threat to public health, especially rodent-derived ones belonging to the genus *Orthohantavirus*, as they cause hemorrhagic fever with renal syndrome or hantavirus cardiopulmonary syndrome [11]. Although bats at the moment are not recognized as primary hosts for emerging hantaviruses, the importance of studying these viruses in bats is unconditional, as several phylogenetic studies indicate that hantaviruses might have first appeared in bats before emerging in rodent species [12, 13]. Hantaviruses were previously identified in bats from Africa—Magboi virus in Sierra Leone [14], Mouyassue virus in Cote d’Ivoire [15], Makokou virus in Gabon [16], Kiwira virus in Tanzania and the Democratic Republic of Congo [17]; Asia—Xuan Son virus in Vietnam [18], Huangpi virus, Longquan virus and Laibin virus in China [12, 19], Quezon virus in the Philippines [20], Sarawak mobatvirus in Malaysia [21]; South America - Buri-tiense virus in Brazil [22]; and Europe—Brno loanvirus (*Loanvirus brunaense*) in Czech Republic, Germany, Austria, and Poland [23, 24]. Interestingly, while most of the bat-derived hantaviruses did not show species specificity, *Loanvirus brunaense* was detected only in common noctule bats (*Nyctalus noctula*) in Straková et al. and Dafalla et al. studies [23, 24]. The common noctule (*N. noctula*) is one of the most common bat species throughout Europe and Asia [25]. This bat species is prone to synanthropy—Printz et al. study showed that common noctules are significantly more abundant in urban areas in comparison to forests [26]. Close contact of common noctules with humans and domestic

animals indicates the importance of screening for pathogens in this bat species, as well as in other synanthropic bat species.

This study aimed at screening and phylogenetic characterization of hantaviruses in synanthropic bats from the Rostov Bat Rehabilitation Center at Don State Technical University, Russia.

## Materials and methods

### Sampling

Sampling was conducted from January to March 2023 at the Rostov Bat Rehabilitation Center (Don State Technical University, Rostov-on-Don, Russia). Bats included in this study belonged to four species: common noctule (*N. noctula*), parti-coloured bat (*Vespertilio murinus*), Kuhl’s pipistrelle (*Pipistrellus kuhlii*), and serotine bat (*Eptesicus serotinus*). All bats included in this study were seized by Don State Technical University bat rehabilitation center volunteers starting from November 2022 to March 2023 from households at the request of the owners. Then, seized colonies and separate individuals were transferred to the rehabilitation center, where they underwent veterinary examination and care. Bats of one species were kept collectively (from 5 to 20 individuals) in one box to ensure socialization during rehabilitation with hibernation included. The diet of bats while in the rehabilitation center consisted of superworms and mealworms. Bats were put into controlled hibernation in late November 2022 at a constant temperature of 10 °C and brought out in early April 2023. A minimum of 0.5 g of fecal samples were collected from boxes with hibernating bats: 182 samples from *N. noctula*, 41 samples from *V. murinus*, 15 samples from *P. kuhlii*, and 8 samples from *E. serotinus*. Sampling was conducted once from each box. Obtained fecal samples were stored at –80 °C. Also, for the study, a minimum of 50 µL of blood samples were collected from 81 male *N. noctula* bats. Blood samples were collected by venipuncture of the brachial vein according to the Smith et al. protocol [27]. Blood sampling from female individuals and certain other bat species was not performed due to ethical concerns. For instance, venipuncture in female bats poses a risk for potentially pregnant individuals (there were only female *E. serotinus* in the rehabilitation center at the time of sampling), and the smaller body size of species like *V. murinus* and *P. kuhlii* made it difficult to collect a sufficient amount of blood safely. After blood sampling, plasma specimens were collected by centrifugation (1000 g, 15 min) and stored at –80 °C. All animals included in this study were clinically healthy according to the basic veterinarian examination and showed no signs of infections.

## Hantaviruses detection: RNA extraction, PCR, and sequencing

For viral RNA extraction, 50 mg of each fecal sample was homogenized in 500 µL of PBS with vortexing until a homogeneous suspension was obtained. Then after centrifuging with 11,000 rpm for 4 min, supernatant was obtained, 100 µL of which was used for further RNA extraction. Viral RNA was extracted from blood serum and homogenized fecal supernatant samples using the RIBO-prep kit (AmpliSens, Moscow, Russia). Reverse transcription was performed using the REVERTA-L kit (AmpliSens, Moscow, Russia).

For PCR screening to detect hantaviruses, we used two pairs of primers for L-segments of hantaviruses from the Klempa et al. study: Han-L-F1 (5'-ATG-TAYGTBAGTGCWGATGC-3') and Han-L-R1 (5'-AACCADTCWGTYCCRTCATC-3'); Han-L-F2 (5'-TGCWGATGCHACI-AARTGGTC-3') and Han-L-R2 (5'-GCRTCRTCWGARTGRTGDGCAA-3') [28]. The synthesis of oligonucleotides, including degenerate primers, was carried out by Evrogen (Moscow, Russia). PCR was conducted using PCR 5X ScreenMix (Evrogen, Moscow, Russia) and T100 Thermal Cycler (BioRad, Hercules, California, USA) with the following thermal cycles for the first pair of primers— 1 cycle of 5 min at 93 °C, 34 cycles of 30 s at 93 °C, 20 s at 55 °C, 30 s at 70 °C, 1 cycle of 5 min at 70 °C and 4 °C at the end of the reaction, and for the second pair of primers— 1 cycle of 5 min at 93 °C, 34 cycles of 30 s at 93 °C, 20 s at 59 °C, 25 s at 70 °C, 1 cycle of 5 min at 70 °C and 4 °C at the end of the reaction. Detection of amplification products was performed in 1.5% agarose gel by horizontal electrophoresis using Gel Doc XR+Gel Documentation System (BioRad, Hercules, California, USA). The prevalence of positive signals was calculated as percentages with 95% confidence intervals (95% CI). Extraction and purification of amplification products from agarose gel were conducted using the N-Gel kit (Biolabmix, Novosibirsk, Russia). Sequencing of obtained amplicons was carried out with SeqStudio Genetic Analyzer (Thermo Fisher Scientific, Waltham, Massachusetts, USA).

## Nucleic and amino acids sequence identity analysis

Identity analyses of the sequenced nucleic acids of L-segments of hantaviruses and corresponding amino acid sequences were conducted using MEGA software (version 11) [29] and previously published L-segment sequences of hantaviruses from GenBank (NCBI). The length of analyzed fragments was 313 bp and 105 amino acids.

## Phylogenetic analysis of detected hantaviruses

Detected and sequenced hantavirus L-segments were used for phylogenetic analysis. The majority of sequences for the phylogenetic analysis were obtained from previously published studies [22–24]. Additionally, 10 more sequences of hantaviruses previously associated with human infections were obtained and included 2 samples of each enlisted: *Orthohantavirus seoulense* [30, 31], *Orthohantavirus tulaense* [32, 33], *Orthohantavirus dobravaense* [34], *Hantaan orthohantavirus* [35], and *Orthohantavirus sinnombreense* [36]. All the data was downloaded from the NCBI GenBank [37] database using the Entrez Direct (version 21.6) package [38]. The dataset included both partial and complete coding sequences (CDS) of the L-segment. A total of 103 sequences were included in the analysis. Multiple sequence alignment was performed using the MAFFT (version 7.525) program [39]. To find the best substitution model ModelFinder (implemented in IQ-TREE2) was used [40]. Phylogenetic tree construction was carried out using IQ-TREE2 (version 2.3.0) [41] under the GTR + F + I + G4 substitution model. Branch support was assessed with 1000 ultrafast bootstrap replicates [42]. The resulting phylogenetic tree was visualized using the Interactive Tree Of Life (iTOL, version 6) software [43].

After conducting phylogenetic analysis, a geographic representation of the spatial distribution of detected bat-derived hantaviruses from Rostov-on-Don in this study and closely related viruses described in previous studies was created. The map was constructed utilizing the *rnaturalearth* package (version 1.0.1) for data sourcing [44] and the *ggplot2* package for visualization [45].

The complete bioinformatic pipeline for the phylogenetic analysis and the creation of a spatial distribution map of studied hantaviruses is provided in the Data Availability Statement.

## Results


Positive signals on the detection of RNA of hantaviruses were found in 4 of 81 blood serum samples (4.94%, 95% CI 0.22–9.66%) and 2 of 182 fecal samples (1.1%, 95% CI 0–2.61%) from *N. noctula* bats. There were no positive signals in samples acquired from other bat species included in this study.

We acquired two unique L-segment sequences of the detected hantaviruses. Five out of the six sequenced L-segments were exact duplicates. The sequence submitted to GenBank under accession number OR934973 was obtained from three blood serum samples and two fecal samples. The second unique sequence, submitted under accession number

OR934974, was derived from a blood serum sample of a male *N. noctula* bat. According to the results of the phylogenetic analysis of detected hantaviruses based on L-segments (Fig. 1), they appeared to be in one clade with *Loanvirus brunaense* detected in *N. noctula* bats from the Czech Republic, Germany, Austria, and Poland (Fig. 2). Nucleic and amino acid sequences of L-segments of detected hantaviruses were 93.6–100% identical to Brno loanviruses (Table S1).

## Discussion


In this study, we conducted a screening of hantaviruses in bats from Rostov Bat Rehabilitation Center and a phylogenetic analysis of detected viruses based on sequenced L-segments. Bat rehabilitation centers provide unique opportunities for studying the microbiota of these animals, which, along with rehabilitation itself, meets the goals of bats' conservation, as information about their microbiota is crucial for bats' health management and anti-epizootic measures to prevent potential spillovers of emerging pathogens



**Fig. 1** Maximum likelihood phylogenetic tree constructed using IQ-TREE2 based on complete and partial L segment nucleotide sequences (353 nt) of hantaviruses. Bootstrap values higher than 67 are displayed. The light blue color range indicates Brno loanviruses (*Loanvirus brunaense*) detected in Rostov-on-Don, and the pink range indicates

Brno loanviruses detected in Central Europe in previous studies. Analyzed viral sequences were also grouped according to the order of the host: blue—Rodentia, green—Chiroptera, solid yellow—Eulipotyphla, and purple—Primates (*Homo sapiens* only)





**Fig. 2** Locations, where Brno loanviruses (*Loanvirus brunaense*) were detected: Czechia (Brno) [23], Germany (Hanover, Rossla, Magdeburg), Austria (Lower Austria), Poland (Rybnik, Lubenia) [24], and Russia (Rostov-on-Don, this study)

[46–48]. Bats included in our study are prone to synanthropy and were initially found by citizens of Rostov-on-Don in their households with the following seizing and transportation to the rehabilitation center where they were put into controlled hibernation. Close contact with humans additionally highlights the importance of screening for potential pathogens in bats [49–51].

According to the results of PCR screening, hantaviruses were detected only in samples from *N. noctula*. We should note that fecal samples were taken from hibernating bats, which could alter the abundance of viruses. Previous studies showed that the abundance of some bacteria and overall microbial diversity was lower in the gut microbiota of hibernating bats, *N. noctula* in particular [52–54]. However, this could not be true related to the viral abundance and diversity and needs to be tested in future studies.

Phylogenetic and nucleic and amino acids sequence identity analyses of L-segment sequences revealed that detected hantaviruses are highly homologous to *Loanvirus brunaense*, which brings some evidence that these are the same bat-derived viruses. Brno loanvirus (*Loanvirus brunaense*) was initially identified in two liver and one kidney samples

collected in 2012 and 2013 from common noctules in Brno (Czech Republic) by Straková et al. [23]. A recent study of the hantaviruses' prevalence in bat populations of Central Europe also detected *Loanvirus brunaense* in tissue samples of common noctules: the hantavirus RNA detection rate ranged between 3 of 20 (15%, 95% CI 0–30.65%) for Poland, 3 of 245 (1.2%, 95% CI 0–2.6%) for Germany and 1 of 207 (0.5%, 95% CI 0–1.43%) for Austria [24]. Interestingly, along with the *N. noctula*, these studies included 24 other bat species of bats, and *Loanvirus brunaense* was identified only in *N. noctula* bats [23, 24], as well as in our study, which included samples from 4 different bat species. These show that *Loanvirus brunaense* could be species-specific to the host. Although *Loanvirus brunaense* has been identified in Central and Eastern Europe, we speculate that this virus could be detected in other regions of the common noctule's habitat. *N. noctula* is a partial and differential migrant bat species capable of overcoming distances up to 1,600 km between hibernation areas [55]. The area of geographical distribution of this bat species lies throughout Eurasia, from the British Isles to Southern Asia, where subspecies of this bat are presented, for example, *N. n. labiatus* [25]. Given

the geographical widespread and migration potential of this bat species, there are several concerns about the emerging potential of detected *Loanvirus brunaense*, which could be put to an end by conducting multiple in vitro and in vivo tests. However, we should mention that according to the phylogenetic analysis detected viruses were not closely related to hantaviruses that cause infections in humans, which is also consistent with previous studies [23, 24].

There were several studies on hantavirus surveillance conducted in Russia. Tkachenko et al. analyzed the public health data on the occurrence of hemorrhagic fever with renal syndrome in Russia collected from 2000 to 2022 and found that there were 4.9 cases per 100,000 population. This incidence rate was not equal in different regions: European Russia accounted for 98.5% of these cases, with an incidence rate of 9.7 per 100,000, while Asian Russia reported 1.5% of cases, with a rate of 0.6 per 100,000. Among the causative agents of these infections, there were six rodent-derived hantaviruses, and 97.7% of hemorrhagic fever with renal syndrome cases were caused by *Puumala orthohantavirus*, the natural reservoir of which is the bank vole (*Myodes glareolus*) [56]. Yashina et al. discovered a novel virus belonging to *Orthohantavirus* genus—Academ Virus in samples 14 out of 18 collected from the Siberian mole (*Talpa altaica*) between 2017 and 2021. Phylogenetic analysis revealed that this virus shared a common evolutionary origin with the Bruges virus (*Orthohantavirus brugesense*), previously identified in the European mole (*Talpa europaea*), and is distantly related to other mole-borne hantaviruses [57, 58]. In addition to these findings, Yashina et al. examined the distribution and phylogeny of the Seewis virus (*Orthohantavirus seewisense*) and Altai virus, belonging to the *Orthohantavirus* genus, in Asian Russia. Both viruses co-circulated in shrew populations in Western Siberia, with the Lena River virus (*Mobatvirus lenaense*) and an *Orthohantavirus seewisense* variant co-circulating in Eastern Siberia. These findings suggest Altai virus and related viruses emerged from ancient cross-species transmission among Sorex shrews in Eurasia [59]. Overall, most of the studies of hantaviruses in wild animals conducted earlier in Russia revealed the wide diversity of rodent- and shrew-derived hantaviruses from *Orthohantavirus* and *Mobatvirus* genera. Bat-derived hantaviruses detected in our study were distinct from these genera and appeared in one clade within the *Loanvirus* genus, which expands the current data on hantavirus diversity in small mammals of Russia.

A limitation of this study is that serum samples from male *N. noctula* were used for hantavirus screening due to ethical restrictions on conducting research in a rehabilitation center. Nevertheless, the overall *Loanvirus brunaense* prevalence rate in common noctules in this study generally corresponds to previous epizootic reports [23, 24],

indicating that *Loanvirus brunaense* prevalence may not be related to the sex of the common noctule. Also, the next studies on hantavirus surveillance in bats in Southern Russia should include inner organ samples in addition to blood serum and fecal samples, as in some previous studies, hantaviruses were detected in tissue samples of bats [17, 60]. In addition, further research should be aimed at full-genomic assay of *Loanvirus brunaense* detected in Rostov-on-Don, as this data will provide valuable information regarding the molecular evolution of hantaviruses in bats and could support the evidence that hantaviruses initially appeared in bats and then were transmitted to rodents [61, 62]. Nonetheless, our screening study by itself without full-genomic assessment provides valuable insights into the presence of *Loanvirus brunaense* and its prevalence in the common noctules' population in Rostov-on-Don, which reveals the greater geographical distribution of this virus.

## Conclusion

In this study, we discovered Brno loanvirus (*Loanvirus brunaense*) in *Nyctalus noctula* bats from Rostov Bat Rehabilitation Center (Rostov-on-Don, Russia) with an overall prevalence of 4.94% (4/81, 95% CI 0.22–9.66%) in blood serum samples and 1.1% (2/182, 95% CI 0–2.61%) in fecal samples. These are the first findings of this virus in Southern Russia and Ciscaucasus/Fore-Caucasus. We did not detect *Loanvirus brunaense* in samples from *Vespertilio murinus*, *Pipistrellus kuhlii*, and *Eptesicus serotinus*, which shows that this virus could be species-specific to the host. These results are consistent with previous studies, where *Loanvirus brunaense* was detected in *N. noctula* bats in the Czech Republic, Germany, Austria, and Poland. This study shows that *Loanvirus brunaense* has a wide area of habitat: from Central Europe to Southern Russia, which borders Asia, and probably it is not limited to these regions, as *N. noctula* bats' geographical distribution lies throughout Eurasia. Further studies with wider screening and genomic assays of *Loanvirus brunaense* in bats could reveal trends in the molecular evolution of hantaviruses and provide valuable data for the control of potential spillovers.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s42770-024-01587-5>.

**Acknowledgements** We acknowledge the support of the team of the bat rehabilitation center of the Don State Technical University. Igor V. Popov expresses his deep gratitude to Sergey A. Nedospasov for his guidance and Olga G. Ilchenko for her valuable advice on bat conservation strategies.

**Funding** The reported study was funded by the Russian Science Foun-



dation (project 23-24-00276).

**Data availability** Detected Brno loanviruses (*Loanvirus brunaense*) L-segment sequences have been deposited in the GenBank under accession numbers OR934973 and OR934974. The bioinformatic pipeline used for the data analysis has been deposited in GitHub: <https://github.com/PopovIIIab/PhoBI>.

## Declarations

**Conflict of interest** The authors declare no competing interests.

**Institutional review board statement** The reported study does not involve the results of any in vivo intervention experiments. Handling of animals at the bat rehabilitation center and sampling were approved by the local ethics committee of Don State Technical University (Protocol No. 5 2022).

## References

- Han H-J, Wen H-I, Zhou C-M, Chen F-F, Luo L-M, Liu J (2015) -w, Yu X-J. Bats as reservoirs of severe emerging infectious diseases. *Virus Res* 205:1–6
- Ye Z-W, Yuan S, Yuen K-S, Fung S-Y, Chan C-P, Jin D-Y (2020) Zoonotic origins of human coronaviruses. *Int J Biol Sci* 16(10):1686
- Popov IV, Ohlopkova OV, Donnik IM, Zolotukhin PV, Umanets A, Golovin SN et al (2023) Detection of coronaviruses in insectivorous bats of Fore-Caucasus, 2021. *Sci Rep* 13(1):2306
- Baker M, Schountz T, Wang LF (2013) Antiviral immune responses of bats: a review. *Zoonoses Public Health* 60(1):104–116
- Brook CE, Rozins C, Guth S, Boots M (2023) Reservoir host immunology and life history shape virulence evolution in zoonotic viruses. *PLoS Biol* 21(9):e3002268
- Popov IV, Mazanko MS, Kulaeva ED, Golovin SN, Malinovkin AV, Aleshukina IS et al (2021) Gut microbiota of bats: pro-mutagenic properties and possible frontiers in preventing emerging disease. *Sci Rep* 11(1):21075
- Arnaout Y, Djelouadi Z, Robardet E, Cappelle J, Cliquet F, Touzalin F et al (2022) Genetic identification of bat species for pathogen surveillance across France. *PLoS ONE* 17(1):e0261344
- Van Brussel K, Holmes EC (2022) Zoonotic disease and virome diversity in bats. *Curr Opin Virol* 52:192–202
- Demian WL, Cormier O, Mossman K (2024) Immunological features of bats: resistance and tolerance to emerging viruses. *Trends Immunol* 45(3):198–210
- Zhou H, Ji J, Chen X, Bi Y, Li J, Wang Q et al (2021) Identification of novel bat coronaviruses sheds light on the evolutionary origins of SARS-CoV-2 and related viruses. *Cell* 184(17):4380–4391e14
- Krüger DH, Schönrich G, Klempa B (2011) Human pathogenic hantaviruses and prevention of infection. *Hum Vaccines* 7(6):685–693
- Guo W-P, Lin X-D, Wang W, Tian J-H, Cong M-L, Zhang H-L et al (2013) Phylogeny and origins of hantaviruses harbored by bats, insectivores, and rodents. *PLoS Pathog* 9(2):e1003159
- Souza W, Bello G, Amarilla A, Alfonso H, Aquino V, Figueiredo LTM (2014) Phylogeography and evolutionary history of rodent-borne hantaviruses. *Infect Genet Evol* 21:198–204
- Weiss S, Witkowski PT, Auste B, Nowak K, Weber N, Fahr J et al (2012) Hantavirus in bat, Sierra Leone. *Emerg Infect Dis* 18(1):159
- Sumibcay L, Kadjo B, Gu SH, Kang HJ, Lim BK, Cook JA et al (2012) Divergent lineage of a novel hantavirus in the banana pipistrelle (*Neoromicia nanus*) in Cote d'Ivoire. *Virol J* 9:1–7
- Witkowski PT, Drexler JF, Kallies R, Ličková M, Bokorova S, Mananga GD et al (2016) Phylogenetic analysis of a newfound bat-borne hantavirus supports a laurasiatherian host association for ancestral mammalian hantaviruses. *Infect Genet Evol* 41:113–119
- Weiss S, Sudi LE, Dux A, Mangu CD, Ntinginya NE, Shirima GM et al (2022) Kiwira Virus, a newfound hantavirus discovered in free-tailed bats (Molossidae) in East and Central Africa. *Viruses* 14(11):2368
- Arai S, Nguyen ST, Boldgiv B, Fukui D, Araki K, Dang CN et al (2013) Novel bat-borne hantavirus, Vietnam. *Emerg Infect Dis* 19(7):1159
- Xu L, Wu J, He B, Qin S, Xia L, Qin M et al (2015) Novel hantavirus identified in black-bearded tomb bats, China. *Infect Genet Evol* 31:158–160
- Arai S, Taniguchi S, Aoki K, Yoshikawa Y, Kyuwa S, Tanaka-Taya K et al (2016) Molecular phylogeny of a genetically divergent hantavirus harbored by the Geoffroy's rousette (*Rousettus amplexicaudatus*), a frugivorous bat species in the Philippines. *Infect Genet Evol* 45:26–32
- Zana B, Kemenesi G, Buzás D, Csorba G, Görföl T, Khan FAA et al (2019) Molecular identification of a novel hantavirus in Malaysian bronze tube-nosed bats (*Murina aenea*). *Viruses* 11(10):887
- Barbosa dos Santos M, Koide Albuquerque N, Patroca da Silva S, Silva da Silva F, Damous Dias D, Brito Mendes S et al (2024) A novel hantavirus identified in bats (*Carollia perspicillata*) in Brazil. *Sci Rep* 14(1):6346
- Straková P, Dufkova L, Širmarová J, Salát J, Bartonička T, Klempa B et al (2017) Novel hantavirus identified in European bat species *Nyctalus noctula*. *Infect Genet Evol* 48:127–130
- Dafalla M, Orłowska A, Keleş SJ, Straková P, Schlottau K, Jeske K et al (2023) Hantavirus Brno loanvirus is highly specific to the common noctule bat (*Nyctalus noctula*) and widespread in Central Europe. *Virus Genes* 59(2):323–332
- Lindecke O, Currie SE, Fasel NJ, Fritze M, Kravchenko K, de Assis CK et al (2020) Common Noctule *Nyctalus noctula* (Schreber, 1774). *Handbook of the mammals of Europe*. Springer, pp 1–25
- Printz L, Tschapka M, Vogeler A (2021) The common noctule bat (*Nyctalus noctula*): population trends from artificial roosts and the effect of biotic and abiotic parameters on the probability of occupation. *J Urban Ecol* 7(1):juab033
- Smith CS, De Jong CE, Field HE (2010) Sampling small quantities of blood from microbats. *Acta Chiropterologica* 12(1):255–258
- Klempa B, Fichet-Calvet E, Lecompte E, Auste B, Aniskin V, Meisel H et al (2006) Hantavirus in African wood mouse, Guinea. *Emerg Infect Dis* 12(5):838
- Tamura K, Stecher G, Kumar S (2021) MEGA11: molecular evolutionary genetics analysis version 11. *Mol Biol Evol* 38(7):3022–3027
- Hofmann J, Ulrich RG, Mehl C, Drewes S, Esser J, Løyen M et al (2024) Hantavirus Disease Cluster caused by Seoul Virus, Germany. *Emerg Infect Dis* 30(1):133
- Hofmann J, Weiss S, Kuhns M, Zinke A, Heinsberger H, Krüger DH (2018) Importation of human Seoul virus infection to Germany from Indonesia. *Emerg Infect Dis* 24(6):1099
- Hofmann J, Kramer S, Herrlinger KR, Jeske K, Kuhns M, Weiss S et al (2021) Tula virus as causative agent of hantavirus disease in immunocompetent person. *Ger Emerg Infect Dis* 27(4):1232
- Reynes JM, Carli D, Boukezia N, Debruyne M, Herti S (2015) Tula hantavirus infection in a hospitalised patient, France, June 2015. *Eurosurveillance* 20(50):30095

34. Zelena H, Strakova P, Heroldova M, Mrazek J, Kastl T, Zakovska A et al (2019) Molecular epidemiology of hantaviruses in the Czech Republic. *Emerg Infect Dis* 25(11):2133
35. Seo J-W, Kim C-M, Yun N-R, Lee Y-M, Panchali MJL, Kim D-M (2021) Utility of nested reverse-transcriptase polymerase chain reaction of clinical specimens for early diagnosis of hemorrhagic fever with renal syndrome. *Am J Trop Med Hyg* 105(5):1285
36. Hecht G, Dale AP, Ruberto I, Adame G, Close R, Snyder S-J et al (2023) Detection of Hantavirus during the COVID-19 pandemic, Arizona, USA, 2020. *Emerg Infect Dis* 29(8):1663
37. Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2016) GenBank Nucleic Acids Res 44(D1):D67–D72
38. Kans J (2024) Entrez direct: E-utilities on the UNIX command line. Entrez programming utilities help [Internet]. National Center for Biotechnology Information (US)
39. Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30(4):772–780
40. Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* 14(6):587–589
41. Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32(1):268–274
42. Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2018) UFBot2: improving the ultrafast bootstrap approximation. *Mol Biol Evol* 35(2):518–522
43. Letunic I, Bork P (2024) Interactive tree of life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. *Nucleic Acids Res.*:gkae268
44. South A, South MA (2017) R Package: naturrearth. World Map Data from Natural Earth, Version 101 URL: <https://cran-project.org/web/packages/rnaturrearth/rnaturrearth.pdf> (accessed 23/08/24)
45. Wickham H, Wickham H (2016) Data analysis. Springer
46. Popov IV, Popov IV, Krikunova AA, Lipilkina TA, Derezhina TN, Chikindas ML et al (2023) Gut microbiota composition of Insectivorous Synanthropic and Fructivorous Zoo bats: a direct metagenomic comparison. *Int J Mol Sci* 24(24):17301
47. Jones S, Bell T, Coleman CM, Harris D, Woodward G, Worledge L et al (2022) Testing bats in rehabilitation for SARS-CoV-2 before release into the wild. *Conserv Sci Pract* 4(7):e12707
48. Minor RL, Döpfer D, Lemley EM, Thurber MI (2023) Factors impacting successful Rehabilitation of Big Brown bats (*Eptesicus fuscus*) in a Wisconsin Wildlife Rehabilitation Center: a 5-Year retrospective. *J Zoo Wildl Med* 54(1):32–39
49. Suu-Ire R, Obodai E, Bel-Nono SO, Ampofo WK, Mazet JA, Goldstein T et al (2022) Surveillance for potentially zoonotic viruses in rodent and bat populations and behavioral risk in an agricultural settlement in Ghana. *One Health Outlook* 4(1):6
50. Alkhovsky S, Lenshin S, Romashin A, Vishnevskaya T, Vyshemirsky O, Bulychева Y et al (2022) SARS-like coronaviruses in horseshoe bats (*Rhinolophus* spp.) in Russia, 2020. *Viruses* 14(1):113
51. Latinne A, Nga NTT, Long NV, Ngoc PTB, Thuy HB, Consortium P et al (2023) One health surveillance highlights circulation of viruses with zoonotic potential in bats, pigs, and humans in Viet Nam. *Viruses* 15(3):790
52. Popov IV, Berezinskaia IS, Popov IV, Martiusheva IB, Tkacheva EV, Gorobets VE et al (2023) Cultivable gut microbiota in Synanthropic bats: shifts of its composition and Diversity Associated with hibernation. *Animals* 13(23):3658
53. Xiao G, Liu S, Xiao Y, Zhu Y, Zhao H, Li A et al (2019) Seasonal changes in gut microbiota diversity and composition in the greater horseshoe bat. *Front Microbiol* 10:2247
54. Liu S, Xiao Y, Wang X, Guo D, Wang Y, Wang Y (2023) Effects of Microhabitat temperature variations on the Gut microbiotas of Free-Living Hibernating animals. *Microbiol Spectr* 11(4):e00433–e00423
55. Lehnert LS, Kramer-Schadt S, Teige T, Hoffmeister U, Popa-Lisseanu A, Bontadina F et al (2018) Variability and repeatability of noctule bat migration in Central Europe: evidence for partial and differential migration. *Proc Royal Soc B* 285(1893):20182174
56. Tkachenko E, Kurashova S, Balkina A, Ivanov A, Egorova M, Leonovich O et al (2023) Cases of hemorrhagic fever with renal syndrome in Russia during 2000–2022. *Viruses* 15(7):1537
57. Yashina LN, Panov VV, Abramov SA, Smetannikova NA, Luchnikova EM, Dupal TA et al (2022) Academ virus, a novel hantavirus in the siberian mole (*Talpa altaica*) from Russia. *Viruses* 14(2):309
58. Laenen L, Vergote V, Kafetzopoulou LE, Wawina TB, Vassou D, Cook JA et al (2018) A novel hantavirus of the European mole, Bruges virus, is involved in frequent Nova virus coinfections. *Genome Biol Evol* 10(1):45–55
59. Yashina LN, Abramov SA, Zhigalin AV, Smetannikova NA, Dupal TA, Krivopalov AV et al (2021) Geographic distribution and phylogeny of soricine shrew-borne Seewis virus and Altai virus in Russia. *Viruses* 13(7):1286
60. Souza PJ, Fernandes J, Coelho TA, Cosentino M, D'arc M, Alves PDG et al (2023) Seba's Short-Tailed Bats (*Carollia perspicillata*): A Natural Host of a Newly Discovered Hantavirus in the Brazilian Atlantic Rainforest. *Carollia perspicillata*
61. Bennett SN, Gu SH, Kang HJ, Arai S, Yanagihara R (2014) Reconstructing the evolutionary origins and phylogeography of hantaviruses. *Trends Microbiol* 22(8):473–482
62. Klempa B, Witkowski PT, Popugaeva E, Auste B, Koivogui L, Fichet-Calvet E et al (2012) Sangassou virus, the first hantavirus isolate from Africa, displays genetic and functional properties distinct from those of other murinae-associated hantaviruses. *J Virol* 86(7):3819–3827

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.