
Phylogenomic and Functional Reassessment of *Pseudogymnoascus destructans* Across Mitochondrial and Nuclear Scales



<u>Ilia Popov, Alexey Ermakov, Igor Popov</u> Don State Technical University, Rostov-on-Don, Russia

Introduction



White-Nose Syndrome (caused by *P. destructans or Pd*) is the reason of 90% mortality among hibernating bats every year



times


North America is the most affected by WNS region There are studies indicating that WNS came to America from Europe European strain of *Pd* is not pathogenic

But there were little studies on the genome of Pd Its genome was a mystery:


- ≈20k genes marked "uncharacterized" in RefSeq
- There were no comparative genomic studies
- What is Pd? Where does it come from?
- Does it have non-pathogenic relatives (except for P. pannorum)?

Beimforde et al. ²⁰¹⁴ :	Aguileta et al. ²⁰¹⁴ :	Judd et al. ²⁰²⁴ :
Leotiomycetes and Sordariomycetes are brothers! They are coupled with equivalent fossil- calibrated divergence times!	Sordariomycetes have 0.01 substitutions per site per million years!	Take the retrospective open data on Global Mean Surface Temperature to complement your study!

Materials & Methods



Main Results



- S. Function unknown

 3. C. Satoshystelle rearrapport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Postmanisticrum modification, prosen humover, chapsenese

 E. Amrice acid transport and medacotien

 Q. Postmanisticrum modification, prosen humover, chapsenese

 E. Amrice acid transport and medacotien

 Q. In Decommit directions and operation and conversion

 Q. Postmanisticrum modification, prosen humover, chapsenese

 E. Amrice acid transport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Postmanisticrum modification, prosen humover, chapsenese

 E. Amrice acid transport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Postmanisticrum modification, prosen humover, chapsenese

 E. Amrice acid transport and medacotien

 Q. Castoshystelle rearrapport and medacotien

 Q. Castoshystelle rearrapport


 Q. Castoshystelle rearrapport
- A complete Pd profile (UnP + ChP)
- **B** "characterized" Pd profile (ChP)
- C comparison: Complete vs. UnP vs. ChP

Outcomes

- Bayesian dating: Pd-P. pannorum split = 28.2 Million years ago (MYA);
 5-species MRCA = 140.9 MYA; Leotiomycetes crown = 298 MYA.
 Comparative genomics: Pd mito genome 32.2 kb, 94.4 % ANI to Pp, but shows complete gene-order reshuffling; 3 Antarctic relatives retain collinearity.
- Geographical signal: Pd isolates = North America; closest relatives = Antarctic
- Functional expansion: Re-annotation boosts known Pd protein functions from 2 137 to 12 206; carbohydrate-metabolism genes now 1 206 (+1 097); secondary-metabolite genes 760 (+708).

Supplementary







GitHubLab journal



Contacts Telegram

annual temperature