

VISIT SUMMARY SHEET

PATIENT FIRST
2600 PAPERMILL ROAD
WYOMISSING PA 196103362
484/220-0051

PAUL J RIVERA JR-WC - 180681*1
407 NORTH WYOMISSING BLVD
READING PA 19610
C484.794.7408

COMPLAINT-L Side Pain
ACC DATE/TIME 10.17.22/07:00 1st CONSULT DATE- 10.17.22
SEEN BY- Janet L. McGee, D.O. ON 10/17/22 08:33
ALLERGIES-NKA
MEDICINES TAKING-Elderberry
BP-152/86 PLSE-80 RESP-20
TMP-98.8
WT-275.12 HT-75 BMI-34.3*
SPO2-97% P-80 O2-N
XRAYs- L Hip TAKEN

Lumbosacral Spine TAKEN

INSTRUCTIONS- Musculo-skeletal Pain

Sprain, Fx, Bruise

Sprain, Hip

X-ray Readings

Smoking/Tobacco Use

BP, Adult

BMI-High

RS 10.17.22 09:50

Date of Injury: 10/17/22

Description of Injury/Incident: stepped over something and leg gave out causing him to twist

Light Duty restrictions beginning on 10/18/22:

Lift/Carry Not Permitted

Climb Not permitted

Push/Pull Less than 10 lbs

Additional no bending or twisting, no prolonged standing or walking

Medications prescribed not expected to cause fatigue

Anticipated full duty release date: 10/27/22

DIAGNOSIS- Sprain of ligaments of lumbar spine, initial encounter

Unspecified sprain of left hip, initial encounter

Return to PATIENT FIRST - WYOMISSING (484/220-0051) on 10/24/22 for Unspecified sprain of left hip, initial encounter

PATIENT FIRST

2600 PAPERMILL ROAD
WYOMISSING PA 196103362
484/220-0051

PAUL RIVERA JR-WC

10.17.22 09:47

Musculo-skeletal Pain

Whether due to injury, infection, exercise or other reasons, muscle or joint pain may be treated with Tylenol, Aspirin, ibuprofen, or prescription pain and/or anti-inflammatory medicines to alleviate symptoms. In addition, in case of injury, intermittent local applications of cold packs during the first 3 days & intermittent local heat afterwards may be helpful. Rest & elevation of the affected part may also be of benefit. Sometimes more serious problems like tumors can cause musculo-skeletal pain. Therefore, if your pain persists for prolonged period (more than 2 wks), worsens, or does not seem to be related to movement, return for re-evaluation & further testing to rule out more serious underlying problems.

PATIENT FIRST

2600 PAPERMILL ROAD
WYOMISSING PA 196103362
484/220-0051

PAUL RIVERA JR-WC

10.17.22 09:47

Sprain, Hip

Do not bear weight on the affected side. Use the crutches and follow your doctor's instructions. INITIAL X-RAYS DO NOT ALWAYS SHOW A FRACTURE. With time, a thin crack (not seen on the first x-ray) may become apparent on rexrax. Therefore, you should return for re-evaluation in 1 week.

Arrange to see PATIENT FIRST (484/220-0051) on 10/24/22

PATIENT FIRST

2600 PAPERMILL ROAD
WYOMISSING PA 196103362
484/220-0051

PAUL RIVERA JR-WC

10.17.22 09:47

Sprain, Fx, Bruise

Keep injured area elevated for one to two days. For local swelling, use ice packs intermittently (15 minutes every 2-3 hours while awake) for the first 72 hours. For pain, use aspirin, Tylenol, ibuprofen or prescribed pain medicine. Heat may be used in the form of warm soaks, compresses or heating pad after the first 72 hours. If an elastic bandage has been applied, it should be used until the pain subsides. Injuries requiring splints, slings or crutches should be followed by a return visit before discontinuing the appliance. If an ace bandage has been applied, watch the area for numbness, a blue or dusky color, increasing pain, throbbing, unusual coldness or increased swelling. If these symptoms develop, remove the bandage and reapply snugly but not too tightly. X-RAYS OF BONES FOR EVIDENCE OF FRACTURE ARE OFTEN DIFFICULT TO INTERPRET IF TAKEN SOON AFTER THE INJURY, ESPECIALLY IF THERE IS A THIN CRACK. THEREFORE, IF THE INJURED PART BECOMES MORE TENDER AND IS MORE PAINFUL AFTER 5-7 DAYS OR DOES NOT IMPROVE AS EXPECTED, RETURN FOR RE-EVALUATION. A REPEAT XRAY MAY HAVE TO BE TAKEN. AFTER TIME, A FRACTURE WILL BECOME MORE EVIDENT ON THE XRAY BECAUSE OF BONE RESORPTION THAT OCCURS IN THE DAYS FOLLOWING INJURY. IF PROBLEMS PERSIST RETURN FOR RECHECK.

Arrange to see PATIENT FIRST (484/220-0051) on 10/24/22