

Angular Momentum in Terms of Toroidal and Poloidal Stream Functions

Loren Matilsky

November 15, 2022

1 Prior Work Considered

In this set of notes, we consider 3D nonlinear dynamo simulations that explored magnetic-field amplitudes at a variety of rotation rates (i.e., enough to make scatter plots of various quantities versus Rossby number and thus theoretically address the activity-rotation relation). We aim to determine what region of parameter space has been explored by global models, and in particular, how the rotation-activity relation may or may not have been addressed. The works considered are:

[Christensen & Aubert \(2006\)](#),

[Christensen et al. \(2009\)](#),

[Strugarek et al. \(2017\)](#),

[Guerrero et al. \(2019\)](#),

[Brun et al. \(2022\)](#),

2 Observations of the Stellar Activity-Rotation Relation (ARR)

The stellar activity-rotation relation (hereafter ARR) has a long history of observation and informs astronomers' generally accepted picture of stellar spin-down. The positive correlation between rotational velocity ($v \sin i$) and chromospheric Ca^+ emission (more or less considered to be linearly proportional to strength of the surface magnetic field) was first noted by [Kraft \(1967\)](#). This correlation was made famous in the context of spin-down by the “Skumanich $t^{1/2}$ ” law. This law states that as a star ages (call its age t), its surface magnetic-field strength and rotation rate both decrease like $t^{-1/2}$ ([Skumanich 1972](#)). Since spin-down is believed to be caused by angular-momentum loss from a stellar wind, there thus appears to

be a negative feedback loop between rotation (which produces magnetic field by a convective dynamo) and magnetic field (which drives the stellar wind and slows the rotation).

In addition to chromospheric emission, coronal emission (X-ray luminosity) is also found to be proportional to the rotation rate (e.g., [Walter 1982](#)). Furthermore, the X-ray data show that for a critical rotation rate (corresponding to a critical mixing-length Rossby number of $Ro \sim 0.1$), the ratio of X-ray luminosity to total luminosity saturated to a value of $L_X/L_* \sim 0.001$. It is still unclear whether this saturation regime corresponds a fundamental dynamo process (for example, the inability of the dynamo to convert additional kinetic energy to magnetic field) or simply geometric effects (for example, the stellar surface becoming so peppered with active regions that it cannot accommodate any more dynamo-produced flux) ([Jardine & Unruh 1999](#)). More recent work ([Reiners et al. 2022](#)) claims that the saturation regime indicates a fundamental dynamo process.

Since the detailed properties of the magnetic fields in stellar interiors are almost completely unconstrained (including for the Sun), it is reasonable to assume that the overall dynamo strength (defined here to be the rms magnetic-field strength, with the mean taken over the full volume of the star) is proportional to the surface magnetic-field strength, and thus to the proxies of Ca^+ and X-ray flux. With these assumptions, the ARR tells us about two stellar dynamo regimes: slow rotation (where the Sun lies), in which dynamo strength increases with more rapid rotation, and fast rotation, in which the dynamo strength is insensitive to the rotation rate (a saturation regime). Figure 2 shows the ARR as reported by [Reiners et al. \(2022\)](#). As the Rossby number Ro (ratio of well-measured rotation period P_{rot} to an ad hoc convective turnover time τ) decreases right to left (i.e., the rotation rate increases), the surface magnetic field strength increases up to a critical Rossby number $Ro = 0.13$. After that, faster rotation only leads to marginally increased field strength.

It should be noted that the “Rossby number” used in plots like Figure 2 is really an empirical parameter chosen to make the ARR collapse onto a single curve for all different stellar masses (e.g., [Noyes et al. 1984](#); [Pizzolato et al. 2003](#); [Wright et al. 2011](#)). In general, the empirical convective turnover time scales like $\tau \sim L_*^{1/2}$, which has some basis in mixing-length theory, but the true dependence of τ on stellar parameters could be far more complicated. In any case, simply plotting the normalized activity emission (e.g., L_X/L_*) with respect to rotation period yields an essentially similar ARR, possibly with different saturation breaks (all in the range of $P_{\text{rot}} \sim 1\text{--}3$ days) for different stellar masses (e.g., [Pizzolato et al. 2003](#)).

3 Parameters Spaces Explored

Global, rotating, magnetized spherical shells are characterized by several non-dimensional parameters. These parameters (which completely characterize the Boussinesq system) are:

$$\text{shell aspect ratio} \quad \equiv \quad \beta \quad \equiv \quad \frac{r_i}{r_o}, \quad (1)$$

$$\text{Rayleigh number} \quad \equiv \quad \text{Ra} \quad \equiv \quad \frac{g_o \alpha \Delta \bar{T} H^3}{\nu_o \kappa_o}, \quad (2)$$

$$\text{thermal Prandtl number} \quad \equiv \quad \text{Pr} \quad = \quad \frac{\nu_o}{\kappa_o}, \quad (3)$$




Figure 1: Copied from [Reiners et al. \(2022\)](#), Figure 5: Magnetic field–rotation relation for solar-like and low-mass stars. Symbols for stars rotating slower than $Ro = 0.13$ are colored red, while those of faster rotators are colored blue. Larger and darker symbols indicate higher stellar mass than smaller and lighter symbols. The gray dashed lines show linear fits separately for the slowly rotating stars ($Ro > 0.13$; $\langle B \rangle = 200 G \times Ro^{-1.25}$) and the fast rotators ($Ro < 0.13$; $\langle B \rangle = 2050 G \times Ro^{-0.11}$). Downward open triangles show upper limits for $\langle B \rangle$.

$$\text{Ekman number} \quad \equiv \quad \text{Ek} \quad = \frac{\nu_o}{\Omega_0 H^2}, \quad (4)$$

$$\text{and} \quad \text{magnetic Prandtl number} \quad \equiv \quad \text{Pr}_m \quad = \frac{\nu_o}{\eta_o}. \quad (5)$$

Here, r is the radius, the subscripts “o” and “i” refer to the outer and inner radii of the shell (respectively), $H = r_o - r_i$ is the shell depth, g is the gravitational acceleration, α is the coefficient of thermal expansion, $\Delta\bar{T}$ is the background (superadiabatic, or adverse) temperature difference between the bottom and top of the shell, ν is the kinematic viscosity, κ the thermal diffusivity, Ω_0 the rotation rate, and η the magnetic diffusivity. The “o” subscripts on the diffusivities and g indicate they are in general functions of r . Technically, this radial dependence makes the parameter space accessed by dynamo simulations infinite-dimensional, so hopefully it’s not too important! Just kidding.

If the system is anelastic, there are additional parameters because of the stratification. For a polytrope describing a convection zone, these are:

$$\text{shell density contrast} \quad \equiv \quad \text{DC} \quad \equiv \frac{\bar{\rho}(r_i)}{\bar{\rho}(r_o)} \quad (6)$$

$$\text{or} \quad \text{number of density scale heights} \quad \equiv \quad N_\rho \quad \equiv \ln(\text{DC}), \quad (7)$$

$$\text{polytropic index} \quad \equiv \quad n \quad \equiv \text{something near } \frac{3}{2}, \quad (8)$$

$$\text{or} \quad \text{ratio of specific heats} \quad \equiv \quad \gamma \quad \equiv \frac{c_p}{c_v} = \frac{n+1}{n} = \text{something near } \frac{5}{3}, \quad (9)$$

$$\text{and} \quad \text{dissipation number} \quad \equiv \quad \text{Di} \quad = \frac{g_o H}{c_p \bar{T}_o}. \quad (10)$$

Here, $\bar{\rho}(r)$ is the background density and c_p is the constant-pressure specific heat. The dissipation number arises because viscous and joule heating are typically neglected in the Boussinesq approximation.

If there is also a stable layer in the system (say, below the convection zone), some other parameters are needed: the transition location and width, along with the polytropic index of the stable layer or (if the stable layer isn’t polytropic) the full entropy gradient $d\bar{S}/dr$ in the stable layer. For polytrope, n determines $d\bar{S}/dr$, which will be a complicated function of r but with an order of magnitude of $[n/(3/2) - 1]c_p/r_i$. Regardless, the stable layer introduces an important, final dimensionless number:

$$\text{buoyancy parameter} \quad \equiv \quad \text{B} \quad \equiv \frac{\widetilde{N^2}}{\Omega_0^2}, \quad (11)$$

$$\text{where} \quad N^2 \equiv \frac{g}{c_p} \frac{d\bar{S}}{dr} \quad (12)$$

is the squared buoyancy frequency and the tilde indicates a volume average over the stable layer.

We note also that some authors use a reduced Rayleigh number that is independent of any diffusion coefficient, replacing the diffusion time combination $(H^2/\nu_o)(H^2/\kappa_o)$ with

$1/\Omega_0^2$:

$$\text{reduced Rayleigh number} \quad \equiv \quad \text{Ra}^* \quad \equiv \quad \frac{g_o \alpha \Delta \bar{T}}{\Omega_0^2 H} = \text{RaPr}^{-1} \text{Ek}^2. \quad (13)$$

Finally, in an astrophysical convective system (like a star or the Earth), the adiabatic temperature difference across the convection zone ($\Delta \bar{T}$) is typically unknown. Instead, we can measure the flux driven through the system, and need to define a “flux-based Rayleigh number” with respect to this flux. For example, we usually know the stellar luminosity L_* and radius r . Throughout the star, the different energy fluxes must add up to $\mathcal{F}(r) = L_*/4\pi r^2$. If the convection is vigorous, the convective heat transport, or enthalpy flux, should dominate the other fluxes, i.e., $\mathcal{F} \sim \bar{\rho} c_p u_r T$, where u_r and T are typical radial velocity and temperature perturbations, respectively. If cold plumes dissipate their energy thermally, we might expect $u_r \sim \kappa_o/H$, or

$$\text{first flux-based Rayleigh number} \quad \equiv \quad \text{Ra}_{\text{F1}} \quad \equiv \quad \frac{g_o \mathcal{F}_o H^4}{c_p \bar{\rho}_o \bar{T}_o \nu_o \kappa_o^2} \quad (14)$$

Note that the Boulder crew ([Featherstone & Hindman 2016a,b](#); [Hindman et al. 2020](#); [Matilsky & Toomre 2020](#); [Matilsky et al. 2020](#); [Korre & Featherstone 2021](#); [Camisassa & Featherstone 2022](#)) use a different “volume-averaged” flux-based Rayleigh number:

$$\text{second flux-based Rayleigh number} \quad \equiv \quad \text{Ra}_{\text{F2}} \quad \equiv \quad \frac{\tilde{g} \tilde{F} H^4}{c_p \tilde{\rho} \tilde{T} \tilde{\nu} \tilde{\kappa}^2}, \quad (15)$$

where the tildes refer to volume averages over the convection zone only and $F(r) \equiv \mathcal{F}(r) - \mathcal{F}_{\text{rad}}(r)$ is the flux convection and conduction must carry in equilibrium. In the Rayleigh models, F is determined by the volumetric heating $Q(r)$, i.e., $F(r) = (1/r^2) \int_{r_i}^r Q(x) x^2 dx$.

There is also a reduced flux-based Rayleigh number, used, for example, in [Christensen & Aubert \(2006\)](#):

$$\text{reduced flux-based Rayleigh number} \quad \equiv \quad \text{Ra}_{\text{F}}^* \quad \equiv \quad \frac{1}{4\pi r_i r_o} \frac{g_o \alpha Q_{\text{adv}}}{\rho_o \Omega_0^3 H^2} \quad (16)$$

where Q_{adv} is the “total” (presumably the volume-averaged) integrated convective heat flux. Physically, such a Rayleigh number would be appropriate if the rotation set the typical radial velocity: $u_r \sim H\Omega_0$.

4 Christensen & Aubert (2006)

[Christensen & Aubert \(2006\)](#) explored saturated Boussinesq dynamos in the context of planetary magnetic fields

References

- Brun, A. S., Strugarek, A., Noraz, Q., Perri, B., Varela, J., Augustson, K., Charbonneau, P., & Toomre, J. 2022, “Powering stellar magnetism: Energy transfers in cyclic dynamos of Sun-like stars”, *The Astrophysical Journal*, 926, 21, doi: [10.3847/1538-4357/ac469b](#)

Table 1: Parameter space explored by prior global dynamo simulations. In [Christensen & Aubert \(2006\)](#), the system is Boussinesq with a fixed temperature difference. Only cases with dipolar dynamos ($f_{\text{dip}} > 0.35$) and sufficiently vigorous convection ($\text{Nu} > 2$, where Nu is the Nusselt number) are considered. The min/max supercriticality was estimated from their Figure 4(a)’s x -axis.

Paper	Christensen & Aubert (2006)	Brun et al. (2022)
Ek min	10^{-6}	4×10^{-5}
Ek max	3×10^{-4}	9.0×10^{-3}
Ra/Ra _{crit} min	≈ 5	5.7
Ra/Ra _{crit} max	≈ 50	73
Pr min	0.1	0.25
Pr max	10	0.25
Pr _m min	0.06	1
Pr _m max	10	2

Camisassa, M. E., & Featherstone, N. A. 2022, “Solar-like to antisolar differential rotation: A geometric interpretation”, *The Astrophysical Journal*, 938, 65, doi: [10.3847/1538-4357/ac879f](#)

Christensen, U. R., & Aubert, J. 2006, “Scaling properties of convection-driven dynamos in rotating spherical shells and application to planetary magnetic fields”, *Geophysical Journal International*, 166, 97, doi: [10.1111/j.1365-246x.2006.03009.x](#)

Christensen, U. R., Holzwarth, V., & Reiners, A. 2009, “Energy flux determines magnetic field strength of planets and stars”, *Nature*, 457, 167, doi: [10.1038/nature07626](#)

Featherstone, N. A., & Hindman, B. W. 2016a, “The spectral amplitude of stellar convection and its scaling in the high-Rayleigh-number regime”, *The Astrophysical Journal*, 818, 32, doi: [10.3847/0004-637x/818/1/32](#)

Featherstone, N. A., & Hindman, B. W. 2016b, “The emergence of solar supergranulation as a natural consequence of rotationally constrained interior convection”, *The Astrophysical Journal*, 830, L15, doi: [10.3847/2041-8205/830/1/L15](#)

Guerrero, G., Zaire, B., Smolarkiewicz, P. K., de Gouveia Dal Pino, E. M., Kosovichev, A. G., & Mansour, N. N. 2019, “What sets the magnetic field strength and cycle period in solar-type stars?”, *The Astrophysical Journal*, 880, 6, doi: [10.3847/1538-4357/ab224a](#)

Hindman, B. W., Featherstone, N. A., & Julien, K. 2020, “Morphological classification of the convective regimes in rotating stars”, *The Astrophysical Journal*, 898, 120, doi: [10.3847/1538-4357/ab9ec2](#)

Jardine, M., & Unruh, Y. C. 1999, “Coronal emission and dynamo saturation”, *Astronomy & Astrophysics*, 346, 883

- Korre, L., & Featherstone, N. A. 2021, “On the dynamics of overshooting convection in spherical shells: Effect of density stratification and rotation”, *The Astrophysical Journal*, 923, 52, doi: [10.3847/1538-4357/ac2dea](https://doi.org/10.3847/1538-4357/ac2dea)
- Kraft, R. P. 1967, “Studies of stellar rotation. V. The dependence of rotation on age among solar-type stars”, *The Astrophysical Journal*, 150, 551, doi: [10.1086/149359](https://doi.org/10.1086/149359)
- Matilsky, L. I., Hindman, B. W., & Toomre, J. 2020, “Revisiting the Sun’s strong differential rotation along radial lines”, *The Astrophysical Journal*, 898, 111, doi: [10.3847/1538-4357/ab9ca0](https://doi.org/10.3847/1538-4357/ab9ca0)
- Matilsky, L. I., & Toomre, J. 2020, “Exploring bistability in the cycles of the solar dynamo through global simulations”, *The Astrophysical Journal*, 892, 106, doi: [10.3847/1538-4357/ab791c](https://doi.org/10.3847/1538-4357/ab791c)
- Noyes, R. W., Hartmann, L. W., Baliunas, S. L., Duncan, D. K., & Vaughan, A. H. 1984, “Rotation, convection, and magnetic activity in lower main-sequence stars”, *The Astrophysical Journal*, 279, 763, doi: [10.1086/161945](https://doi.org/10.1086/161945)
- Pizzolato, N., Maggio, A., Micela, G., Sciortino, S., & Ventura, P. 2003, “The stellar activity-rotation relationship revisited: Dependence of saturated and non-saturated x-ray emission regimes on stellar mass for late-type dwarfs”, *Astronomy & Astrophysics*, 397, 147, doi: [10.1051/0004-6361:20021560](https://doi.org/10.1051/0004-6361:20021560)
- Reiners, A., Shulyak, D., Käpylä, P. J., Ribas, I., Nagel, E., Zechmeister, M., Caballero, J. A., Shan, Y., Fuhrmeister, B., Quirrenbach, A., Amado, P. J., Montes, D., Jeffers, S. V., Azzaro, M., Béjar, V. J. S., Chaturvedi, P., Henning, T., Kürster, M., & Pallé, E. 2022, “Magnetism, rotation, and nonthermal emission in cool stars”, *Astronomy & Astrophysics*, 662, A41, doi: [10.1051/0004-6361/202243251](https://doi.org/10.1051/0004-6361/202243251)
- Skumanich, A. 1972, “Time scales for Ca II emission decay, rotational braking, and lithium depletion”, *The Astrophysical Journal*, 171, 565, doi: [10.1086/151310](https://doi.org/10.1086/151310)
- Strugarek, A., Beaudoin, P., Charbonneau, P., Brun, A. S., & do Nascimento, J.-D. 2017, “Reconciling solar and stellar magnetic cycles with nonlinear dynamo simulations”, *Science*, 357, 185, doi: [10.1126/science.aal3999](https://doi.org/10.1126/science.aal3999)
- Walter, F. M. 1982, “On the coronae of rapidly rotating stars. III - an improved coronal rotation-activity relation in late type dwarfs”, *The Astrophysical Journal*, 253, 745, doi: [10.1086/159675](https://doi.org/10.1086/159675)
- Wright, N. J., Drake, J. J., Mamajek, E. E., & Henry, G. W. 2011, “The stellar-activity-rotation relationship and the evolution of stellar dynamos”, *The Astrophysical Journal*, 743, 48, doi: [10.1088/0004-637x/743/1/48](https://doi.org/10.1088/0004-637x/743/1/48)