

# One Stop for Physics Practice for **NEET**

Monday - Friday ; 10 PM

**1 Series = Questions from Top Books**

## Vectors

### Properties & Resolution

**1**




**Tamanna Chaudhary**  
(Physics Expert)



**physics\_tcarmy**



**a.sane.hurricane**



# Tamanna Chaudhary

Expert in NEET UG

One of the most followed Educators, Tamanna, won the 2020 Unacademy People's Choice Award. Her student secured 99.56 percentile in Physics.

**98M** Watch mins

**2M** Watch mins (last 30 days)

**254K** Followers

**14K** Dedications



# 777



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

A force of 5 N acts on a particle along a direction making an angle of  $60^\circ$  with vertical. Its vertical component be

- A) 10 N
- B) 3 N
- C) 4 N
- D) 2.5 N



Position of a particle in a rectangular-co-ordinate system is (3, 2, 5). Then its position vector will be

- A)  $3\hat{i} + 5\hat{j} + 2\hat{k}$
- B)  $3\hat{i} + 2\hat{j} + 5\hat{k}$
- C)  $5\hat{i} + 3\hat{j} + 2\hat{k}$
- D) None of these



If  $\vec{A} = 2\hat{i} + 4\hat{j} - 5\hat{k}$  the direction of cosines of the vector  $\vec{A}$  are

- A)  $\frac{2}{\sqrt{45}}, \frac{4}{\sqrt{45}} \text{ and } \frac{-5}{\sqrt{45}}$
- B)  $\frac{1}{\sqrt{45}}, \frac{2}{\sqrt{45}} \text{ and } \frac{3}{\sqrt{45}}$
- C)  $\frac{4}{\sqrt{45}}, 0 \text{ and } \frac{4}{\sqrt{45}}$
- D)  $\frac{3}{\sqrt{45}}, \frac{2}{\sqrt{45}} \text{ and } \frac{5}{\sqrt{45}}$



How many minimum number of coplanar vectors having different magnitudes can be added to give zero resultant

- A) 2
- B) 3
- C) 4
- D) 5



Given vector  $\vec{A} = 2\hat{i} + 3\hat{j}$ , the angle between  $\vec{A}$  and y-axis is

- A)  $\tan^{-1} 3/2$
- B)  $\tan^{-1} 2/3$
- C)  $\sin^{-1} 2/3$
- D)  $\cos^{-1} 2/3$



If a unit vector is represented by  $0.5\hat{i} + 0.8\hat{j} + c\hat{k}$ , then the value of  $c$  is

- A) 1
- B)  $\sqrt{0.11}$
- C)  $\sqrt{0.01}$
- D)  $\sqrt{0.39}$





Which of the following is a scalar quantity

- A) Displacement
- B) Electric field
- C) Acceleration
- D) Work

7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

1. A vector has component along the  $X$ -axis equal to 25 unit and along the  $Y$ -axis equal to 60 unit. Find the magnitude and direction of the vector.



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

7. Write the unit vector in the direction of  $\vec{A} = 5\vec{i} + \vec{j} - 2\vec{k}$ .

7777



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

1. Is a vector necessarily changed if it is rotated through an angle ?

7777



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

8. Can a vector have zero component along a line and still have nonzero magnitude ?

7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

4. The component of a vector is
- (a) always less than its magnitude
  - (b) always greater than its magnitude
  - (c) always equal to its magnitude
  - (d) none of these.



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

10. Suppose  $\vec{a}$  is a vector of magnitude 4.5 unit due north.  
What is the vector (a)  $3\vec{a}$ , (b)  $-4\vec{a}$ ?



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

# One Stop for Physics Practice for **NEET**

Monday - Friday ; 10 PM

1 Series = Questions from Top Books

## Vectors

### Components of a Vector

2



**Tamanna Chaudhary**  
(Physics Expert)



**physics\_tarmy**



**a.sane.hurricane**





# Tamanna Chaudhary

Expert in NEET UG

One of the most followed Educators, Tamanna, won the 2020 Unacademy People's Choice Award. Her student secured 99.56 percentile in Physics.

**98M** Watch mins

**2M** Watch mins (last 30 days)

**254K** Followers

**14K** Dedications



# 7777



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

# One Stop for Physics Practice for NEET

1 Series = Questions from Top Books



Tamanna Chaudhary  
117K subscribers

## Books Covered in this Session-



HC Verma

- Done in Vectors-1



SL Arora



DC Pandey



TC Selected



NEWTON'S APPLE



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

# One Stop for Physics Practice for **NEET**

**1 Series = Questions from Top Books**



Tamanna Chaudhary ✓  
117K subscribers

## Details about the series-


**Guaranteed Question Practice of 7000+ Questions**

**Classes: Monday to Friday at 10 PM**

**PDF (without annotations) - Provided on my Telegram Channel**

**Topic-wise, Chapter-wise, Unit-wise and Full Mock Questions included**

**Can be followed by all NEET Aspirants**



**Use ARMYLIVE code to join**



**physics\_tarmy**



**a.sane.hurricane**

The component of a vector  $\vec{A}$  along  $y$ -axis will have maximum value if

- (1)  $\vec{A}$  makes an angle of  $30^\circ$  with  $y$ -axis
- (2)  $\vec{A}$  makes an angle of  $60^\circ$  with  $y$ -axis
- (3)  $\vec{A}$  makes an angle of  $0^\circ$  with  $y$ -axis
- (4)  $\vec{A}$  makes an angle of  $90^\circ$  with  $y$ -axis



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

The angle that the vector  $\vec{OA} = 5\hat{i} + 5\hat{j}$  makes with y-axis is

- (1)  $30^\circ$
- (2)  $45^\circ$
- (3)  $60^\circ$
- (4)  $\tan^{-1}\left(\frac{1}{2}\right)$



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

If  $\vec{A} = 2\hat{i} + 3\hat{j}$  and  $\vec{B} = 5\hat{i} + 7\hat{j}$ , then the vector having the magnitude of  $\vec{A}$  and direction of  $\vec{B}$  would be

- (1)  $\left(\frac{74}{13}\right)^{1/2} (2\hat{i} + 3\hat{j})$       (2)  $\sqrt{74} (5\hat{i} + 7\hat{j})$   
(3)  $\left(\frac{13}{74}\right)^{1/2} (5\hat{i} + 7\hat{j})$       (4)  $5\sqrt{13} \hat{i} + 7\sqrt{13} \hat{j}$



The unit vector of a vector  $\vec{A} = 2\hat{i} + 3\hat{j} + 4\hat{k}$  is

(1)  $\frac{9}{\sqrt{29}}(\hat{i} + \hat{j} + \hat{k})$       (2)  $\sqrt{29}(2\hat{i} + 3\hat{j} + 4\hat{k})$

(3)  $\frac{1}{\sqrt{29}}(2\hat{i} + 3\hat{j} + 4\hat{k})$       (4)  $9\sqrt{29}(\hat{i} + \hat{j} + \hat{k})$



If  $\vec{A} = a\hat{i} + 0.5\hat{j} + 0.5\hat{k}$  is unit vector, then value of 'a' would be

(1)  $\frac{1}{\sqrt{2}}$

(2)  $\sqrt{2}$

(3)  $\frac{1}{2}$

(4) 2



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane



A force  $F = (6\hat{i} - 8\hat{j} + 10\hat{k})$  N produces an acceleration of  $1 \text{ m/s}^2$  in a body. The mass of body would be :

- (a) 200 kg                      (b) 20 kg  
(c)  $10\sqrt{2}$  kg              (d)  $6\sqrt{2}$  kg



7777



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

What is the angle between  $\hat{i} + \hat{j} + \hat{k}$  and  $\hat{j}$ ?

- (a)  $0$                       (b)  $45^\circ$   
(c)  $60^\circ$                     (d) None of these



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

13. What is the property of two vectors  $\vec{P}$  and  $\vec{Q}$  if  $\vec{P} + \vec{Q} = \vec{P} - \vec{Q}$  ?

- (a)  $P$  is null vector      (b)  $Q$  is null vector  
(c)  $P$  is proper vector    (d)  $Q$  is proper vector



A child takes 8 steps towards east and 6 steps towards north. If each step is equal to 1 cm, then the magnitude of displacement is :

- (a) 14 m                      (b) 0.1 m  
(c) 10 m                     (d) none of these



The arbitrary number '-2' is multiplied with vector  $\vec{A}$  then :

- (a) the magnitude of vector will be doubled and direction will be same
- (b) the magnitude of vector will be doubled and direction will be opposite
- (c) the magnitude of vector and its direction remain constant
- (d) none of the above



The position vector of a moving particle at time  $t$  is  $\vec{r} = 3\hat{i} + 4t^2\hat{j} - t^3\hat{k}$ . Its displacement during the time interval  $t = 1\text{ s}$  to  $t = 3\text{ s}$  is :

- (a)  $\hat{j} - \hat{k}$                       (b)  $3\hat{i} + 4\hat{j} - \hat{k}$   
(c)  $9\hat{i} + 36\hat{j} - 27\hat{k}$             (d)  $32\hat{j} - 26\hat{k}$



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

A man walks 20 m at an angle of  $60^\circ$  north-east. How far towards east has he travelled ?

- (a) 10 m                      (b) 20 m  
(c)  $20\sqrt{3}$  m              (d)  $\frac{10}{\sqrt{3}}$  m



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

EXAMPLE 3. A particle has a displacement of 12 m towards east and 5 m towards the north and then 6 m vertically upward. Find the magnitude of the sum of these displacements.



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane



EXAMPLE 19. If  $\vec{A} = 3\hat{i} + 4\hat{j}$  and  $\vec{B} = 7\hat{i} + 24\hat{j}$ , find a vector having the same magnitude as  $\vec{B}$  and parallel to  $\vec{A}$ .



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

EXAMPLE 21. One of the rectangular components of a velocity of  $80 \text{ kmh}^{-1}$  is  $40 \text{ kmh}^{-1}$ . Find the other component.



Use ARMYLIVE code to join



physics\_tarmy



a.sane.hurricane

4. Find the value of  $\lambda$  in the unit vector

$$0.4 \hat{i} + 0.8 \hat{j} + \lambda \hat{k}.$$



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

A velocity of  $10 \text{ ms}^{-1}$  has its Y-component  $5\sqrt{2} \text{ ms}^{-1}$ .  
Calculate its X-component.



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

# One Stop for Physics Practice for **NEET**


Monday - Friday ; 10 PM

**1 Series = Questions from Top Books**

## Vectors

### Vectorial Addition/ Subtraction

**3**



**Tamanna Chaudhary**  
(Physics Expert)



**physics\_tarmy**



**a.sane.hurricane**



# Tamanna Chaudhary

Expert in NEET UG

One of the most followed Educators, Tamanna, won the 2020 Unacademy People's Choice Award. Her student secured 99.56 percentile in Physics.

**98M** Watch mins

**2M** Watch mins (last 30 days)

**254K** Followers

**14K** Dedications



# 7777



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane



# One Stop for Physics Practice for **NEET**

**1 Series = Questions from Top Books**



Tamanna Chaudhary ✓  
117K subscribers

## Details about the series-

**Guaranteed Question Practice of 7000+ Questions**

**Classes: Monday to Friday at 10 PM**

**PDF (without annotations) - Provided on my Telegram Channel**

**Topic-wise, Chapter-wise, Unit-wise and Full Mock Questions included**

**Can be followed by all NEET Aspirants**

NEWTON'S APPLE



**Use ARMYLIVE code to join**



**physics\_tcarmy**



**a.sane.hurricane**

# One Stop for Physics Practice for **NEET**

**1 Series = Questions from Top Books**



Tamanna Chaudhary ✓  
117K subscribers

## Books Covered in this Session-



**HC Verma**



**SL Arora**



**DC Pandey**



**TC Selected**



NEWTON'S APPLE



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane



If the sum of two unit vectors is a unit vector, then magnitude of difference is

A)  $\sqrt{2}$

B)  $\sqrt{3}$

C)  $1/\sqrt{2}$

D)  $\sqrt{5}$



7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

$$\vec{A} = 2\hat{i} + \hat{j}, B = 3\hat{j} - \hat{k} \text{ and } \vec{C} = 6\hat{i} - 2\hat{k}.$$

Value of

$$\vec{A} - 2\vec{B} + 3\vec{C} \text{ would be}$$

A)  $20\hat{i} + 5\hat{j} + 4\hat{k}$

B)  $20\hat{i} - 5\hat{j} - 4\hat{k}$

C)  $4\hat{i} + 5\hat{j} + 20\hat{k}$

D)  $5\hat{i} + 4\hat{j} + 10\hat{k}$



For the resultant of the two vectors to be maximum, what must be the angle between them

- A)  $0^\circ$
- B)  $60^\circ$
- C)  $90^\circ$
- D)  $180^\circ$



7777



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

Let the angle between two nonzero vectors  $\vec{A}$  and  $\vec{B}$  be  $120^\circ$  and resultant be  $\vec{C}$

- A)  $|\vec{C}|$  must be equal to  $|\vec{A} - \vec{B}|$
- B)  $|\vec{C}|$  must be less than  $|\vec{A} - \vec{B}|$
- C)  $|\vec{C}|$  must be greater than  $|\vec{A} - \vec{B}|$
- D)  $|\vec{C}|$  may be equal to  $|\vec{A} - \vec{B}|$

NEWTON'S APPLE



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

What vector must be added to the two vectors  $\hat{i} - 2\hat{j} + 2\hat{k}$  and  $2\hat{i} + \hat{j} - \hat{k}$ , so that the resultant may be a unit vector along x-axis

- A)  $2\hat{i} + \hat{j} - \hat{k}$
- B)  $-2\hat{i} + \hat{j} - \hat{k}$
- C)  $2\hat{i} - \hat{j} + \hat{k}$
- D)  $-2\hat{i} - \hat{j} - \hat{k}$



Assertion : Vector addition of two vectors  $\vec{A}$  and  $\vec{B}$  is commutative.  
Reason :  $\vec{A} + \vec{B} = \vec{B} + \vec{A}$

- A) If both assertion and reason are true and the reason is the correct explanation of the assertion.
- B) If both assertion and reason are true but reason is not the correct explanation of the assertion.
- C) If assertion is true but reason is false.
- D) If the assertion and reason both are false.
- E) If assertion is false but reason is true.

NEWTON'S APPLE



Use ARMYLIVE code to join




physics\_tcarmy



a.sane.hurricane

The magnitudes of vectors  $\vec{OA}$ ,  $\vec{OB}$  and  $\vec{OC}$  in figure (2-W3) are equal. Find the direction of  $\vec{OA} + \vec{OB} - \vec{OC}$ .




6. Two vectors have magnitudes 3 unit and 4 unit respectively. What should be the angle between them if the magnitude of the resultant is (a) 1 unit, (b) 5 unit and (c) 7 unit.





5. Refer to figure (2-E1). Find (a) the magnitude, (b)  $x$  and  $y$  components and (c) the angle with the  $X$ -axis of the resultant of  $\vec{OA}$ ,  $\vec{BC}$  and  $\vec{DE}$ .



2. Let  $\vec{A}$  and  $\vec{B}$  be the two vectors of magnitude 10 unit each. If they are inclined to the  $X$ -axis at angles  $30^\circ$  and  $60^\circ$  respectively, find the resultant.



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

7. Can you add two vectors representing physical quantities having different dimensions? Can you multiply two vectors representing physical quantities having different dimensions?



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

2. Find the resultant of the three vectors shown in figure (2-W1).




Figure 2-W1

NEWTON'S APPLE



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

When the angle between two vectors of equal magnitude is  $2\pi / 3$ , prove that the magnitude of the resultant is equal to either.



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

At what angle do the two forces  $(P + Q)$  and  $(P - Q)$  act so that the resultant is  $\sqrt{3P^2 + Q^2}$ . (Ans.  $60^\circ$ )



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

Find the angle between two vectors  $\vec{P}$  and  $\vec{Q}$  if resultant of the vectors is given by  $R^2 = P^2 + Q^2$ .



7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

Given three coplanar vectors  $\vec{a} = 4\hat{i} - \hat{j}$ ,  
 $\vec{b} = -3\hat{i} + 2\hat{j}$  and  $\vec{c} = -3\hat{j}$ . Find the magnitude of  
the sum of the three vectors. [Ans.  $\sqrt{5}$ ]





Problem 14. If  $\vec{a} + \vec{b} = \vec{c}$  and  $|\vec{a}| + |\vec{b}| = |\vec{c}|$ , what can we say about the direction of these vectors?



Use ARMYLIVE code to join



physics\_tcarmy



a.sane.hurricane

Problem 8. Under what condition does the equality :

$$|\vec{A} + \vec{B}| = |\vec{A} - \vec{B}| \text{ hold good ?}$$



NEWTON'S APPLE

7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane


The maximum resultant of two vectors is 26 unit and minimum resultant is 16 unit, then the magnitude of each vector is :

- (a) 21, 5                      (b) 13, 13  
(c) 20, 6                      (d) none of these



Two vectors  $\vec{P}$  and  $\vec{Q}$  have equal magnitude of 10 unit. They are oriented as shown in figure. The resultant of these vector is :

- (a) 10 unit
- (b)  $10\sqrt{2}$  unit
- (c) 12 unit
- (d) none of the above



7777



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane

A cyclist is moving on a circular path with constant speed. What is the change in its velocity after it has described an angle of  $30^\circ$ ?

- (a)  $v\sqrt{2}$  (b)  $v(0.3\sqrt{3})$   
(c)  $v\sqrt{3}$  (d) None of these




If two forces of equal magnitude 4 units acting at a point and the angle between them is  $120^\circ$  then the magnitude and direction of the sum of the two vectors are :

- (a) 4,  $\theta = \tan^{-1}(1.73)$       (b) 4,  $\theta = \tan^{-1}(0.73)$   
(c) 2,  $\theta = \tan^{-1}(1.73)$       (d) 6,  $\theta = \tan^{-1}(0.73)$



The angle between  $\vec{A}$  and the resultant of  $(\vec{A} + \vec{B})$  and  $(\vec{A} - \vec{B})$  will be :

- (a)  $0^\circ$                       (b)  $\tan^{-1}\left(\frac{A}{B}\right)$   
(c)  $\tan^{-1}\left(\frac{B}{A}\right)$               (d)  $\tan^{-1}\left(\frac{A-B}{A+B}\right)$



Obtain the magnitude and direction cosines of vector  $(\vec{A} - \vec{B})$ , if  $\vec{A} = 2\hat{i} + 3\hat{j} + \hat{k}$ ,  $\vec{B} = 2\hat{i} + 2\hat{j} + 3\hat{k}$ :

(a)  $0, \frac{1}{\sqrt{5}}, \frac{-2}{\sqrt{5}}$

(b)  $0, \frac{2}{\sqrt{5}}, \frac{1}{\sqrt{5}}$

(c)  $0, 0, \frac{1}{\sqrt{5}}$

(d) none of these



Use ARMYLIVE code to join



physics\_tcmrmy



a.sane.hurricane