

# 15. Treatments in many time periods. The problem.

Ian Lundberg  
Cornell Info 6751: Causal Inference in Observational Settings  
Fall 2022

13 Oct 2022

# Learning goals for today

At the end of class, you will be able to:

1. Present treatments that unfold over time in DAGs
2. Recognize the difficulties of treatment-induced confounding

## Treatments in many time periods: Motivation

Suppose you teach second grade

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle
- ▶ In December, you hope every child can read a picture book

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle
- ▶ In December, you hope every child can read a picture book

Task: Draw this in a DAG

# Treatments in many time periods: Motivation

Child cannot  
sound out  
words in  
September




# Treatments in many time periods: Motivation

Assign parent  
volunteer to  
read with child  
in September




Child cannot  
sound out  
words in  
September


# Treatments in many time periods: Motivation




# Treatments in many time periods: Motivation




# Treatments in many time periods: Motivation



Treatments in many time periods: A general problem




## Treatments in many time periods: A general problem



This causal structure occurs

- ▶ when a policymaker targets treatment  $A_k$  at time  $k$  given confounders  $L_k$  measured at that time
- ▶ in observational settings where treatments unfold over time

# Treatments in many time periods: A general problem




This causal structure occurs

- ▶ when a policymaker targets treatment  $A_k$  at time  $k$  given confounders  $L_k$  measured at that time
- ▶ in observational settings where treatments unfold over time

Goal: Study the outcome  $Y$  would be realized on average if  $A_0, \dots, A_k$  are set to the values  $a_0, \dots, a_k$ .


Treatments in many time periods:  
The curse of dimensionality



Each  $A_k$  is binary. How many potential outcomes are there?



# Treatments in many time periods: The curse of dimensionality



Each  $A_k$  is binary. How many potential outcomes are there?

- ▶  $\bar{a} = (0, 0)$ : No reading with a parent
- ▶  $\bar{a} = (1, 0)$ : Read in September, not October
- ▶  $\bar{a} = (0, 1)$ : Read in October, not September
- ▶  $\bar{a} = (1, 1)$ : Always read with a parent

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

There are then  $2^9 = 512$  potential outcomes  $Y^{a_0, \dots, a_8}$  for each child

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

There are then  $2^9 = 512$  potential outcomes  $Y^{a_0, \dots, a_8}$  for each child

This is why we focus on **treatment strategies**

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

We would then assign treatment  $A_k = g(L_k)$



# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

## Example:

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

We would then assign treatment  $A_k = g(L_k)$

This involves **many treatments**, but only **one strategy**.

## Treatment strategy: Exercise

Use math to define the following strategy:

Assign a parent volunteer to read with a child  $A_k = 1$  if and only if the child struggles sounding out words  $L_k = 0$  and the child did not receive this support last month  $A_{k-1} = 0$

## Treatment strategy: Exercise

Use math to define the following strategy:

Assign a parent volunteer to read with a child  $A_k = 1$  if and only if the child struggles sounding out words  $L_k = 0$  and the child did not receive this support last month  $A_{k-1} = 0$

$$g(L_k, A_{k-1}) = \mathbb{I}(L_k = 0, A_{k-1} = 0)$$

Treatment strategy: Static and dynamic

## Treatment strategy: Static and dynamic

A **static** strategy assigns treatments in advance

- Example: Always treat.  $g() = 1$

# Treatment strategy: Static and dynamic

A **static** strategy assigns treatments in advance


- ▶ Example: Always treat.  $g() = 1$

A **dynamic** strategy assigns treatments as a function of the changing values of confounding variables

- ▶ Example: Treat if has difficulty sounding out words.  
 $g(L_k) = \mathbb{I}(L_k = 0)$


Identification: The adjustment set

Identification: The adjustment set






## Identification: The adjustment set




1. What is the sufficient adjustment set to identify

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$

## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set




## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set

## Identification: The adjustment set




**(2) has no solution!**

1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?


$U$  is unobserved and cannot be part of the adjustment set

Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed


## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- What happens if you adjust for  $L_1$ ?


## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$

## Identification: The adjustment set




A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?




## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?
  - ▶ A backdoor path remains:  $A_1 \leftarrow L_1 \rightarrow Y$

## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?
  - ▶ A backdoor path remains:  $A_1 \leftarrow L_1 \rightarrow Y$

Next class: How to correctly adjust for treatment-induced confounding

# Learning goals for today

At the end of class, you will be able to:

1. Present treatments that unfold over time in DAGs
2. Recognize the difficulties of treatment-induced confounding

Let me know what you are thinking

[tinyurl.com/CausalQuestions](https://tinyurl.com/CausalQuestions)

Office hours TTh 11am-12pm and at  
[calendly.com/ianlundberg/office-hours](https://calendly.com/ianlundberg/office-hours)  
Come say hi!