

## 8. Parametric g-formula: Categorical treatments

Ian Lundberg

Cornell Info 6751: Causal Inference in Observational Settings  
Fall 2022

15 Sep 2022

# Learning goals for today

At the end of class, you will be able to:

1. Estimate causal effects by outcome modeling with the parametric g-formula
2. See how this generalizes a common use of regression

## Recap: Nonparametric identification

# Recap: Nonparametric identification

Three key assumptions:

Consistency	$Y_i = Y_i^{A_i}$
Exchangeability	$A \perp\!\!\!\perp \{Y^a\} \mid \vec{L}$
Positivity	$P(A = a \mid \vec{L} = \vec{\ell}) > 0$

# Recap: Nonparametric identification

Three key assumptions:

Consistency	$Y_i = Y_i^{A_i}$
Exchangeability	$A \perp\!\!\!\perp \{Y^a\} \mid \vec{L}$
Positivity	$P(A = a \mid \vec{L} = \vec{\ell}) > 0$

These assumptions yield **nonparametric identification**:  
a consistent estimator exists using observable sample means

Recap: Nonparametric identification in a proof

## Recap: Nonparametric identification in a proof

$$E(Y^a)$$

## Recap: Nonparametric identification in a proof

$$E(Y^a) = \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell})$$

rules of probability



## Recap: Nonparametric identification in a proof

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}) && \text{rules of probability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}, A = a) && \text{exchangeability} \end{aligned}$$

## Recap: Nonparametric identification in a proof

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}) && \text{rules of probability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}, A = a) && \text{exchangeability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) && \text{consistency} \end{aligned}$$

## Recap: Nonparametric identification in a proof

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}) && \text{rules of probability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}, A = a) && \text{exchangeability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) && \text{consistency} \end{aligned}$$

Positivity ensures  $E(Y \mid \vec{L} = \vec{\ell}, A = a)$  can be estimated from data.

## Recap: Nonparametric identification in a proof

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}) && \text{rules of probability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y^a \mid \vec{L} = \vec{\ell}, A = a) && \text{exchangeability} \\ &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) && \text{consistency} \end{aligned}$$

Positivity ensures  $E(Y \mid \vec{L} = \vec{\ell}, A = a)$  can be estimated from data.

**What we gained:** In an infinite sample, we can estimate causal effects by taking means and aggregating!

## Recap: Nonparametric identification with two $E$ s

For simplicity, rewrite the previous result with two expectations

## Recap: Nonparametric identification with two Es

For simplicity, rewrite the previous result with two expectations

$$E(Y^a) = \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a)$$

## Recap: Nonparametric identification with two Es

For simplicity, rewrite the previous result with two expectations

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) \\ &= E\left(E(Y \mid \vec{L}, A = a)\right) \end{aligned}$$

## Recap: Nonparametric identification with two Es

For simplicity, rewrite the previous result with two expectations

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) \\ &= E\left(E(Y \mid \vec{L}, A = a)\right) \end{aligned}$$

There are two pieces



## Recap: Nonparametric identification with two Es

For simplicity, rewrite the previous result with two expectations

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) \\ &= E\left(E(Y \mid \vec{L}, A = a)\right) \end{aligned}$$

There are two pieces

- Inner expectation: Within groups
  - Mean over units within each stratum of confounders

## Recap: Nonparametric identification with two Es

For simplicity, rewrite the previous result with two expectations

$$\begin{aligned} E(Y^a) &= \sum_{\vec{\ell}} P(\vec{L} = \vec{\ell}) E(Y \mid \vec{L} = \vec{\ell}, A = a) \\ &= E\left(E(Y \mid \vec{L}, A = a)\right) \end{aligned}$$

There are two pieces

- ▶ Inner expectation: Within groups
  - ▶ Mean over units within each stratum of confounders
- ▶ Outer expectation: Across groups
  - ▶ Mean of that over the population distribution of  $\vec{L}$

Recap: Nonparametric identification. An estimator

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

Recap: Nonparametric identification. An estimator

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

This points toward a **non-parametric plug-in estimator**:

Recap: Nonparametric identification. An estimator

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

This points toward a **non-parametric plug-in estimator**:

- Inner expectation: Sample mean in each subgroup

## Recap: Nonparametric identification. An estimator

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

This points toward a **non-parametric plug-in estimator**:

- ▶ Inner expectation: Sample mean in each subgroup
- ▶ Outer expectation: Mean over full sample

## Recap: Nonparametric identification. An estimator

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

This points toward a **non-parametric plug-in estimator**:

- ▶ Inner expectation: Sample mean in each subgroup
- ▶ Outer expectation: Mean over full sample

Resulting estimator:

$$\hat{E}(Y^a) = \frac{1}{n} \sum_{i=1}^n \hat{E}(Y \mid \vec{L} = \vec{\ell}_i, A = a)$$

## Recap: Nonparametric identification. An estimator

Putting this in words:

$$\hat{E}(Y^a) = \frac{1}{n} \sum_{i=1}^n \hat{E}(Y \mid \vec{L} = \vec{\ell}_i, A = a)$$

- ▶ Sample average over units  $i$
- ▶ For each unit, take the sample average outcome among
  - ▶ Units with their covariate values  $\vec{\ell}_i$
  - ▶ But who have the relevant treatment value  $A = a$

It is all just sample means, aggregated a certain way.



Why move to **parametric** estimation?

# Why move to **parametric** estimation?

Some reasons for a parametric model:

# Why move to **parametric** estimation?

Some reasons for a parametric model:

# Why move to **parametric** estimation?

Some reasons for a parametric model:

1. **Empty cells:** Despite positivity in the population, some sample strata have no units with  $A = a$ . Nonparametric estimation is impossible

# Why move to **parametric** estimation?

Some reasons for a parametric model:

1. **Empty cells:** Despite positivity in the population, some sample strata have no units with  $A = a$ . Nonparametric estimation is impossible
2. **Continuous confounders:** If any variables in  $\vec{L}$  are continuous, then each stratum of  $\vec{L}$  contains only one unit. Empty cells are inevitable

# Why move to **parametric** estimation?

Some reasons for a parametric model:

1. **Empty cells:** Despite positivity in the population, some sample strata have no units with  $A = a$ . Nonparametric estimation is impossible
2. **Continuous confounders:** If any variables in  $\vec{L}$  are continuous, then each stratum of  $\vec{L}$  contains only one unit. Empty cells are inevitable
3. **Estimation variance:** Even if you can do nonparametric estimation, it involves estimating numerous means. This may be a high-variance approach, resulting in extensive statistical uncertainty

The parametric g-formula

## The parametric g-formula

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$



# The parametric g-formula

Nonparametric identification yielded this estimand:

$$E(Y^a) = E\left(E(Y \mid \vec{L}, A = a)\right)$$

What if we estimated  $E(Y \mid \vec{L} = \vec{\ell}, A = a)$  with a regression model?

The parametric g-formula: Simple case

$$L \longrightarrow A \longrightarrow Y$$


The parametric g-formula: Simple case

$$L \longrightarrow A \longrightarrow Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$


## The parametric g-formula: Simple case

$$L \longrightarrow A \longrightarrow Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

## The parametric g-formula: Simple case

$$L \xrightarrow{\quad} A \xrightarrow{\quad} Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

# The parametric g-formula: Simple case

$$L \longrightarrow A \longrightarrow Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$


Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

Estimator in words:

# The parametric g-formula: Simple case

$$L \longrightarrow A \longrightarrow Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$


Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

Estimator in words:

1. Estimate the regression model

# The parametric g-formula: Simple case

$$L \xrightarrow{\quad} A \xrightarrow{\quad} Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

Estimator in words:

1. Estimate the regression model
2. Change all treatment values to 1



# The parametric g-formula: Simple case

$$L \rightarrow A \rightarrow Y$$


Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

Estimator in words:

1. Estimate the regression model
2. Change all treatment values to 1
3. Predict for everyone

# The parametric g-formula: Simple case

$$L \xrightarrow{\quad} A \xrightarrow{\quad} Y$$

Nonparametric identification  $E(Y^a) = E(E(Y \mid L, A = a))$

Parametric model  $E(Y \mid L, A) = \alpha + \gamma L + \beta A$

Estimator for the potential outcome under treatment ( $A = 1$ ):

$$\hat{E}(Y^1) = \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right)$$

Estimator in words:

1. Estimate the regression model
2. Change all treatment values to 1
3. Predict for everyone
4. Take the sample mean

The parametric g-formula: Connection to  $\hat{\beta}$

The parametric g-formula: Connection to  $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\hat{E}(Y^1) - \hat{E}(Y^0)$$

## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\hat{E}(Y^1) - \hat{E}(Y^0) = \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right) \right)$$

## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 \right) \right)\end{aligned}$$

## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 \right) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \hat{\beta}\end{aligned}$$



## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 \right) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \hat{\beta} \\ &= \hat{\beta}\end{aligned}$$

## The parametric g-formula: Connection to $\hat{\beta}$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 \right) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \hat{\beta} \\ &= \hat{\beta}\end{aligned}$$

With OLS, the parametric g-formula collapses on the coefficient.

The parametric g-formula is more general

The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\hat{E}(Y^1) - \hat{E}(Y^0)$$

The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\hat{E}(Y^1) - \hat{E}(Y^0) = \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 + \hat{\eta} \times 1 \times \ell_i \right) \right)$$

## The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) = & \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 + \hat{\eta} \times 1 \times \ell_i \right) \right) \\ & - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 + \hat{\eta} \times 0 \times \ell_i \right) \right)\end{aligned}$$



# The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 + \hat{\eta} \times 1 \times \ell_i \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 + \hat{\eta} \times 0 \times \ell_i \right) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \left( \hat{\beta} + \hat{\eta} \ell_i \right)\end{aligned}$$

# The parametric g-formula is more general

Suppose we add the  $A \times L$  interaction to the model.

$$E(Y \mid A, L) = \alpha + \gamma L + \beta A + \eta AL$$

Estimator for the effect  $E(Y^1) - E(Y^0)$ :

$$\begin{aligned}\hat{E}(Y^1) - \hat{E}(Y^0) &= \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 1 + \hat{\eta} \times 1 \times \ell_i \right) \right) \\ &\quad - \left( \frac{1}{n} \sum_{i=1}^n \left( \hat{\alpha} + \hat{\gamma} \ell_i + \hat{\beta} \times 0 + \hat{\eta} \times 0 \times \ell_i \right) \right) \\ &= \frac{1}{n} \sum_{i=1}^n \left( \hat{\beta} + \hat{\eta} \ell_i \right)\end{aligned}$$

The g-formula no longer collapses to a coefficient!

The parametric g-formula: Word of warning

## The parametric g-formula: Word of warning

The parametric g-formula does not replace any assumptions.

# The parametric g-formula: Word of warning

The parametric g-formula does not replace any assumptions.

You need:

## Identification Assumptions (causal)

Consistency  $Y_i = Y_i^{A_i}$

Exchangeability  $A \perp\!\!\!\perp \{Y^a\} \mid \vec{L}$

Positivity  $P(A = a \mid \vec{L} = \vec{\ell}) > 0$

## Estimation Assumptions (statistical)

Estimation  $E(Y \mid A, \vec{L})$  follows the assumed model

# The parametric g-formula: Word of warning

The parametric g-formula does not replace any assumptions.

You need:

## Identification Assumptions (causal)

Consistency  $Y_i = Y_i^{A_i}$

Exchangeability  $A \perp\!\!\!\perp \{Y^a\} \mid \vec{L}$

Positivity  $P(A = a \mid \vec{L} = \vec{\ell}) > 0$

## Estimation Assumptions (statistical)

Estimation  $E(Y \mid A, \vec{L})$  follows the assumed model

To use a parametric model, we **added** the bottom assumption.

# The parametric g-formula: Word of warning

The parametric g-formula does not replace any assumptions.

You need:

## Identification Assumptions (causal)

Consistency  $Y_i = Y_i^{A_i}$

Exchangeability  $A \perp\!\!\!\perp \{Y^a\} \mid \vec{L}$

Positivity  $P(A = a \mid \vec{L} = \vec{\ell}) > 0$

## Estimation Assumptions (statistical)

Estimation  $E(Y \mid A, \vec{L})$  follows the assumed model

To use a parametric model, we **added** the bottom assumption.

- But testable! Diagnostics and out-of-sample performance

## The parametric g-formula: Summary



# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- Allows you to use a model for  $E(Y \mid A, \vec{L})$

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics
- ▶ The estimator is operationalized by prediction:

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics
- ▶ The estimator is operationalized by prediction:
  - ▶ Estimate the regression

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics
- ▶ The estimator is operationalized by prediction:
  - ▶ Estimate the regression
  - ▶ Modify the treatment

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics
- ▶ The estimator is operationalized by prediction:
  - ▶ Estimate the regression
  - ▶ Modify the treatment
  - ▶ Make new predictions

# The parametric g-formula: Summary

The parametric g-formula is how we answer  
nonparametric causal questions  
with  
parametric outcome modeling tools

- ▶ Allows you to use a model for  $E(Y \mid A, \vec{L})$
- ▶ The model is exactly as in standard non-causal statistics
- ▶ The estimator is operationalized by prediction:
  - ▶ Estimate the regression
  - ▶ Modify the treatment
  - ▶ Make new predictions
  - ▶ Average over the sample



# Learning goals for today

At the end of class, you will be able to:

1. Estimate causal effects by outcome modeling with the parametric g-formula
2. See how this generalizes a common use of regression

Let me know what you are thinking

[tinyurl.com/CausalQuestions](https://tinyurl.com/CausalQuestions)

Office hours TTh 11am-12pm and at  
[calendly.com/ianlundberg/office-hours](https://calendly.com/ianlundberg/office-hours)  
Come say hi!