

## 15. Treatments in many time periods

Ian Lundberg

Cornell Info 6751: Causal Inference in Observational Settings  
Fall 2022

13 Oct 2022

# Learning goals for today

At the end of class, you will be able to:

1. Present treatments that unfold over time in DAGs
2. Reason about the sequential ignorability assumption
3. Apply inverse probability weighting to treatments over time

## Treatments in many time periods: Motivation

Suppose you teach second grade

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle
- ▶ In December, you hope every child can read a picture book

# Treatments in many time periods: Motivation

Suppose you teach second grade

- ▶ Every month, you assess children's ability to sound out words
- ▶ You assign parent volunteers to read with those who struggle
- ▶ In December, you hope every child can read a picture book

Task: Draw this in a DAG

# Treatments in many time periods: Motivation

Child cannot  
sound out  
words in  
September




# Treatments in many time periods: Motivation

Assign parent  
volunteer to  
read with child  
in September




Child cannot  
sound out  
words in  
September


# Treatments in many time periods: Motivation




# Treatments in many time periods: Motivation




## Treatments in many time periods: Motivation



Treatments in many time periods: A general problem




## Treatments in many time periods: A general problem



This causal structure occurs

- ▶ when a policymaker targets treatment  $A_k$  at time  $k$  given confounders  $L_k$  measured at that time
- ▶ in observational settings where treatments unfold over time

# Treatments in many time periods: A general problem




This causal structure occurs

- ▶ when a policymaker targets treatment  $A_k$  at time  $k$  given confounders  $L_k$  measured at that time
- ▶ in observational settings where treatments unfold over time

Goal: Study the outcome  $Y$  would be realized on average if  $A_0, \dots, A_k$  are set to the values  $a_0, \dots, a_k$ .


Treatments in many time periods:  
The curse of dimensionality



Each  $A_k$  is binary. How many potential outcomes are there?



# Treatments in many time periods: The curse of dimensionality



Each  $A_k$  is binary. How many potential outcomes are there?

- ▶  $\bar{a} = (0, 0)$ : No reading with a parent
- ▶  $\bar{a} = (1, 0)$ : Read in September, not October
- ▶  $\bar{a} = (0, 1)$ : Read in October, not September
- ▶  $\bar{a} = (1, 1)$ : Always read with a parent

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

There are then  $2^9 = 512$  potential outcomes  $Y^{a_0, \dots, a_8}$  for each child

# Treatments in many time periods: The curse of dimensionality

Suppose the teacher can assign (or not) a parent volunteer to read with a child in each of 9 months in the school year

$$A_0, \dots, A_8$$

There are then  $2^9 = 512$  potential outcomes  $Y^{a_0, \dots, a_8}$  for each child

This is why we focus on **treatment strategies**

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

**Example:**

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

We would then assign treatment  $A_k = g(L_k)$



# Treatment strategy

A treatment strategy is a counterfactual policy rule  $g()$  for assigning the treatment

## Example:

Assign a parent volunteer to read with a child whenever the child struggles sounding out words

$$g(L_k) = \mathbb{I}(L_k = 0)$$

We would then assign treatment  $A_k = g(L_k)$

This involves **many treatments**, but only **one strategy**.

## Treatment strategy: Exercise

Use math to define the following strategy:

Assign a parent volunteer to read with a child  $A_k = 1$  if and only if the child struggles sounding out words  $L_k = 0$  and the child did not receive this support last month  $A_{k-1} = 0$

## Treatment strategy: Exercise

Use math to define the following strategy:

Assign a parent volunteer to read with a child  $A_k = 1$  if and only if the child struggles sounding out words  $L_k = 0$  and the child did not receive this support last month  $A_{k-1} = 0$

$$g(L_k, A_{k-1}) = \mathbb{I}(L_k = 0, A_{k-1} = 0)$$

Treatment strategy: Static and dynamic

## Treatment strategy: Static and dynamic

A **static** strategy assigns treatments in advance

- ▶ Example: Always treat.  $g() = 1$

# Treatment strategy: Static and dynamic

A **static** strategy assigns treatments in advance


- ▶ Example: Always treat.  $g() = 1$

A **dynamic** strategy assigns treatments as a function of the changing values of confounding variables

- ▶ Example: Treat if has difficulty sounding out words.  
 $g(L_k) = \mathbb{I}(L_k = 0)$


Identification: The adjustment set

Identification: The adjustment set






## Identification: The adjustment set




1. What is the sufficient adjustment set to identify

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$

## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set

## Identification: The adjustment set




1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set




## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set


## Identification: The adjustment set



1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?

$U$  is unobserved and cannot be part of the adjustment set

## Identification: The adjustment set




**(2) has no solution!**

1. What is the sufficient adjustment set to identify
  - a) The total effect of  $A_0$  on  $Y$
  - b) The total effect of  $A_1$  on  $Y$
2. We want to identify the effect of  $\bar{A} = (A_0, A_1)$  on  $Y$   
Can we jointly block all backdoor paths between  $\bar{A}$  and  $Y$ ?


$U$  is unobserved and cannot be part of the adjustment set

Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed


## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- What happens if you adjust for  $L_1$ ?


## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- What happens if you adjust for  $L_1$ ?
  - You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$

## Identification: The adjustment set




A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?




## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?
  - ▶ A backdoor path remains:  $A_1 \leftarrow L_1 \rightarrow Y$

## Identification: The adjustment set



A joint adjustment set for  $\bar{A}$  is doomed

- ▶ What happens if you adjust for  $L_1$ ?
  - ▶ You block a causal path:  $A_0 \rightarrow \boxed{L_1} \rightarrow Y$
  - ▶ You open a backdoor path:  $A_0 \rightarrow \boxed{L_1} \leftarrow U \rightarrow Y$
- ▶ What happens if you don't adjust for  $L_1$ ?
  - ▶ A backdoor path remains:  $A_1 \leftarrow L_1 \rightarrow Y$

To proceed, we need a different adjustment set in each time period

# Notation

- ▶  $\bar{A}_k = (A_0, A_1, \dots, A_k)$  treatments up to time  $k$
- ▶  $\bar{L}_k = (L_0, L_1, \dots, L_k)$  confounders up to time  $k$
- ▶  $g()$  treatment strategy
- ▶  $Y^g$  potential outcome under that strategy

Identification: Sequential ignorability

# Identification: Sequential ignorability


$$\begin{array}{ccccccc}
 \text{Potential} & & & & & & \\
 \text{outcome} & & & & & & \\
 \text{under} & \text{is} & & & & & \\
 \text{assignment} & \text{independent} & \text{treatment} & \text{given} & \text{treatments up to } k-1 \text{ followed} & \text{rule } g & \text{and confounders} \\
 \text{rule } g & \text{of} & \text{at time } k & & & & \text{up to time } k \\
 \\
 Y^g & \perp\!\!\!\perp & A_k & | & \bar{A}_{k-1} = g(\bar{A}_{k-2}, \bar{L}_{k-1}), & & \bar{L}_k
 \end{array}$$

for all assignment rules  $g$  and time periods  $k = 1, \dots, K$

# Identification: Sequential ignorability

$$\begin{array}{ccccccc}
 \text{Potential} & & \text{is} & & & & \\
 \text{outcome} & & \text{independent} & & \text{treatment} & & \\
 \text{under} & & \text{of} & & \text{at time } k & & \\
 \text{assignment} & & & & \text{given} & & \\
 \text{rule } g & & & & & & \\
 Y^g & \perp\!\!\!\perp & A_k & | & \bar{A}_{k-1} = g(\bar{A}_{k-2}, \bar{L}_{k-1}), & \bar{L}_k & \\
 & & & & \text{treatments up to } k-1 \text{ followed rule } g & \text{and confounders} & \\
 & & & & & \text{up to time } k & 
 \end{array}$$


for all assignment rules  $g$  and time periods  $k = 1, \dots, K$



# Identification: Sequential ignorability

$$\begin{array}{ccccccc}
 \text{Potential} & & \text{is} & & & & \\
 \text{outcome} & & \text{independent} & & \text{treatment} & & \\
 \text{under} & & \text{of} & & \text{at time } k & & \\
 \text{assignment} & & & & \text{given} & & \\
 \text{rule } g & & & & & & \\
 Y^g & \perp\!\!\!\perp & A_k & | & \bar{A}_{k-1} = g(\bar{A}_{k-2}, \bar{L}_{k-1}), & \bar{L}_k & \\
 & & & & \text{treatments up to } k-1 \text{ followed rule } g & \text{and confounders} & \\
 & & & & & \text{up to time } k & 
 \end{array}$$


for all assignment rules  $g$  and time periods  $k = 1, \dots, K$



# Identification: Sequential ignorability

$$\begin{array}{ccccccc}
 \text{Potential} & & \text{is} & & & & \\
 \text{outcome} & & \text{independent} & & \text{treatment} & & \text{and confounders} \\
 \text{under} & & \text{of} & & \text{at time } k & & \text{up to time } k \\
 \text{assignment} & & & & \text{given} & & \\
 \text{rule } g & & & & & & \\
 Y^g & \perp\!\!\!\perp & A_k & | & \bar{A}_{k-1} = g(\bar{A}_{k-2}, \bar{L}_{k-1}), & \bar{L}_k & \\
 & & & & \text{treatments up to } k-1 \text{ followed rule } g & & 
 \end{array}$$

for all assignment rules  $g$  and time periods  $k = 1, \dots, K$






# Identification: Sequential ignorability

$$\begin{array}{ccccccc}
 \begin{array}{c} \text{Potential} \\ \text{outcome} \\ \text{under} \\ \text{assignment} \\ \text{rule } g \end{array} & \begin{array}{c} \text{is} \\ \text{independent} \\ \text{of} \end{array} & \begin{array}{c} \text{treatment} \\ \text{at time } k \end{array} & \begin{array}{c} \text{given} \\ \\ \end{array} & \begin{array}{c} \text{treatments up to } k-1 \text{ followed rule } g \\ \\ \end{array} & \begin{array}{c} \text{and confounders} \\ \text{up to time } k \end{array} \\
 Y^g & \perp\!\!\!\perp & A_k & | & \bar{A}_{k-1} = g(\bar{A}_{k-2}, \bar{L}_{k-1}), & \bar{L}_k
 \end{array}$$

for all assignment rules  $g$  and time periods  $k = 1, \dots, K$




## Estimation: Two strategies

1. Inverse probability weighting (+ marginal structural models)
2. Structural nested mean models (coming next class)

Inverse probability weighting: DAG motivation


# Inverse probability weighting: DAG motivation

We observe data from this model




# Inverse probability weighting: DAG motivation

We observe data from this model




We want this




# Inverse probability weighting: DAG motivation

We observe data from this model



We want this



1. How would you weight to estimate the effect of  $A_0$ ?
2. How would you weight to estimate the effect of  $A_1$ ?

We will combine these

## Inverse probability weighting

In time 0, define an inverse probability of treatment weight

$$W^{A_0} = \frac{1}{P(A_0 \mid L_0)}$$

such that  $A_0$  does not depend on  $L_0$  after weighting

# Inverse probability weighting

In time 0, define an inverse probability of treatment weight

$$W^{A_0} = \frac{1}{P(A_0 \mid L_0)}$$

such that  $A_0$  does not depend on  $L_0$  after weighting

In time 1, do it again

$$W^{A_1} = \frac{1}{P(A_1 \mid \bar{A}_{k-1}, \bar{L}_1)}$$



# Inverse probability weighting

In time 0, define an inverse probability of treatment weight

$$W^{A_0} = \frac{1}{P(A_0 | L_0)}$$

such that  $A_0$  does not depend on  $L_0$  after weighting

In time 1, do it again

$$W^{A_1} = \frac{1}{P(A_1 | \bar{A}_{k-1}, \bar{L}_1)}$$

Continue through all time periods.

# Inverse probability weighting

In time 0, define an inverse probability of treatment weight

$$W^{A_0} = \frac{1}{P(A_0 | L_0)}$$

such that  $A_0$  does not depend on  $L_0$  after weighting

In time 1, do it again

$$W^{A_1} = \frac{1}{P(A_1 | \bar{A}_{k-1}, \bar{L}_1)}$$

Continue through all time periods.


Define the overall weight as the product

$$W^{\bar{A}} = \prod_{k=0}^K \frac{1}{P(A_k | \bar{A}_{k-1}, \bar{L}_k)}$$


# Inverse probability weighting

$$W^{\bar{A}} = \prod_{k=0}^K \frac{1}{P(A_k \mid \bar{A}_{k-1}, \bar{L}_k)}$$

Takes us from this



to this pseudo-population



# Inverse probability weighting with marginal structural models

Finally, we can put a model on top of the weighting.

$$E(Y^{\bar{a}}) = E(Y \mid \bar{A} = \bar{a}) = h(\bar{a})$$

for some function  $h()$  that pools information.

**Example:** Outcomes depend on the proportion of periods treated

$$h(\bar{a}) = \frac{1}{K+1} \sum_{k=0}^K a_k$$

# Learning goals for today

At the end of class, you will be able to:

1. Present treatments that unfold over time in DAGs
2. Reason about the sequential ignorability assumption
3. Apply inverse probability weighting to treatments over time

## Real example: Neighborhood disadvantage

Wodtke, G. T., Harding, D. J., & Elwert, F. (2011). [Neighborhood effects in temporal perspective: The impact of long-term exposure to concentrated disadvantage on high school graduation](#). *American Sociological Review*, 76(5), 713-736.

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

- ▶ Define a neighborhood as a Census tract



# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

- ▶ Define a neighborhood as a Census tract
- ▶ Score that neighborhood along several dimensions
  - ▶ poverty
  - ▶ unemployment
  - ▶ welfare receipt
  - ▶ female-headed households
  - ▶ education
  - ▶ occupational structure

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

- ▶ Define a neighborhood as a Census tract
- ▶ Score that neighborhood along several dimensions
  - ▶ poverty
  - ▶ unemployment
  - ▶ welfare receipt
  - ▶ female-headed households
  - ▶ education
  - ▶ occupational structure
- ▶ Scale by the first principle component

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

- ▶ Define a neighborhood as a Census tract
- ▶ Score that neighborhood along several dimensions
  - ▶ poverty
  - ▶ unemployment
  - ▶ welfare receipt
  - ▶ female-headed households
  - ▶ education
  - ▶ occupational structure
- ▶ Scale by the first principle component
- ▶ Categorize in 5 quintiles

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

How does the neighborhood in which a child lives affect that child's probability of high school completion?

- ▶ Define a neighborhood as a Census tract
- ▶ Score that neighborhood along several dimensions
  - ▶ poverty
  - ▶ unemployment
  - ▶ welfare receipt
  - ▶ female-headed households
  - ▶ education
  - ▶ occupational structure
- ▶ Scale by the first principle component
- ▶ Categorize in 5 quintiles

This 5-value treatment is “neighborhood disadvantage”

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

Neighborhoods are experienced over time:

$$\bar{a}$$

is a trajectory of neighborhood disadvantage over ages 2, 3, ..., 17

The authors study the effect of neighborhood disadvantage,

$$\begin{aligned} E(Y_{\bar{a}} - Y_{\bar{a}'}) &= E(Y_{\bar{a}}) - E(Y_{\bar{a}'}) \\ &= P(Y_{\bar{a}} = 1) - P(Y_{\bar{a}'} = 1), \end{aligned} \quad (1)$$

Example:


$\bar{a}$  is residence in the most advantaged neighborhood each year  
and

$\bar{a}'$  is residence in the most disadvantaged neighborhood each year

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

Problem: Neighborhoods  $A_1$  shape family characteristics  $L_2$ , which confound where people live in the future  $A_2$



**Figure 1.** Causal Graphs for Exposure to Disadvantaged Neighborhoods with Two Waves of Follow-up

Note:  $A_k$  = neighborhood context,  $L_k$  = observed time-varying confounders,  $U$  = unobserved factors,  $Y$  = outcome.

**Table 2.** Time-Dependent Sample Characteristics


Variable	Blacks ( <i>n</i> = 834)			Nonblacks ( <i>n</i> = 1,259)		
	Age 1	Age 10	Age 17	Age 1	Age 10	Age 17
NH disadvantage index, percent						
1st quintile	3.48	3.60	3.48	13.34	19.14	20.65
2nd quintile	3.24	3.72	6.00	19.46	18.67	21.84
3rd quintile	5.28	5.88	7.79	26.13	23.27	22.48
4th quintile	14.87	18.11	18.47	26.13	23.99	21.13
5th quintile	73.14	68.71	64.27	14.93	14.93	13.90
FU head's marital status, percent						
Unmarried	33.93	44.84	52.04	5.88	11.36	15.09
Married	66.07	55.16	47.96	94.12	88.64	84.91
FU head's employment status, percent						
Unemployed	27.22	32.61	33.09	8.10	8.02	9.69
Employed	72.78	67.39	66.91	91.90	91.98	90.31
Public assistance receipt, percent						
Did not receive AFDC	81.06	75.66	82.37	96.27	96.19	97.93
Received AFDC	18.94	24.34	17.63	3.73	3.81	2.07
Homeownership, percent						
Do not own home	69.66	53.48	50.12	40.19	22.32	20.73
Own home	30.34	46.52	49.88	59.81	77.68	79.27
FU income in \$1,000s, mean	19.68	25.04	27.45	32.59	46.65	57.50
FU head's work hours, mean	30.08	26.82	27.51	42.65	40.84	40.68
FU size, mean	5.75	5.32	4.81	4.22	4.69	4.33
Cum. residential moves, mean	.32	2.48	3.64	.32	2.16	3.02

*Note:* NH = neighborhood; FU = family unit. Statistics reported for children not lost to follow-up before age 20 (first imputation dataset).

# Real example: Neighborhood disadvantage

Wodtke et al. 2011

Problem: Neighborhoods  $A_1$  shape family characteristics  $L_2$ , which confound where people live in the future  $A_2$



**Figure 1.** Causal Graphs for Exposure to Disadvantaged Neighborhoods with Two Waves of Follow-up

Note:  $A_k$  = neighborhood context,  $L_k$  = observed time-varying confounders,  $U$  = unobserved factors,  $Y$  = outcome.



# Real example: Neighborhood disadvantage

Wodtke et al. 2011

Solution: MSM-IPW

$$w_i = \prod_{k=1}^K \frac{1}{P(A_k = a_{ki} \mid \bar{A}_{k-1} = \bar{a}_{(k-1)i}, \bar{L}_k = \bar{l}_{ki})}. \quad (4)$$

Also with stabilized weights

$$sw_i = \prod_{k=1}^K \frac{P(A_k = a_{ki} \mid \bar{A}_{k-1} = \bar{a}_{(k-1)i}, L_0 = l_0)}{P(A_k = a_{ki} \mid \bar{A}_{k-1} = \bar{a}_{(k-1)i}, \bar{L}_k = \bar{l}_{ki})}, \quad (5)$$

# Real example: Neighborhood disadvantage


Wodtke et al. 2011

Marginal structural model: Logit

- ▶ 5-category treatment entered numerically
- ▶ Baseline covariates included due to stabilized weights
- ▶ Weights adjust for time-varying confounding

# Real example: Neighborhood disadvantage

Wodtke et al. 2011



**Figure 3.** Predicted Probability of High School Graduation by Neighborhood Exposure

History

Note: NH = Neighborhood

# Learning goals for today

At the end of class, you will be able to:

1. Present treatments that unfold over time in DAGs
2. Reason about the sequential ignorability assumption
3. Apply inverse probability weighting to treatments over time

Let me know what you are thinking

[tinyurl.com/CausalQuestions](https://tinyurl.com/CausalQuestions)

Office hours TTh 11am-12pm and at  
[calendly.com/ianlundberg/office-hours](https://calendly.com/ianlundberg/office-hours)  
Come say hi!