

Lorum ipsum

Author 1 Name, Author 2 Name *

March 2, 2020

This paper is currently in draft mode. Check source to change options.

Abstract

Lorem ipsum dolor sit amet, dicta iudicabit consequat ex vix, veniam legimus appetere has id, an pri graece epicuri detraxit. Ea aliquam expetendis posidonium eos, nam invenire corrumpit imperdiet ei. Et constituto dissentias usu, mel solum erant et. Mel dolorem menandri in.

This is just psuedo text.

2010 Mathematics Subject Classification: Primary 47H05, 52A41, 90C25; Secondary 15A09, 26A51, 26B25, 26E60, 47H09, 47A63. **Keywords:**

1 Schur's Cone and Its Polar

Example 1.1 Define

$$Q = \left\{ x \in \mathbb{R}^n \left| \sum_{i=1}^k x_i \geq 0, k = 1, \dots, n-1, x_1 + \dots + x_n = 0 \right. \right\}. \quad (1.1) \quad \{eqn:cone-1\}$$

Then

$$Q^\circ = \{y \in \mathbb{R}^n | \langle y, x \rangle \geq 0 \forall x \in Q\} \quad (1.2)$$

$$= \{y \in \mathbb{R}^n | y_1 \geq y_2 \geq \dots \geq y_n\}. \quad (1.3)$$

*Subject type, Some Department of Some University, Location of the University, Country. E-mail: author.name@university.edu.

Proof. We have $Q = H(\mathbb{R}_+^{n-1})$, where

$$H = \begin{bmatrix} 1 & & & \\ -1 & 1 & & \\ & -1 & & \\ & & \ddots & 1 \\ & & & -1 \end{bmatrix}, \quad (1.4)$$

H is a bi-diagonal matrix with zeros on the diagonal and -1 on its sub-diagonals. Indeed, for $x = (x_1, \dots, x_{n-1}) \in \mathbb{R}_+^{n-1}$, we have

$$y = Hx \quad (1.5)$$

$$= \begin{bmatrix} 1 \\ -1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} x_1 + \begin{bmatrix} 0 \\ 1 \\ -1 \\ \vdots \\ 0 \end{bmatrix} x_2 + \dots + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ -1 \end{bmatrix} x_{n-1} \quad (1.6)$$

$$= \begin{bmatrix} x_1 \\ x_2 - x_1 \\ x_3 - x_2 \\ \vdots \\ -x_{n-2} + x_{n-1} \\ -x_{n-1} \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix}, \quad (1.7)$$

where $y_1 \geq 0, y_2 + y_1 \geq 0, \dots, y_{n-1} + \dots + y_1 \geq 0$, and $y_1 + \dots + y_n = 0$. So $y \in Q$. Hence $H(\mathbb{R}_+^{n-1}) \subseteq Q$. Similarly, one can write for $y \in Q$, $\exists x \in \mathbb{R}_+^{n-1}$ with $y = Hx$. In fact, $x = y_1, x_2 = y_1 + y_2, \dots, x_{n-1} = y_1 + \dots + y_{n-1}, -x_{n-1} = y_n$ because $y_1 + \dots + y_n = 0$. In particular, $x \in \mathbb{R}_+^{n-1}$.

Note that $y \in Q^\circ \iff y^\top a_i \geq 0$ for $i = 1, \dots, n-1$. Where

$$a_1 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, a_2 = \begin{bmatrix} 0 \\ 1 \\ -1 \\ \vdots \\ 0 \end{bmatrix}, \dots, a_n = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ -1 \end{bmatrix}. \quad (1.8)$$

Now $y^\top a_i \geq 0 \iff y_i - y_{i+1} \geq 0$, for $i = 1, \dots, n-1$. Hence $Q^\circ = \{y \in \mathbb{R}^n | y_1 \geq y_2 \geq \dots \geq y_n\}$.

■

2 Lorentz Cone and its Polar

Example 2.1

$$L = \{(\xi, t) \in \mathbb{R}^n \mid \|\xi\| \leq t, \xi \in \mathbb{R}^{n-1}\}. \quad (2.1)$$

Then the polar cone of L is $L^\circ = \{(x, s) \mid \langle (x, s), (\xi, t) \rangle \geq 0\} = L$.

Proof.

Let $(x, s) \in L^\circ$. Then $\langle (x, s), (\xi, t) \rangle \geq 0, \forall (\xi, t) \in L$, i. e. $\langle x, \xi \rangle + st \geq 0$.

For $t > 0$, we have $\langle x, \xi/t \rangle \geq -s$, i.e. $\langle x, -\xi/t \rangle \leq s$. Using $\|\xi\| \leq t$ and by taking supremum over $\|\xi/t\| \leq 1$, we obtain $\|x\| \leq s$.

Hence $L^\circ = L$. ■

Corollary 2.2 *Let $A \in \mathbb{R}^{n \times n}$ be positive definite. Define*

$$L_A = \{(\xi, t) \in \mathbb{R}^n \mid \sqrt{\langle \xi, A\xi \rangle} \leq t\}.$$

Then $(L_A)^\circ = \{(\xi, t) \mid \sqrt{\langle \xi, A^{-1}\xi \rangle} \leq t\}$.

3 Testing the references

This book by Nesterov [2]. This book by Heinz [1].

4 Testing figures and images

In Figure 1, it has Figure 1a on the left and Figure 1b on the right.



(a) Alto the unicorn.



{fig:two-test-fig-1} (b) Alto and Minty.

{fig:two-test-fig-2}

Figure 1: The above 2 images are test images for the figure, subfigure latex commands.

{fig:two-test-figures}

Acknowledgments

XW were supported by NSERC Discovery grants.

References

- [1] H. H. BAUSCHKE AND P. L. COMBETTES, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, CMS Books in Mathematics, Springer International Publishing, Cham, 2017.
- [2] Y. NESTEROV, *Lectures on Convex Optimization*, vol. 137 of Springer Optimization and Its Applications, Springer International Publishing, 2018.