

The background of the entire image is a light pink color. In the upper half, there is a faint, darker pink silhouette of a world map. A thick, vibrant pink ribbon is draped diagonally across the lower half of the image, starting from the right side and extending towards the bottom left. The ribbon has a slight 3D effect with shadows.

AWARENESS ON

BREAST CANCER



BREAST CANCER



PREVALENCE/BURDEN



2nd most common
cancer in women



1 in 8 women are diagnosed with
breast cancer

SIGNS AND SYMPTOMS



Change in the size/shape
of the breast



Pain in any area of the
breast



Fluid discharge from
nipples other than
breast milk



Any new lump in
the breast

PREVENTION TIPS



Regularly self-examine
your breasts and exercise
regularly



Avoid alcohol & smoking



Regular clinical check-up
every 3 months



Scanning of breast region
(Mammography, MRI)

WHAT IF YOU GET BREAST CANCER



Treatment with medicine
(Chemotherapy)



Surgery and radiation

References: 1. Breast Cancer—Patient Version. Available at: <https://www.cancer.gov/types/breast>. Accessed on: 14.09.2022. 2. Breast Cancer in Young Women. Available at: https://www.cdc.gov/cancer/breast/young_women/bringyourbrave/breast_cancer_young_women/index.htm. Accessed on: 15.09.2022. 3. Breast cancer. Available at: <https://www.who.int/news-room/fact-sheets/detail/breast-cancer>. Accessed on: 14.09.2022. 4. If You Have Breast Cancer. Available at: <https://www.cancer.org/cancer/breast-cancer/if-you-have-breast-cancer.html>. Accessed on: 14.09.2022. 5. Breast Cancer Awareness. Available at: <https://www.cdc.gov/cancer/dccp/resources/features/breastcancerawareness/index.htm>. Accessed on: 14.09.2022.

The content is not intended nor recommended as a substitute for medical advice, diagnosis, or treatment. Always seek the advice of your physician or other qualified healthcare professional regarding any medical questions or condition.

From makers of:

LycorRed[®]
Sachet
Preg
(L-arginine 3gm + Lycopene 4mg + DHA 200mg)

In High Risk Pregnancy



LycorRed[®]

Preg



Sachet

L-arginine 3g + Lycopene 4mg + DHA 200mg

L-arginine + Antioxidant supplementation during pregnancy reduces the risk of pre-eclampsia¹



- In
- Pre-eclampsia
- IUGR
- Oligohydramnios



Usage:-

1 sachet twice daily during 2nd trimester

1. British Medical Journal 2011;342: d2901



Jagsonpal Pharmaceuticals Ltd.

T-210 J, Shahpur Jat, New Delhi, Delhi 110049

Disclaimer: The contents of this scientific issue are developed by Passi HealthCom Pvt. Ltd. exclusively for Jagsonpal Pharmaceuticals Ltd. Although great care has been taken in compiling and checking the information, the authors, Passi HealthCom Pvt. Ltd. and its agents and sponsors shall not be responsible, or in any way liable for any errors, omissions or inaccuracies in this publication whether arising from negligence or otherwise, however, or for any consequences arising therefrom.