

SC 220: Groups and Linear Algebra: Autumn 2020. B.Tech Sem-III

Gautam Dutta

December 6, 2020

Eigenvalues and Eigenvectors
Lecture 22

Eigenvalues and Eigenvectors

- *Eg.1*

Let $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.

Eigenvalues and Eigenvectors

- *Eg.1*

Let $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.

$$A \begin{pmatrix} x \\ 0 \end{pmatrix} = 2 \begin{pmatrix} x \\ 0 \end{pmatrix} \quad \text{and} \quad A \begin{pmatrix} 0 \\ y \end{pmatrix} = 3 \begin{pmatrix} 0 \\ y \end{pmatrix}$$

Eigenvalues and Eigenvectors

- *Eg.1*

Let $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.

$$A \begin{pmatrix} x \\ 0 \end{pmatrix} = 2 \begin{pmatrix} x \\ 0 \end{pmatrix} \quad \text{and} \quad A \begin{pmatrix} 0 \\ y \end{pmatrix} = 3 \begin{pmatrix} 0 \\ y \end{pmatrix}$$

- We say that the action of A on the vectors $\begin{pmatrix} x \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is simple. It just gives a vector parallel to the given vector.

Eigenvalues and Eigenvectors

- *Eg.1*

Let $A = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$.

$$A \begin{pmatrix} x \\ 0 \end{pmatrix} = 2 \begin{pmatrix} x \\ 0 \end{pmatrix} \quad \text{and} \quad A \begin{pmatrix} 0 \\ y \end{pmatrix} = 3 \begin{pmatrix} 0 \\ y \end{pmatrix}$$

- We say that the action of A on the vectors $\begin{pmatrix} x \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ y \end{pmatrix}$ is simple. It just gives a vector parallel to the given vector.
All other vectors get rotated under the action of A .

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

Let $v_1 = \begin{pmatrix} x \\ x \end{pmatrix}$,

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

Let $v_1 = \begin{pmatrix} x \\ x \end{pmatrix}$, $v_2 = \begin{pmatrix} x \\ -x \end{pmatrix}$.

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

Let $v_1 = \begin{pmatrix} x \\ x \end{pmatrix}$, $v_2 = \begin{pmatrix} x \\ -x \end{pmatrix}$.

Then $Av_1 = 4v_1$ and $Av_2 = 2v_2$.

For any other vector v , Av is not parallel to v .

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

Let $v_1 = \begin{pmatrix} x \\ x \end{pmatrix}$, $v_2 = \begin{pmatrix} x \\ -x \end{pmatrix}$.

Then $Av_1 = 4v_1$ and $Av_2 = 2v_2$.

For any other vector v , Av is not parallel to v .

- *Eg.3*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{Then} \quad A \begin{pmatrix} x \\ 0 \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$

- *Eg.2*

Now consider

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 3 \end{pmatrix}$$

Let $v_1 = \begin{pmatrix} x \\ x \end{pmatrix}$, $v_2 = \begin{pmatrix} x \\ -x \end{pmatrix}$.

Then $Av_1 = 4v_1$ and $Av_2 = 2v_2$.

For any other vector v , Av is not parallel to v .

- *Eg.3*

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{Then} \quad A \begin{pmatrix} x \\ 0 \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$$

For any other vector, v , Av is not parallel to v .

Eg.4

$$A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

Eg.4

$$A = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

Under the action of A every vector in \mathbb{R}^2 gets rotated by an angle θ . We can't find any vector in \mathbb{R}^2 such that $Av = \lambda v$ for some number λ .

- *Def.* Eigenvector and Eigenvalue.

Let V be a n -dimensional vector space over a field F .

- *Def.* Eigenvector and Eigenvalue.

Let V be a n -dimensional vector space over a field F . Let T be a linear operator on V .

- *Def.* Eigenvector and Eigenvalue.

Let V be a n -dimensional vector space over a field F . Let T be a linear operator on V .

A non-zero vector $v \in V$ is called an eigenvector of T if $Tv = \lambda v$, $\lambda \in F$.

- *Def.* Eigenvector and Eigenvalue.

Let V be a n -dimensional vector space over a field F . Let T be a linear operator on V .

A non-zero vector $v \in V$ is called an eigenvector of T if $Tv = \lambda v$, $\lambda \in F$.

λ is called the eigenvalue of T corresponding to the eigenvector v .

- How do we find the eigenvalues and eigenvector of T ?

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X,$$

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X, \implies (A - \lambda \mathbb{1})X = 0.$$

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X, \implies (A - \lambda \mathbb{1})X = 0.$$

- The equation has a non-trivial solution if $A - \lambda \mathbb{1}$ is a non-invertible matrix.i.e

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X, \quad \implies \quad (A - \lambda \mathbb{1})X = 0.$$

- The equation has a non-trivial solution if $A - \lambda \mathbb{1}$ is a non-invertible matrix.i.e

$$\det(A - \lambda \mathbb{1}) = 0 \quad (1)$$

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X, \implies (A - \lambda \mathbb{1})X = 0.$$

- The equation has a non-trivial solution if $A - \lambda \mathbb{1}$ is a non-invertible matrix.i.e

$$\det(A - \lambda \mathbb{1}) = 0 \quad (1)$$

This equation is an n^{th} degree polynomial in λ .

- How do we find the eigenvalues and eigenvector of T ?
Let A be the matrix representation of T and X be the matrix representation of the vector v in the basis B of V .
Then

$$AX = \lambda X, \implies (A - \lambda \mathbb{1})X = 0.$$

- The equation has a non-trivial solution if $A - \lambda \mathbb{1}$ is a non-invertible matrix.i.e

$$\det(A - \lambda \mathbb{1}) = 0 \quad (1)$$

This equation is an n^{th} degree polynomial in λ .
It is called the characteristic equation for A

- If the underlying vector space is over the complex field then the characteristic equation has n solutions, (possibly repeated)

- If the underlying vector space is over the complex field then the characteristic equation has n solutions, (possibly repeated) yielding n eigenvalues of A .

- If the underlying vector space is over the complex field then the characteristic equation has n solutions, (possibly repeated) yielding n eigenvalues of A .
- If the underlying field F is not \mathbb{C} then the characteristic equation may yield $k \leq n$ eigenvalues in F .

- If the underlying vector space is over the complex field then the characteristic equation has n solutions, (possibly repeated) yielding n eigenvalues of A .
- If the underlying field F is not \mathbb{C} then the characteristic equation may yield $k \leq n$ eigenvalues in F .
- The eigenvalues are also called the characteristic values of T .

- If the underlying vector space is over the complex field then the characteristic equation has n solutions, (possibly repeated) yielding n eigenvalues of A .
- If the underlying field F is not \mathbb{C} then the characteristic equation may yield $k \leq n$ eigenvalues in F .
- The eigenvalues are also called the characteristic values of T .
- The eigenvectors v_i satisfying $Tv_i = \lambda_i v_i$ are also called the characteristic vectors of T .

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

$$\det \begin{pmatrix} 4 - \lambda & 2 & 0 \\ 2 & 4 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{pmatrix} = 0$$

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

$$\det \begin{pmatrix} 4 - \lambda & 2 & 0 \\ 2 & 4 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{pmatrix} = 0 \quad (3 - \lambda)[(4 - \lambda)^2 - 4] = 0$$

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

$$\det \begin{pmatrix} 4 - \lambda & 2 & 0 \\ 2 & 4 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{pmatrix} = 0 \quad (3 - \lambda)[(4 - \lambda)^2 - 4] = 0$$

Eigenvalues are $\lambda_1 = 2, \lambda_2 = 3, \lambda_3 = 6$.

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

$$\det \begin{pmatrix} 4 - \lambda & 2 & 0 \\ 2 & 4 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{pmatrix} = 0 \quad (3 - \lambda)[(4 - \lambda)^2 - 4] = 0$$

Eigenvalues are $\lambda_1 = 2, \lambda_2 = 3, \lambda_3 = 6$.

Corresponding eigenvectors are

Eg.1

$$A = \begin{pmatrix} 4 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad AX = \lambda X$$

To find a nontrivial X we must have $\det(A - \lambda \mathbb{1}) = 0$ i.e

$$\det \begin{pmatrix} 4 - \lambda & 2 & 0 \\ 2 & 4 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{pmatrix} = 0 \quad (3 - \lambda)[(4 - \lambda)^2 - 4] = 0$$

Eigenvalues are $\lambda_1 = 2, \lambda_2 = 3, \lambda_3 = 6$.

Corresponding eigenvectors are

$$X_1 = \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Consider

$$A = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 4 & 1 \\ 0 & 0 & 5 \end{pmatrix}$$

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Consider

$$A = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 4 & 1 \\ 0 & 0 & 5 \end{pmatrix}$$

The characteristic equation is

$$\det \begin{pmatrix} 2 - \lambda & 3 & 0 \\ 0 & 4 - \lambda & 1 \\ 0 & 0 & 5 - \lambda \end{pmatrix} = 0$$

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Consider

$$A = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 4 & 1 \\ 0 & 0 & 5 \end{pmatrix}$$

The characteristic equation is

$$\det \begin{pmatrix} 2-\lambda & 3 & 0 \\ 0 & 4-\lambda & 1 \\ 0 & 0 & 5-\lambda \end{pmatrix} = 0 \implies (2-\lambda)(4-\lambda)(5-\lambda) = 0$$

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Consider

$$A = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 4 & 1 \\ 0 & 0 & 5 \end{pmatrix}$$

The characteristic equation is

$$\det \begin{pmatrix} 2 - \lambda & 3 & 0 \\ 0 & 4 - \lambda & 1 \\ 0 & 0 & 5 - \lambda \end{pmatrix} = 0 \implies (2 - \lambda)(4 - \lambda)(5 - \lambda) = 0$$

So the eigenvalues are $\lambda_1 = 2$, $\lambda_2 = 4$ and $\lambda_3 = 5$.

Eg. 2:

If A is an upper(lower) triangular matrix, then eigenvalues are diagonal elements.

Consider

$$A = \begin{pmatrix} 2 & 3 & 0 \\ 0 & 4 & 1 \\ 0 & 0 & 5 \end{pmatrix}$$

The characteristic equation is

$$\det \begin{pmatrix} 2-\lambda & 3 & 0 \\ 0 & 4-\lambda & 1 \\ 0 & 0 & 5-\lambda \end{pmatrix} = 0 \implies (2-\lambda)(4-\lambda)(5-\lambda) = 0$$

So the eigenvalues are $\lambda_1 = 2$, $\lambda_2 = 4$ and $\lambda_3 = 5$. The corresponding eigenvectors are

$$X_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}; \quad X_2 = \begin{pmatrix} 3 \\ 2 \\ 0 \end{pmatrix}; \quad X_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$$

0 Eigenvalues:

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

- $\implies A$ is not invertible.

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

- $\implies A$ is not invertible.
- $\implies AX = 0$ has non-trivial solutions.

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

- $\implies A$ is not invertible.
- $\implies AX = 0$ has non-trivial solutions.
- The solution space of this equation consists of eigenvectors of A corresponding to the 0-eigenvalue.

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

- $\implies A$ is not invertible.
- $\implies AX = 0$ has non-trivial solutions.
- The solution space of this equation consists of eigenvectors of A corresponding to the 0-eigenvalue. This indicates the following proposition.

0 Eigenvalues:

- If 0 is one of the eigenvalues of A then

$$\det(A - 0 \cdot \mathbb{1}) = 0 \implies \det A = 0$$

- $\implies A$ is not invertible.
- $\implies AX = 0$ has non-trivial solutions.
- The solution space of this equation consists of eigenvectors of A corresponding to the 0-eigenvalue. This indicates the following proposition.
- *Proposition 35*
An $n \times n$ matrix is invertible if and only if none of its eigenvalue is 0.

- *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

- *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

- *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors. Then

- *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

- *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

$$T_1(c_1 v_1 + c_2 v_2 + \dots + c_k v_k) = 0$$

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

$$T_1(c_1 v_1 + c_2 v_2 + \dots + c_k v_k) = 0$$

$$\therefore c_1(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3) \dots (\lambda_1 - \lambda_k)v_1 = 0$$

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

$$T_1(c_1 v_1 + c_2 v_2 + \dots + c_k v_k) = 0$$

$$\therefore c_1(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3) \dots (\lambda_1 - \lambda_k)v_1 = 0$$

Since $v_1 \neq 0$ and $\lambda_1 \neq \lambda_2, \lambda_3, \dots, \lambda_k$,

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

$$T_1(c_1 v_1 + c_2 v_2 + \dots + c_k v_k) = 0$$

$$\therefore c_1(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3) \dots (\lambda_1 - \lambda_k)v_1 = 0$$

Since $v_1 \neq 0$ and $\lambda_1 \neq \lambda_2, \lambda_3, \dots, \lambda_k$,

we have $c_1 = 0$.

• *Proposition 36*

Eigenvectors corresponding to distinct eigenvalues are linearly independent.

Proof:

Let $\lambda_1, \lambda_2, \dots, \lambda_k$ be k distinct eigenvalues of an operator T .

Let v_1, v_2, \dots, v_k be the corresponding eigenvectors.

Then

$$(T - \lambda_i \mathbb{1})v_i = 0, \quad i = 1, 2, \dots, k$$

Let $c_1 v_1 + c_2 v_2 + \dots + c_k v_k = 0$.

Consider the operator

$$T_1 = (T - \lambda_2 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

$$T_1(c_1 v_1 + c_2 v_2 + \dots + c_k v_k) = 0$$

$$\therefore c_1(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3) \dots (\lambda_1 - \lambda_k)v_1 = 0$$

Since $v_1 \neq 0$ and $\lambda_1 \neq \lambda_2, \lambda_3, \dots, \lambda_k$,

we have $c_1 = 0$.

.....contd.

.....contd.

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



- If an eigenvalue repeats k times then we may not get k linearly independent eigenvectors.

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



- If an eigenvalue repeats k times then we may not get k linearly independent eigenvectors.
- *Eg.*

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



- If an eigenvalue repeats k times then we may not get k linearly independent eigenvectors.
- Eg. If $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ Then the characteristic equation is

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



- If an eigenvalue repeats k times then we may not get k linearly independent eigenvectors.
- Eg. If $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ Then the characteristic equation is $(1 - \lambda)^2 = 0$. So the eigenvalue $\lambda = 1$ has multiplicity 2.

.....contd.

Now considering the operator

$$T_2 = (T - \lambda_1 \mathbb{1})(T - \lambda_3 \mathbb{1}) \dots (T - \lambda_k \mathbb{1})$$

we can show that $c_2 = 0$.

Likewise $c_3 = \dots = c_k = 0$.

$\therefore v_1, v_2, \dots, v_k$ are linearly independent.



- If an eigenvalue repeats k times then we may not get k linearly independent eigenvectors.
- *Eg.* If $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ Then the characteristic equation is $(1 - \lambda)^2 = 0$. So the eigenvalue $\lambda = 1$ has multiplicity 2. But the solution space of the homogeneous equation $(A - \lambda \mathbb{1})X = 0$ is only one dimensional yielding only one linearly independent eigenvector viz. $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Diagonalization

Diagonalization

Let A be a $n \times n$ matrix.

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

$$P^{-1}P = \mathbb{1} \implies R_i C_j = \delta_{ij}.$$

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

$$P^{-1}P = \mathbb{1} \implies R_i C_j = \delta_{ij}.$$

$$\therefore P^{-1}AP = P^{-1}(AC_1, AC_2, \dots, AC_n)$$

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

$$P^{-1}P = \mathbb{1} \implies R_i C_j = \delta_{ij}.$$

$$\begin{aligned} \therefore P^{-1}AP &= P^{-1}(AC_1, AC_2, \dots, AC_n) \\ &= \begin{pmatrix} R_1 \\ \vdots \\ R_n \end{pmatrix} (\lambda_1 C_1 \ \dots \ \lambda_n C_n) \end{aligned}$$

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

$$P^{-1}P = \mathbb{1} \implies R_i C_j = \delta_{ij}.$$

$$\therefore P^{-1}AP = P^{-1}(AC_1, AC_2, \dots, AC_n)$$

$$\begin{aligned} &= \begin{pmatrix} R_1 \\ \vdots \\ R_n \end{pmatrix} (\lambda_1 C_1 \dots \lambda_n C_n) = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix} \\ &= D \end{aligned}$$

Diagonalization

Let A be a $n \times n$ matrix. Let C_1, C_2, \dots, C_n be n linearly independent eigenvectors of A corresponding to the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

Construct the $n \times n$ matrix $P = (C_1, C_2, \dots, C_n)$.

P is invertible since its columns C_i are linearly independent.

Let R_1, R_2, \dots, R_n be the rows of P^{-1} .

$$P^{-1}P = \mathbb{1} \implies R_i C_j = \delta_{ij}.$$

$$\begin{aligned} \therefore P^{-1}AP &= P^{-1}(AC_1, AC_2, \dots, AC_n) \\ &= \begin{pmatrix} R_1 \\ \vdots \\ R_n \end{pmatrix} (\lambda_1 C_1 \dots \lambda_n C_n) = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{pmatrix} \\ &= D \end{aligned}$$

where D is a diagonal matrix with the diagonal elements as the eigenvalues of A .

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3 - \lambda & 0 & -1 \\ 0 & 4 - \lambda & 0 \\ -1 & 0 & 3 - \lambda \end{pmatrix} = 0$

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0$$

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0 \implies \lambda = 4 \text{ or } \lambda = 2.$$

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0 \implies \lambda = 4 \text{ or } \lambda = 2.$$

Eigenvalue 4 occurs with multiplicity 2.

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0 \implies \lambda = 4 \text{ or } \lambda = 2.$$

Eigenvalue 4 occurs with multiplicity 2.

Eigenvectors: For $\lambda = 2$,

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0 \implies \lambda = 4 \text{ or } \lambda = 2.$$

Eigenvalue 4 occurs with multiplicity 2.

Eigenvectors: For $\lambda = 2$, $C_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

For $\lambda = 4$ we get two linearly independent eigenvectors.

Eg.1:

$$A = \begin{pmatrix} 3 & 0 & -1 \\ 0 & 4 & 0 \\ -1 & 0 & 3 \end{pmatrix}$$

Characteristic equation : $\det \begin{pmatrix} 3-\lambda & 0 & -1 \\ 0 & 4-\lambda & 0 \\ -1 & 0 & 3-\lambda \end{pmatrix} = 0$

$$\therefore (4-\lambda)[(3-\lambda)^2 - 1] = 0 \implies \lambda = 4 \text{ or } \lambda = 2.$$

Eigenvalue 4 occurs with multiplicity 2.

Eigenvectors: For $\lambda = 2$, $C_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

For $\lambda = 4$ we get two linearly independent eigenvectors.

$$C_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}; \quad C_3 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

Using C_1, C_2, C_3 we construct

Using C_1, C_2, C_3 we construct $P = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix}$.

Using C_1, C_2, C_3 we construct $P = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix}$.

$$P^{-1} = \begin{pmatrix} \frac{1}{2} & 0 & -\frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{pmatrix}.$$

Using C_1, C_2, C_3 we construct $P = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix}$.

$$P^{-1} = \begin{pmatrix} \frac{1}{2} & 0 & -\frac{1}{2} \\ 0 & 1 & 0 \\ \frac{1}{2} & 0 & \frac{1}{2} \end{pmatrix}.$$

$$P^{-1}AP = \text{diag}(4, 4, 2)$$

$$\text{Eg.2: } A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}.$$

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{matrix} y = 2z \\ x = 9z \end{matrix}$$

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{matrix} y = 2z \\ x = 9z \end{matrix}$$

$$\therefore C_1 = \begin{pmatrix} 9 \\ 2 \\ 1 \end{pmatrix}.$$

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{matrix} y = 2z \\ x = 9z \end{matrix}$$

$$\therefore C_1 = \begin{pmatrix} 9 \\ 2 \\ 1 \end{pmatrix}. \text{ For } \lambda = 2 \text{ we get } C_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{aligned} y &= 2z \\ x &= 9z \end{aligned}$$

$$\therefore C_1 = \begin{pmatrix} 9 \\ 2 \\ 1 \end{pmatrix}. \text{ For } \lambda = 2 \text{ we get } C_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Though this eigenvalue has multiplicity 2 there is only one corresponding eigenvector.

Eg.2: $A = \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$. Eigenvalues are 2, 2 and 3.

Eigenvector for $\lambda = 3$:

$$\begin{pmatrix} -1 & 4 & 1 \\ 0 & -1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{aligned} y &= 2z \\ x &= 9z \end{aligned}$$

$$\therefore C_1 = \begin{pmatrix} 9 \\ 2 \\ 1 \end{pmatrix}. \text{ For } \lambda = 2 \text{ we get } C_2 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

Though this eigenvalue has multiplicity 2 there is only one corresponding eigenvector. We can't construct the matrix P such that $P^{-1}AP = \text{diag}(2, 2, 3)$.

So A is not diagonalizable.

Conclusion

Conclusion

- If all the eigenvalues of a matrix A are distinct, then it is diagonalizable.

Conclusion

- If all the eigenvalues of a matrix A are distinct, then it is diagonalizable.
- If an eigenvalue has multiplicity k , then A is diagonalizable only if the eigenspace for this eigenvalue is k -dimensional.

Conclusion

- If all the eigenvalues of a matrix A are distinct, then it is diagonalizable.
- If an eigenvalue has multiplicity k , then A is diagonalizable only if the eigenspace for this eigenvalue is k -dimensional.
- Let T be a linear operator on a vector space V . If the matrix representation of T is diagonalizable then the range of T is spanned by the eigenvectors of the nonzero eigenvalues of T .

Conclusion

- If all the eigenvalues of a matrix A are distinct, then it is diagonalizable.
- If an eigenvalue has multiplicity k , then A is diagonalizable only if the eigenspace for this eigenvalue is k -dimensional.
- Let T be a linear operator on a vector space V . If the matrix representation of T is diagonalizable then the range of T is spanned by the eigenvectors of the nonzero eigenvalues of T .
 $\text{kernel } T$ is spanned by the eigenvectors of the 0-eigenvalue of T .

- **Def.: Similarity**

A matrix A is similar to matrix B if \exists an invertible matrix P such that $B = PAP^{-1}$.

- **Def.: Similarity**

A matrix A is similar to matrix B if \exists an invertible matrix P such that $B = PAP^{-1}$.

- Similarity is an equivalence relation on the set of matrices.

- **Def.: Similarity**

A matrix A is similar to matrix B if \exists an invertible matrix P such that $B = PAP^{-1}$.

- Similarity is an equivalence relation on the set of matrices.
- Similar matrices have the same characteristic polynomial. Hence they will have the same set of eigenvalues.