


BMJ Open Efficacy in predicting mortality of patients with heart failure using heart rate before intensive care unit discharge: a retrospective cohort study from MIMIC-IV Database

Chia-Ying Hsiao,^{1,2} Min-I Su,^{2,3,4} Yu-Cheng Chang,^{5,6} Ying-Hsiang Lee,^{2,7,8} Po-Lin Lin,^{9,10} Wei-Ru Chiou ^{2,3}

To cite: Hsiao C-Y, Su M-I, Chang Y-C, *et al.* Efficacy in predicting mortality of patients with heart failure using heart rate before intensive care unit discharge: a retrospective cohort study from MIMIC-IV Database. *BMJ Open* 2024;**14**:e087725. doi:10.1136/bmjopen-2024-087725

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-087725>).

Received 19 April 2024
Accepted 22 November 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Wei-Ru Chiou;
clementchiou@gmail.com

ABSTRACT

Objective Heart rate serves as a critical prognostic factor in heart failure (HF) patients. We hypothesise that elevated heart rate in critically ill HF patients on discharge from the intensive care unit (ICU) could be linked to adverse outcomes.

Design A single-centre retrospective cohort study.

Setting This study used data collected between 2008 and 2019 from the Medical Information Mart for Intensive Care IV (V.2.0) Database.

Participants From the 76 943 ICU stays, we enrolled 2365 patients in this study.

Primary and secondary outcome measures We observed correlations between in-hospital mortality and ICU discharge heart rate of 83.56 ± 15.81 beats per minute (bpm) (survivors) vs 93.84 ± 17.28 bpm (non-survivors, $p < 0.001$). Total mortality showed similar trends, with 83.67 ± 15.36 bpm (survivors) vs 85.23 ± 17.25 bpm (non-survivors, $p = 0.027$), as did ICU readmissions at 83.55 ± 15.77 bpm (non-readmitted) vs 88.64 ± 17.49 bpm (readmitted, $p < 0.001$).

Results Given multivariate analysis, the ICU discharge heart rate strongly predicted in-hospital mortality (HR 1.032 (95% CI 1.022 to 1.041), $p < 0.001$), total mortality (HR 1.008 (95% CI 1.004 to 1.013), $p < 0.001$) and ICU readmission (HR 1.018 (95% CI 1.010 to 1.025), $p < 0.001$). Patients with an ICU discharge heart rate > 90 bpm demonstrated significantly higher in-hospital mortality (HR 2.639 (95% CI 1.898 to 3.669), $p < 0.001$), total mortality (HR 1.342 (95% CI 1.163 to 1.550), $p < 0.001$) and ICU readmission rates (HR 1.781 (95% CI 1.413 to 2.243), $p < 0.001$).

Conclusion The findings suggest that HF patients with an elevated heart rate (> 90 bpm) at ICU discharge are more likely to experience increased in-hospital mortality, total mortality and ICU readmissions, indicating potential negative outcomes.

INTRODUCTION

Heart rate is related to disease and stress, and the maximum heart rate is an indicator of short-term mortality in critical illness.¹ A

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The cohort comprised of a large and extensive coverage of 'real-world' intensive care unit (ICU) cases.
- ⇒ Lack of detailed information on medications and echocardiography.
- ⇒ Since this study employed an observational data, we could only study the general mortality rate and not particular causes of death.

heart rate ≥ 90 beats per minute (bpm) at the time of multiple organ dysfunction diagnosis is an independent risk factor for increased 28-day mortality.¹ Patients with a sustained heart rate > 95 bpm for over 12 hours during any 24-hour period in the intensive care unit (ICU) experienced a higher incidence of major cardiac events and longer ICU stays²

Heart failure (HF) and ICU

According to the Romanian Acute Heart Failure Syndromes registry, 10.7% of patients with the condition required ICU care, with admission to the ICU being linked to an increased risk of in-hospital mortality (17.3% vs 6.5%, $p = 0.002$).³ Within a Taiwan national database of 192 201 patients who were admitted to the ICU, a total of 25 263 (13.14%) patients were readmitted. HF was identified as a significant risk factor for readmission to the ICU (HR 3.365 (95% CI 3.047 to 3.717), $p \leq 0.001$).⁴

Heart rate and HF

Higher heart rate was related to the development of regional and global left ventricular (LV) dysfunction independent of subclinical atherosclerosis and coronary heart disease.⁵ In the Organized Program to Initiate Life-saving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF) study,

the mean heart rate was 87 ± 21.5 bpm in survivors and 89 ± 22.7 bpm in non-survivors during hospitalisation. Multivariable analysis showed that every 10 bpm increase in heart rate, between 65 and 110 bpm, was associated with higher in-hospital mortality (OR 1.094 (95% CI 1.062 to 1.127), $p < 0.0001$).⁶ In the Get With the Guidelines-Heart Failure (GWTG-HF) study, in-hospital mortality risk is calculated by summing points assigned to each predictor variable, with a total score ranging from 0 to 100. Heart rate is one of these factors: a rate of 79 bpm or below is assigned 0 points, while a rate of 105 bpm or higher is assigned 8 points.⁷

In 2001, the Cardiac Insufficiency Bisoprolol Study II (CIBIS II) trial demonstrated that heart rate can not only function as an indirect marker of disease but may also directly contribute to the development of HF.⁸ The Systolic Heart failure Treatment with the IF Inhibitor Ivabradine Trial (SHIFT) trial showed that ivabradine treatment helps systolic HF patients with sinus rhythm and a heart rate of ≥ 70 bpm.⁹ In patients with worsening HF, the first hospital admissions were lower in the group with the lowest heart rate (70 to < 72 bpm) compared with those with the highest baseline heart rate (≥ 87 bpm). In the placebo group, patients with a heart rate ≥ 87 bpm had more than double the risk of cardiovascular death or hospital admission for worsening HF compared with those with the lowest heart rate (HR 2.34, 95% CI 1.84 to 2.98, $p < 0.001$). Event rates increased by 3% for every 1 bpm rise from baseline and by 16% for every 5 bpm increase.¹⁰ Ivabradine treatment is associated with reduced risks of cardiovascular mortality, all-cause mortality and HF rehospitalisation within 1 year among patients with acute decompensated HF with reduced ejection fraction (HFrEF) in real-world populations.¹¹ Among patients with HFrEF, ivabradine treatment had a more notable benefit on haemodynamic stability.¹²

Limitations of previous studies

Based on the aforementioned information, we summarise the following key points. First, heart rhythm plays a crucial role as an important prognostic factor in patients with HF. Second, pharmacological management aimed at controlling the heart rhythm in hospitalised HF patients holds the potential to improve their prognosis. Third, it has been observed that over 10% of patients hospitalised with acute HF require intensive care, and they have a higher mortality rate. However, there is currently a lack of relevant studies investigating the association between post-treatment heart rhythm control in critically ill HF patients on transfer from the ICU and their future clinical outcomes.

Aims of the current study

Our objective in the present study is to address this research gap by exploring the relationship between the heart rate before ICU discharge and mortality in HF patients with the Medical Information Mart for Intensive Care IV (MIMIC IV) Database, thereby providing a more

comprehensive understanding and guidance for clinical practice.

METHODS

Patient and public involvement

Patients and the public were not directly involved in either the design or the implementation of this study, as it relied on previously collected data.

Data source

This cohort study was conducted using data from the MIMIC-IV V.2.0 Database, which was collected between 2008 and 2019.^{13 14} MIMIC-IV is a publicly available and real-world clinical database that was maintained by the Beth Israel Deaconess Medical Center during the same period. It includes records of over 200 000 emergency department admissions and over 60 000 ICU stays. One author, M-IS, has finished the Collaborative Institutional Training Initiative examination (certification number: 42188048) and achieved access to the database for data extraction. The code for extracting data can be found on GitHub (<https://github.com/MIT-LCP/mimic-iv>).¹⁵

Eligibility criteria

Our study included only adult patients who were admitted to the ICU due to HF and older than 18 years. The diagnosis of HF was determined using the International Classification of Diseases, 10th Revision, and the following codes were considered: I500, I501, I509, I110, I130 and I132. In cases where a patient was admitted to the ICU multiple times, only data from the first ICU stay were included. We eliminated from our analysis the patients who were discharged from the ICU within 24 hours and those with missing data for vital signs such as blood pressure and heart rate within 24 hours before discharge (figure 1).

Definitions of heart rate and outcomes measures

The ICU discharge heart rate was defined as the last heart rate before discharge from the ICU, and the average ICU discharge heart rate was defined as the 24-hour average heart rate before discharge from the ICU. The primary outcomes were in-hospital mortality and total mortality. The total mortality was populated from state death records. The secondary outcome was ICU readmission in the same hospitalisation. ICU readmission was defined as when a patient was readmitted to the ICU after an interval of more than 8 hours since the previous discharge from the ICU.

Demographic and clinical characteristics

In our study, we included demographic and clinical characteristics recorded in the ICU. We also calculated various scores within the first 24 hours of ICU admission and vital signs such as heart rate, blood pressure, respiratory rate, SpO₂, urine output and serum creatinine were extracted throughout the entire ICU stay (table 1). These covariates, which included basic demographic information and




Figure 1 Study flowchart. ICD-10, International Classification of Diseases, 10th Revision; ICU, intensive care unit; MIMIC-IV, Medical Information Mart for Intensive Care IV.

clinical characteristics of patients, were chosen based on previous related studies.

Statistical analysis

Continuous variables are expressed as the mean and SD. Categorical variables are presented as the number and percentage (%). The χ^2 test or Fisher's exact test was used to test comparisons between groups for categorical variables, whereas the t test was used for comparisons of continuous variables as appropriate. We used receiver operating characteristic (ROC) curves to identify the last heart rate before leaving the ICU cut-off values for each study outcome. Cox regression analysis was conducted to assess the association between heart rate and outcomes, with the results expressed as HR and corresponding 95% CIs. Kaplan-Meier survival curves were produced for three outcomes: in-hospital mortality, total mortality and ICU readmission. All analyses were performed in R (V.4.2.3) and SPSS (V.20). A threshold of $p < 0.05$ (two sided) was considered statistically significant.

RESULTS

Study population

Considering the 76 943 admissions registered in the ICU, we enrolled 2365 adult HF patients in this study (table 1). During the follow-up period of 140.26 ± 246.02 days, the average length of ICU stay was 4.49 ± 5.10 days and the mean duration of total hospital stay was 13.91 ± 12.44 days. The female proportion was 40.47% ($n=957$), the mean age

of the study population was 69.7 ± 12.69 years and the body mass index (BMI) was 29.83 ± 8.31 kg/m². The mean heart rate during admission to the ICU was 87.66 ± 20.46 bpm, the mean ICU discharge heart rate was 84.26 ± 16.12 bpm and the mean 24-hour average ICU discharge heart rate was 83.82 ± 14.49 bpm. The underlying diseases of the enrolled population included myocardial infarction (40.13%, $n=949$), cerebrovascular disease (14.25%, $n=337$) and diabetes mellitus (47.15%, $n=1115$).

Patient follow-up

The in-hospital mortality was 6.81% ($n=161$), total mortality was 37.76% ($n=893$) and the ICU readmission rate was 14.04% ($n=332$). In terms of in-hospital/total mortality and ICU readmission, no significant difference was observed across the sexes. Patients who were younger and had a lower BMI demonstrated a lower rate of in-hospital mortality ($p=0.001$) and total mortality ($p < 0.001$) (table 2). The in-hospital mortality, total mortality and ICU readmission were all related to lower ICU discharge heart rate, 24-hour average ICU discharge heart rate and ICU admission heart rate. Regarding the outcome of in-hospital mortality, the ICU discharge heart rate was 83.56 ± 15.81 bpm (survival) vs 93.84 ± 17.28 bpm (dead) ($p < 0.001$), and the 24-hour average ICU discharge heart rate was 83.20 ± 14.29 bpm (survival) vs 92.41 ± 14.55 bpm (dead) ($p < 0.001$). Regarding the outcome of total mortality, the ICU discharge heart rate was 83.67 ± 15.36 bpm (survival) vs 85.23 ± 17.25 bpm

Table 1 Baseline characteristics

Variable	All patients, n=2365
Sex (female) (%)	957 (40.47)
Age (mean (SD))	69.7 (12.69)
Weight (kg) (mean (SD))	84.86 (25.34)
Height (cm) (mean (SD))	168.45 (10.97)
BMI (kg/m ²) (mean (SD))	29.83 (8.31)
MBP (mm Hg) (mean (SD))	84.13 (19.15)
ICU discharge heart rate (bpm) (mean (SD))	84.26 (16.12)
ICU discharge 24-hour average heart rate (bpm) (mean (SD))	83.82 (14.49)
ICU admission heart rate (bpm) (mean (SD))	87.66 (20.46)
Temperature (°C) (mean (SD))	36.67 (0.69)
SpO ₂ (%) (mean (SD))	96.69 (3.77)
Urine output (mL) (mean (SD))	1770.94 (1317.76)
Creatinine (mg/dL) (mean (SD))	1.17 (0.96)
Haemoglobin (g/L) (mean (SD))	101.4 (22.6)
Myocardial infarction (%)	949 (40.13)
Cerebrovascular disease (%)	337 (14.25)
Chronic pulmonary disease (%)	767 (32.43)
Diabetes mellitus (%)	1115 (47.15)
Renal disease (%)	1046 (44.23)
GCS (mean (SD))	13.03 (2.96)
SOFA Score (mean (SD))	5.87 (3.48)
CCI (mean (SD))	7.90 (2.57)
APS III (mean (SD))	48.96 (19.23)
SAPSII Score (mean (SD))	39.29 (11.69)
Follow-up period (day) (mean (SD))	140.26 (246.02)
Length of ICU stay (day) (mean (SD))	4.49 (5.10)
Length of hospital stay (day) (mean (SD))	13.91 (12.44)
In-hospital mortality (%)	161 (6.81)
Total mortality (%)	893 (37.76)
ICU readmission (%)	332 (14.04)

APS III, Acute Physiology Score III; BMI, body mass index; CCI, Charlson Comorbidity Score; GCS, Glasgow Coma Scale; ICU, intensive care unit; MBP, mean blood pressure; SAPSII, Simplified Acute Physiology Score II; SOFA, Sequential Organ Failure Assessment.

(dead) ($p=0.027$), and the 24-hour average ICU discharge heart rate was 83.18 ± 14.00 bpm vs 84.89 ± 15.21 bpm ($p=0.006$). Regarding the outcome of ICU readmission, the ICU discharge heart rate was 83.55 ± 15.77 bpm (not ICU readmission) vs 88.64 ± 17.49 bpm (ICU readmission) ($p<0.001$), and the 24-hour average ICU discharge heart rate was 83.15 ± 14.42 bpm vs 87.94 ± 14.23 bpm ($p<0.001$) (table 2). Online supplemental figure S1 shows the area under the curve for the study outcomes. According to the ROC curve, the optimal cut-off points for in-hospital mortality, total mortality and ICU readmission were 87.5 bpm, 91.5 bpm and 89.5 bpm, respectively. Thus, for this

study, we selected 90 bpm as the threshold for the correlation with our outcomes of interest.

Mortality and ICU readmission

The primary and secondary outcomes were significantly higher in the group with a heart rate >90 bpm (table 3). The univariate analysis showed that elevated ICU discharge heart rate was associated with significantly increased in-hospital mortality (HR 1.034 (95% CI 1.025 to 1.042), $p<0.001$), total mortality (HR 1.005 (95% CI 1.001 to 1.009), $p=0.012$) and ICU readmission rate (HR 1.018 (95% CI 1.012 to 1.024), $p<0.001$). After multivariate analysis, the ICU discharge heart rate was also observed to be a strong predictor of in-hospital mortality (HR 1.032 (95% CI 1.022 to 1.041), $p<0.001$), total mortality (HR 1.008 (95% CI 1.004 to 1.013), $p<0.001$) and ICU readmission (HR 1.018 (95% CI 1.010 to 1.025), $p<0.001$). An ICU discharge heart rate greater than 90 bpm was significantly related to in-hospital mortality (HR 2.639 (95% CI 1.898 to 3.669), $p<0.001$), total mortality (HR 1.342 (95% CI 1.163 to 1.550), $p<0.001$) and ICU readmission (HR 1.781 (95% CI 1.413 to 2.243), $p<0.001$) (online supplemental table S1). The Kaplan-Meier analysis also showed that a heart rate >90 bpm was significantly related to in-hospital mortality ($p<0.001$), total mortality ($p<0.001$) and ICU readmission ($p<0.001$) (online supplemental figure S1).

DISCUSSION

In this clinical investigation, we were able to demonstrate that heart rate at ICU discharge was a predictor of mortality in HF patients in the MIMIC IV group. Consistent with prior research on the general population or specific patient groups, we found that a higher heart rate before ICU discharge was associated with increased in-hospital mortality among ICU patients. This association was statistically significant with an adjusted HR of 1.032 for each 1 bpm increase in heart rate in the ICU. A study has shown that heart rate on admission can predict in-hospital mortality in patients with ischaemic stroke, with a HR of 4.42 for a 10 bpm increase in heart rate.¹⁶ Furthermore, high heart rate 24–36 hours after admission has also been associated with in-hospital mortality in patients with HF.¹⁷

A study found that keeping the heart rate below 100 bpm within the first day of admission could decrease mortality in ICU patients.¹⁸ Another study showed that heart rate measured 24 hours before ICU discharge was independently linked to in-hospital and posthospital mortality after ICU discharge.¹⁹ One study using the MIMIC-III Database showed a significant association of prolonged elevated heart rate with decreased survival in a large and heterogeneous cohort of ICU patients.²⁰ A separate study using the MIMIC-IV Database demonstrated that the group of patients with rheumatic heart disease who survived had a lower heart rate.²¹ The HF mortality prediction models have served to advance

Table 2 Baseline characteristics between the total population and primary/secondary outcomes

Variable	In-hospital mortality			Total mortality			ICU readmission		
	Survival	Dead	P value	Survival	Dead	P value	No	Yes	P value
Patient (%)	2204 (93.20)	161 (6.81)		1472 (62.24)	893 (37.76)		2033 (86.96)	332 (14.04)	
Sex (F) (%)	887 (40.25)	70 (43.48)	0.454	575 (39.06)	382 (42.78)	0.077	822 (40.43)	135 (40.66)	0.952
Age (mean (SD))	69.31 (12.68)	75.10 (11.61)	<0.001	67.41 (12.58)	73.49 (11.93)	<0.001	69.76 (12.70)	69.38 (12.64)	0.619
Weight (kg) (mean (SD))	85.37 (25.46)	77.88 (22.58)	<0.001	88.02 (26.08)	79.64 (23.15)	<0.001	85.04 (25.32)	83.73 (25.46)	0.382
Height (cm) (mean (SD))	168.52 (11.00)	167.52 (10.56)	0.264	169.20 (10.90)	167.20 (10.98)	<0.001	168.54 (10.95)	167.90 (11.09)	0.33
BMI (kg/m ²) (mean (SD))	29.98 (8.35)	27.69 (7.40)	0.001	30.66 (8.40)	28.46 (7.98)	<0.001	29.86 (8.30)	29.63 (8.42)	0.64
MBP (mmHg) (mean (SD))	84.11 (19.24)	84.39 (17.92)	0.857	84.43 (18.76)	83.64 (19.78)	0.331	84.05 (19.12)	84.61 (19.39)	0.622
ICU discharge heart rate (bpm) (mean (SD))	83.56 (15.81)	93.84 (17.28)	<0.001	83.67 (15.36)	85.23 (17.25)	0.027	83.55 (15.77)	88.64 (17.49)	<0.001
ICU discharge 24-hour average heart rate (bpm) (mean (SD))	83.20 (14.29)	92.41 (14.55)	<0.001	83.18 (14.00)	84.89 (15.21)	0.006	83.15 (14.42)	87.94 (14.23)	<0.001
ICU admission heart rate (bpm) (mean (SD))	87.13 (20.27)	94.90 (21.66)	<0.001	86.86 (20.16)	88.98 (20.88)	0.015	87.09 (20.21)	91.16 (21.63)	0.001
Temperature (°C) (mean (SD))	36.67 (0.69)	36.68 (0.71)	0.814	36.65 (0.71)	36.70 (0.65)	0.045	36.66 (0.68)	36.74 (0.72)	0.037
SpO ₂ (%) (mean (SD))	96.76 (3.67)	95.70 (4.92)	0.008	96.99 (3.50)	96.20 (4.14)	<0.001	96.71 (3.74)	96.55 (4.00)	0.459
Urine output (mL) (mean (SD))	1802.33 (1333.75)	1341.22 (981.83)	<0.001	1862.26 (1290.47)	1620.42 (1348.83)	<0.001	1792.70 (1325.29)	1637.70 (1264.41)	0.047
Creatinine (mg/dL) (mean (SD))	1.18 (0.98)	1.08 (0.67)	0.07	1.12 (0.99)	1.27 (0.91)	<0.001	1.19 (1.00)	1.06 (0.72)	0.004
Haemoglobin (g/L) (mean (SD))	101.8 (22.7)	96.0 (20.8)	0.002	103.2 (23.3)	98.5 (21.1)	<0.001	101.5 (22.8)	100.6 (21.5)	0.494
Myocardial infarction (%)	887 (40.25)	62 (38.51)	0.678	588 (39.95)	361 (40.43)	0.829	806 (39.65)	143 (43.07)	0.251
Cerebrovascular disease (%)	298 (13.52)	39 (24.22)	<0.001	194 (13.18)	143 (16.01)	0.06	281 (13.82)	56 (16.87)	0.15
Chronic pulmonary disease (%)	714 (32.40)	53 (32.92)	0.931	435 (29.55)	332 (37.18)	<0.001	650 (31.97)	117 (35.24)	0.255
Diabetes mellitus (%)	1048 (47.55)	67 (41.61)	0.164	670 (45.52)	445 (49.83)	0.046	960 (47.22)	155 (46.69)	0.859
Renal disease (%)	965 (43.79)	81 (50.31)	0.118	564 (38.32)	482 (53.98)	<0.001	892 (43.88)	154 (46.39)	0.404
GCS (mean (SD))	13.15 (2.85)	11.32 (3.71)	<0.001	13.27 (2.88)	12.64 (3.04)	<0.001	13.08 (2.93)	12.73 (3.11)	0.051
SOFA Score (mean (SD))	5.79 (3.47)	6.84 (3.48)	<0.001	5.62 (3.43)	6.28 (3.52)	<0.001	5.82 (3.44)	6.14 (3.70)	0.128
CCI (mean (SD))	7.81 (2.55)	9.25 (2.45)	<0.001	7.27 (2.43)	8.95 (2.45)	<0.001	7.83 (2.53)	8.38 (2.77)	<0.001
APS III (mean (SD))	48.10 (18.86)	60.75 (20.42)	<0.001	45.35 (18.24)	54.92 (19.35)	<0.001	48.33 (18.84)	52.86 (21.06)	<0.001
SAPSII Score (mean (SD))	38.84 (11.65)	45.43 (10.35)	<0.001	37.13 (11.51)	42.86 (11.09)	<0.001	39.09 (11.56)	40.55 (12.35)	0.034

APS III, Acute Physiology Score III; BMI, body mass index; CCI, Charlson Comorbidity Score; GCS, Glasgow Coma Scale; ICU, intensive care unit; MBP, mean blood pressure; SAPSII, Simplified Acute Physiology Score II; SOFA, Sequential Organ Failure Assessment.

Table 3 Outcome analysis by heart rate ≤ 90 bpm vs heart rate > 90 bpm

In-hospital mortality				
Heart rate	Survival	Dead	Total	P value
≤ 90 bpm	1519 (95.84)	66 (4.16)	1585	<0.001
> 90 bpm	685 (87.82)	95 (12.18)	780	
Total	2204 (93.19)	161 (6.81)		
Total mortality				
Heart rate	Survival	Dead	Total	P value
≤ 90 bpm	1021 (64.42)	564 (35.58)	1585	0.002
> 90 bpm	451 (57.82)	329 (42.18)	780	
Total	1472 (62.24)	893 (37.76)		
ICU readmission				
Heart rate	No	Yes	Total	P value
≤ 90 bpm	1404 (88.58)	181 (11.42)	1585	<0.001
> 90 bpm	629 (80.64)	151 (19.36)	780	
Total	2033 (85.96)	332 (14.04)		

bpm, beats per minute; ICU, intensive care unit.

different perspectives on whether heart rate is a predictor or not. Some models suggest a strong impact on prognosis, such as the OPTIMIZE-HF study and another study that uses a GWTG-HF risk score^{6 7 22} and both used heart rate at admission as a predictor. To our knowledge, there has been no report linking ICU discharge heart rate and mortality among HF patients in the ICU. Our study is the first to explore this important clinical issue. In our study, a higher last heart rate before ICU discharge was significantly associated with in-hospital/total mortality and ICU readmission. Heart rate greater than and equal to 90 bpm was strongly associated with increased in-hospital/total mortality and ICU readmission rates.

It is well-established that there is a link between heart rate and mortality. Multiple theories have been proposed to explain the association, including low physical fitness, elevated blood pressure, reduced variability in heart rate and decreased baroreceptor sensitivity in individuals with high heart rate.^{23 24} However, the underlying causes of this correlation have yet to be fully understood.²⁵ The significance of heart rate as a prognostic indicator in acute decompensated heart failure (ADHF) remains a topic of debate. Unlike its predictive role in chronic systolic HF, the role of heart rate in ADHF is much less clear. This is partly due to variations in the timing of heart rate measurement during an acute episode and differences in the outcome measures used in various studies, such as in-hospital mortality and readmission.²⁶ When patients with ADHF presented at the emergency department with a higher heart rate, a higher risk of death within 7 days was linked.²⁷ Patients with ADHF who had a higher heart rate on arriving at the hospital were more likely to have a higher risk of death while in the hospital, with the lowest risk observed at heart rates between 70 and 75 bpm.²⁸

The SHIFT trial has proved that heart rate plays a role in the progression of HF,¹⁰ according to the speculation about a fundamental connection between a higher heart rate and the development²⁹ or worsening of HF,⁸ with theories ranging from the impact on myocardial energy³⁰ to the potential benefits of reducing arterial afterload by reducing heart rate.³¹

Our findings are consistent with a large retrospective cohort study that examined HF patients who had been treated for ADHF. This study found that a higher heart rate at discharge after treatment was linked to a higher risk of death and rehospitalisation, with an even greater risk within the first 30 days following discharge.³² Using heart rate at discharge as a predictor of risk has several benefits, including being routinely available, being easy and reliable to determine, and reflective of the patient's most stable condition, which may be more relevant for future prognosis.

Despite the large sample size and detailed data, this study has several limitations that are common to all retrospective cohort studies. These include missing data, the potential for bias in the data collection and analysis and the inability to account for all possible confounding factors. Given that the data we used originated from an observational database, and furthermore, we lacked detailed information on echocardiography and whether cardiac implantable electronic devices were installed. Due to significant variability in the timing and dosage of pharmacological treatment for HF within existing databases, analysing therapeutic efficacy and establishing causal relationships have been challenging. Therefore, this study focuses on physiological parameters and biochemical markers collected within the first 24 hours of ICU admission as primary variables for analysis. This approach aims to minimise bias arising from inconsistent treatment protocols or clinical judgement, ensuring greater reliability and comparability of the results. The results reported in our study should be regarded only as reference points and must be further verified through additional research.

CONCLUSIONS

Given the association between elevated heart rate at ICU discharge in patients with HF and increasing in-hospital mortality, total mortality and ICU readmission, it is evident that a heart rate exceeding 90 bpm correlates with an increase in these outcomes. It is important to conduct further research to determine if reducing heart rate could be an effective treatment strategy for HF patients discharged from the ICU.

Author affiliations

¹Department of Internal Medicine, Division of Nephrology, Taitung MacKay Memorial Hospital, Taitung, Taiwan

²Department of Medicine, MacKay Medical College, New Taipei, Taiwan

³Department of Internal Medicine, Division of Cardiology, Taitung MacKay Memorial Hospital, Taitung, Taiwan

⁴Graduate Institute of Business Administration, College of Management, National Dong Hwa University, Hualien, Taiwan

⁵Department of Bioinformatics and Medical Engineering, Asia University, Taichung, Taiwan

⁶Department of Cardiology, Asia University Hospital, Taichung, Taiwan

⁷Cardiovascular Center, MacKay Memorial Hospital, Taipei, Taiwan

⁸Department of Artificial Intelligence and Medical Application, MacKay Junior College of Medicine, Nursing, and Management, Taipei, Taiwan

⁹Department of Internal Medicine, Division of Cardiology, Hsinchu MacKay Memorial Hospital, Hsinchu, Taiwan

¹⁰Department of Nursing, MacKay Junior College of Medicine Nursing and Management, Taipei, Taiwan

Acknowledgements Special thanks to Lang-Yin Zeng for her efforts on this article.

Contributors Conceptualisation: C-YH, M-IS, Y-HL, P-LL and W-RC. Methodology: M-IS and Y-CC. Investigation: M-IS, Y-HL and P-LL. Data curation: M-IS and Y-CC. Formal analysis: W-RC. Writing—original draft preparation: W-RC. Writing—review and editing: C-YH, M-IS, Y-CC, Y-HL, P-LL and W-RC. Visualisation: W-RC. Guarantor: M-IS. All authors have read and agreed to the published version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval The accumulation of patient data and the establishment of the research asset were reviewed by the institutional review board at the Beth Israel Deaconess Medical Center, who granted a waiver of informed consent and approved the data sharing initiative

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The data used to support the findings of this study are included within the article and its online supplemental information files.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Wei-Ru Chiou <http://orcid.org/0000-0003-1133-5714>

REFERENCES

- Hoke RS, Müller-Werdan U, Lautenschläger C, *et al.* Heart rate as an independent risk factor in patients with multiple organ dysfunction: a prospective, observational study. *Clin Res Cardiol* 2012;101:139–47.
- Sander O, Welters ID, Foëx P, *et al.* Impact of prolonged elevated heart rate on incidence of major cardiac events in critically ill patients with a high risk of cardiac complications. *Crit Care Med* 2005;33:81–8.
- Chioncel O, Ambrosy AP, Filipescu D, *et al.* Patterns of intensive care unit admissions in patients hospitalized for heart failure. *J Cardiovasc Med (Hagerstown)* 2015;16:331–40.
- Lai J-I, Lin H-Y, Lai Y-C, *et al.* Readmission to the intensive care unit: a population-based approach. *J Formos Med Assoc* 2012;111:504–9.
- Opdahl A, Ambale Venkatesh B, Fernandes VRS, *et al.* Resting heart rate as predictor for left ventricular dysfunction and heart failure: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol* 2014;63:1182–9.
- Abraham WT, Fonarow GC, Albert NM, *et al.* Predictors of in-hospital mortality in patients hospitalized for heart failure: insights from the Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF). *J Am Coll Cardiol* 2008;52:347–56.
- Peterson PN, Rumsfeld JS, Liang L, *et al.* A Validated Risk Score for In-Hospital Mortality in Patients With Heart Failure From the American Heart Association Get With the Guidelines Program. *Circ Cardiovascular Quality and Outcomes* 2010;3:25–32.
- Lechat P, Hulot JS, Escolano S, *et al.* Heart rate and cardiac rhythm relationships with bisoprolol benefit in chronic heart failure in CIBIS II Trial. *Circulation* 2001;103:1428–33.
- Swedberg K, Komajda M, Böhm M, *et al.* Ivabradine and outcomes in chronic heart failure (SHIFT): a randomised placebo-controlled study. *Lancet* 2010;376:875–85.
- Böhm M, Swedberg K, Komajda M, *et al.* Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *Lancet* 2010;376:886–94.
- Liao C-T, Huang J-L, Liang H-W, *et al.* The association between ivabradine and adverse cardiovascular events in acute decompensated HFrEF patients. *ESC Heart Fail* 2021;8:4199–210.
- Lee Y-H, Lin P-L, Chiou W-R, *et al.* Combination of ivabradine and sacubitril/valsartan in patients with heart failure and reduced ejection fraction. *ESC Heart Fail* 2021;8:1204–15.
- Johnson A, Bulgarelli L, Pollard T, *et al.* MIMIC-IV [PhysioNet]. 2020. Available: <https://physionet.org/content/mimiciv/1.0/>
- Goldberger AL, Amaral LA, Glass L, *et al.* PhysioBank, PhysioToolkit, and PhysioNet: components of a new research resource for complex physiologic signals. *Circulation* 2000;101:E215–20.
- Johnson AE, Stone DJ, Celi LA, *et al.* The MIMIC Code Repository: enabling reproducibility in critical care research. *J Am Med Inform Assoc* 2018;25:32–9.
- Erdur H, Scheitz JF, Grittner U, *et al.* Heart rate on admission independently predicts in-hospital mortality in acute ischemic stroke patients. *Int J Cardiol* 2014;176:206–10.
- Lancellotti P, Ancion A, Magne J, *et al.* Elevated heart rate at 24–36h after admission and in-hospital mortality in acute in non-arrhythmic heart failure. *Int J Cardiol* 2015;182:426–30.
- Kara D, Akinci SB, Babaoglu G, *et al.* Increased heart rate on first day in Intensive Care Unit is associated with increased mortality. *Pak J Med Sci* 2016;32:1402–7.
- Grander W, Müllauer K, Koller B, *et al.* Heart rate before ICU discharge: a simple and readily available predictor of short- and long-term mortality from critical illness. *Clin Res Cardiol* 2013;102:599–606.
- Sandfort V, Johnson AEW, Kunz LM, *et al.* Prolonged Elevated Heart Rate and 90-Day Survival in Acutely Ill Patients: Data From the MIMIC-III Database. *J Intensive Care Med* 2019;34:622–9.
- Xu Y, Han D, Huang T, *et al.* Predicting ICU Mortality in Rheumatic Heart Disease: Comparison of XGBoost and Logistic Regression. *Front Cardiovasc Med* 2022;9:847206.
- Yagyu T, Kumada M, Nakagawa T. Novel risk stratification with time course assessment of in-hospital mortality in patients with acute heart failure. *PLoS One* 2017;12:e0187410.
- Zhang D, Shen X, Qi X. Resting heart rate and all-cause and cardiovascular mortality in the general population: a meta-analysis. *CMAJ* 2016;188:E53–63.
- Zhang D, Wang W, Li F. Association between resting heart rate and coronary artery disease, stroke, sudden death and noncardiovascular diseases: a meta-analysis. *CMAJ* 2016;188:E384–92.
- Reunanen A, Karjalainen J, Ristola P, *et al.* Heart rate and mortality. *J Intern Med* 2000;247:231–9.
- Vollmert T, Hellmich M, Gassanov N, *et al.* Heart rate at discharge in patients with acute decompensated heart failure is a predictor of mortality. *Eur J Med Res* 2020;25:47.
- Lee DS, Stiitt A, Austin PC, *et al.* Prediction of heart failure mortality in emergent care: a cohort study. *Ann Intern Med* 2012;156:767–75.
- Bui AL, Grau-Sepulveda MV, Hernandez AF, *et al.* Admission heart rate and in-hospital outcomes in patients hospitalized for heart failure in sinus rhythm and in atrial fibrillation. *Am Heart J* 2013;165:567–74.
- Ho JE, Larson MG, Ghorbani A, *et al.* Long-term cardiovascular risks associated with an elevated heart rate: the Framingham Heart Study. *J Am Heart Assoc* 2014;3:e000668.
- Levine HJ. Optimum heart rate of large failing hearts. *Am J Cardiol* 1988;61:633–6.
- Kelly RP, Ting CT, Yang TM, *et al.* Effective arterial elastance as index of arterial vascular load in humans. *Circulation* 1992;86:513–21.
- Laskey WK, Alomari I, Cox M, *et al.* Heart rate at hospital discharge in patients with heart failure is associated with mortality and rehospitalization. *J Am Heart Assoc* 2015;4:e001626.