

## Distortion of imprecise probabilities

**David Nieto-Barba**  
davidphbarba@outlook.com


**Ignacio Montes**  
imontes@uniovi.es

**Enrique Miranda**  
mirandaenrique@uniovi.es




**20th International Conference on Information Processing and  
Management of Uncertainty in Knowledge-Based Systems**


# Motivation



# Motivation




Given  $P_0 \in \mathbb{P}(\mathcal{X})$ ,  $\delta > 0$ ,  
 $d : \mathbb{P}(\mathcal{X}) \times \mathbb{P}(\mathcal{X}) \rightarrow [0, \infty)$ ,  
 $d$  convex and continuous  
 $\Rightarrow B_d^\delta(P_0) = \mathcal{M}(\underline{P})$




Given  $\underline{P}$ ,  $\delta > 0$ ,

# Motivation



Given  $P_0 \in \mathbb{P}(\mathcal{X})$ ,  $\delta > 0$ ,  
 $d : \mathbb{P}(\mathcal{X}) \times \mathbb{P}(\mathcal{X}) \rightarrow [0, \infty)$ ,  
 $d$  convex and continuous  
 $\Rightarrow B_d^\delta(P_0) = \mathcal{M}(\underline{P})$



Given  $\underline{P}$ ,  $\delta > 0$ ,  
  
?

## ① Preliminaries

## ② Distortion of imprecise models

## ③ Imprecise total variation

## ④ Conclusions

## ① Preliminaries

## ② Distortion of imprecise models

## ③ Imprecise total variation

## ④ Conclusions

## Preliminaries: lower probabilities

- Let  $\mathcal{X} = \{x_1, \dots, x_n\}$  be a *possibility space*.

## Preliminaries: lower probabilities

- Let  $\mathcal{X} = \{x_1, \dots, x_n\}$  be a *possibility space*.
- A function  $\underline{P} : \mathcal{P}(\mathcal{X}) \rightarrow [0, 1]$  is called a ***lower probability*** whenever:

$$\underline{P}(\emptyset) = 0, \quad \underline{P}(\mathcal{X}) = 1 \quad \text{and} \quad A \subseteq B \Rightarrow \underline{P}(A) \leq \underline{P}(B).$$



## Preliminaries: lower probabilities

- Let  $\mathcal{X} = \{x_1, \dots, x_n\}$  be a *possibility space*.
- A function  $\underline{P} : \mathcal{P}(\mathcal{X}) \rightarrow [0, 1]$  is called a ***lower probability*** whenever:

$$\underline{P}(\emptyset) = 0, \quad \underline{P}(\mathcal{X}) = 1 \quad \text{and} \quad A \subseteq B \Rightarrow \underline{P}(A) \leq \underline{P}(B).$$

Also known as (normalised) capacity, monotone game or fuzzy measure.

## Preliminaries: lower probabilities

- Let  $\mathcal{X} = \{x_1, \dots, x_n\}$  be a *possibility space*.
- A function  $\underline{P} : \mathcal{P}(\mathcal{X}) \rightarrow [0, 1]$  is called a **lower probability** whenever:

$$\underline{P}(\emptyset) = 0, \quad \underline{P}(\mathcal{X}) = 1 \quad \text{and} \quad A \subseteq B \Rightarrow \underline{P}(A) \leq \underline{P}(B).$$

Also known as (normalised) capacity, monotone game or fuzzy measure.

- It has an associated **credal set**:

$$\mathcal{M}(\underline{P}) = \{P \in \mathbb{P}(\mathcal{X}) : P(A) \geq \underline{P}(A), \forall A \subseteq \mathcal{X}\}.$$

## Preliminaries: lower probabilities

- Let  $\mathcal{X} = \{x_1, \dots, x_n\}$  be a *possibility space*.
- A function  $\underline{P} : \mathcal{P}(\mathcal{X}) \rightarrow [0, 1]$  is called a **lower probability** whenever:

$$\underline{P}(\emptyset) = 0, \quad \underline{P}(\mathcal{X}) = 1 \quad \text{and} \quad A \subseteq B \Rightarrow \underline{P}(A) \leq \underline{P}(B).$$

Also known as (normalised) capacity, monotone game or fuzzy measure.

- It has an associated **credal set**:

$$\mathcal{M}(\underline{P}) = \{P \in \mathbb{P}(\mathcal{X}) : P(A) \geq \underline{P}(A), \forall A \subseteq \mathcal{X}\}.$$

- A lower probability is said to be **coherent** if:

$$\underline{P}(A) = \min\{P(A) : P \in \mathcal{M}(\underline{P})\}, \quad \forall A \subseteq \mathcal{X}.$$

## Preliminaries: lower probabilities

Some particular properties that a coherent lower probability may satisfy are:

❶ **2-monotonicity.** For every  $A, B \subseteq \mathcal{X}$ :

$$\underline{P}(A \cup B) + \underline{P}(A \cap B) \geq \underline{P}(A) + \underline{P}(B).$$

## Preliminaries: lower probabilities

Some particular properties that a coherent lower probability may satisfy are:

❶ **2-monotonicity.** For every  $A, B \subseteq \mathcal{X}$ :

$$\underline{P}(A \cup B) + \underline{P}(A \cap B) \geq \underline{P}(A) + \underline{P}(B).$$

❷  **$k$ -monotonicity.** For every  $A_1, \dots, A_p \subseteq \mathcal{X}$  and  $1 \leq p \leq k$ :

$$\underline{P}\left(\bigcup_{i=1}^p A_i\right) \geq \sum_{I \subseteq \{1, \dots, p\}} (-1)^{|I|+1} \underline{P}\left(\bigcap_{i \in I} A_i\right).$$

## Preliminaries: lower probabilities

Some particular properties that a coherent lower probability may satisfy are:

❶ **2-monotonicity.** For every  $A, B \subseteq \mathcal{X}$ :

$$\underline{P}(A \cup B) + \underline{P}(A \cap B) \geq \underline{P}(A) + \underline{P}(B).$$

❷  **$k$ -monotonicity.** For every  $A_1, \dots, A_p \subseteq \mathcal{X}$  and  $1 \leq p \leq k$ :

$$\underline{P}\left(\bigcup_{i=1}^p A_i\right) \geq \sum_{I \subseteq \{1, \dots, p\}} (-1)^{|I|+1} \underline{P}\left(\bigcap_{i \in I} A_i\right).$$

❸ **minitivity.** For every  $A, B \subseteq \mathcal{X}$ :

$$\underline{P}(A \cap B) = \min\{\underline{P}(A), \underline{P}(B)\}.$$

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.



## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.
- It is said to be **coherent** if:

$$\underline{P}(f) = \min_{P \in \mathcal{M}(\underline{P})} E_P(f) \quad \forall f \in \mathcal{L}(\mathcal{X}),$$

where  $E_P$  denotes the expectation operator w.r.t.  $P \in \mathbb{P}(\mathcal{X})$

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.
- It is said to be **coherent** if:

$$\underline{P}(f) = \min_{P \in \mathcal{M}(\underline{P})} E_P(f) \quad \forall f \in \mathcal{L}(\mathcal{X}),$$

where  $E_P$  denotes the expectation operator w.r.t.  $P \in \mathbb{P}(\mathcal{X})$  and  $\mathcal{M}(\underline{P}) := \{P \in \mathbb{P}(\mathcal{X}) : E_P(f) \geq \underline{P}(f) \forall f \in \mathcal{L}(\mathcal{X})\}$ .

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.
- It is said to be **coherent** if:

$$\underline{P}(f) = \min_{P \in \mathcal{M}(\underline{P})} E_P(f) \quad \forall f \in \mathcal{L}(\mathcal{X}),$$

where  $E_P$  denotes the expectation operator w.r.t.  $P \in \mathbb{P}(\mathcal{X})$  and  $\mathcal{M}(\underline{P}) := \{P \in \mathbb{P}(\mathcal{X}) : E_P(f) \geq \underline{P}(f) \forall f \in \mathcal{L}(\mathcal{X})\}$ .

- It is said to be **2-monotone** if:

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.
- It is said to be **coherent** if:

$$\underline{P}(f) = \min_{P \in \mathcal{M}(\underline{P})} E_P(f) \quad \forall f \in \mathcal{L}(\mathcal{X}),$$

where  $E_P$  denotes the expectation operator w.r.t.  $P \in \mathbb{P}(\mathcal{X})$  and  $\mathcal{M}(\underline{P}) := \{P \in \mathbb{P}(\mathcal{X}) : E_P(f) \geq \underline{P}(f) \forall f \in \mathcal{L}(\mathcal{X})\}$ .

- It is said to be **2-monotone** if:

$$\underline{P}(f \wedge g) + \underline{P}(f \vee g) \geq \underline{P}(f) + \underline{P}(g).$$

## Preliminaries: lower previsions

- Let  $\mathcal{L}(\mathcal{X})$  be the set of real-valued functionals on  $\mathcal{X}$  or *gambles*.
- Any function  $\underline{P} : \mathcal{L}(\mathcal{X}) \rightarrow \mathbb{R}$  is called *lower prevision*.
- It is said to be **coherent** if:

$$\underline{P}(f) = \min_{P \in \mathcal{M}(\underline{P})} E_P(f) \quad \forall f \in \mathcal{L}(\mathcal{X}),$$

where  $E_P$  denotes the expectation operator w.r.t.  $P \in \mathbb{P}(\mathcal{X})$  and  $\mathcal{M}(\underline{P}) := \{P \in \mathbb{P}(\mathcal{X}) : E_P(f) \geq \underline{P}(f) \forall f \in \mathcal{L}(\mathcal{X})\}$ .

- It is said to be **2-monotone** if:

$$\underline{P}(f \wedge g) + \underline{P}(f \vee g) \geq \underline{P}(f) + \underline{P}(g).$$

In this case, lower probabilities  $\leftrightarrow$  lower previsions (Choquet integral).

① Preliminaries

② Distortion of imprecise models

③ Imprecise total variation

④ Conclusions

# Distortion procedures

	$P_0 \in \mathbb{P}(\mathcal{X})$	$\underline{P}$
(a)	$B_d^\delta(P_0) = \{Q \in \mathbb{P}(\mathcal{X}) : d(Q, P_0) \leq \delta\}$ lower $\downarrow$ envelope $\underline{Q}(A) = \inf\{P(A) : P \in B_d^\delta(P_0)\}$	$\bigcup_{P \in \mathcal{M}(\underline{P})} B_d^\delta(P)$ lower $\downarrow$ envelope $\underline{Q}(A)$

# Distortion procedures

	$P_0 \in \mathbb{P}(\mathcal{X})$	$\underline{P}$
(a)	$B_d^\delta(P_0) = \{Q \in \mathbb{P}(\mathcal{X}) : d(Q, P_0) \leq \delta\}$ lower $\downarrow$ envelope $\underline{Q}(A) = \inf\{P(A) : P \in B_d^\delta(P_0)\}$	$\bigcup_{P \in \mathcal{M}(\underline{P})} B_d^\delta(P)$ lower $\downarrow$ envelope $\underline{Q}(A)$
(b)	$f_\delta : [0, 1] \rightarrow [0, 1]$ increasing, $f_\delta(0) = 0, f_\delta(1) = 1$ $\downarrow$ $\underline{Q}_\delta(A) := f_\delta(P_0(A))$	$f_\delta : [0, 1] \rightarrow [0, 1]$ increasing, $f_\delta(0) = 0, f_\delta(1) = 1$ $\downarrow$ $\underline{Q}_\delta(A) := f_\delta(\underline{P}(A))$



# Desirable properties of distortion procedures

❶ *Expansion*. Given  $\delta_1 > \delta_2 > 0$ :

$$\underline{Q}_{\delta_1}(\underline{P}) \leq \underline{Q}_{\delta_2}(\underline{P}).$$

# Desirable properties of distortion procedures

❶ *Expansion*. Given  $\delta_1 > \delta_2 > 0$ :

$$\underline{Q}_{\delta_1}(\underline{P}) \leq \underline{Q}_{\delta_2}(\underline{P}).$$

❷ *Aggregation*. For any  $\delta_1, \delta_2 > 0$ :

$$\underline{Q}_{\delta_2 + \delta_1}(\underline{P}) = \underline{Q}_{\delta_2}(\underline{Q}_{\delta_1}(\underline{P})).$$

# Desirable properties of distortion procedures

❶ *Expansion*. Given  $\delta_1 > \delta_2 > 0$ :

$$\underline{Q}_{\delta_1}(\underline{P}) \leq \underline{Q}_{\delta_2}(\underline{P}).$$

❷ *Aggregation*. For any  $\delta_1, \delta_2 > 0$ :

$$\underline{Q}_{\delta_2 + \delta_1}(\underline{P}) = \underline{Q}_{\delta_2}(\underline{Q}_{\delta_1}(\underline{P})).$$

❸ *Structure preservation*. For every  $\delta > 0$ :

$\underline{P}$  coherent (resp. 2-monotone,  $k$ -monotone, minitive)  $\Rightarrow$  so is  $\underline{Q}_{\delta}(\underline{P})$ .

# Desirable properties of distortion procedures

❶ **Expansion.** Given  $\delta_1 > \delta_2 > 0$ :

$$\underline{Q}_{\delta_1}(\underline{P}) \leq \underline{Q}_{\delta_2}(\underline{P}).$$

❷ **Aggregation.** For any  $\delta_1, \delta_2 > 0$ :

$$\underline{Q}_{\delta_2 + \delta_1}(\underline{P}) = \underline{Q}_{\delta_2}(\underline{Q}_{\delta_1}(\underline{P})).$$

❸ **Structure preservation.** For every  $\delta > 0$ :

$\underline{P}$  coherent (resp. 2-monotone,  $k$ -monotone, minitive)  $\Rightarrow$  so is  $\underline{Q}_{\delta}(\underline{P})$ .

❹ **Commutativity.** If  $\underline{P}$  is coherent, then for any  $\delta > 0$ :

$$\mathcal{M}(\underline{Q}_{\delta}(\underline{P})) = \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_{\delta}(P)).$$

## ① Preliminaries

## ② Distortion of imprecise models

## ③ Imprecise total variation

## ④ Conclusions

# Total Variation distance

## Definition (Total Variation distance)

$d_{TV} : \mathbb{P}(\mathcal{X}) \times \mathbb{P}(\mathcal{X}) \rightarrow [0, \infty)$  given by:

$$d_{TV}(P, Q) = \max_{A \subseteq \mathcal{X}} |P(A) - Q(A)| \quad \forall P, Q \in \mathbb{P}(\mathcal{X}).$$

# Total Variation distance

## Definition (Total Variation distance)

$d_{TV} : \mathbb{P}(\mathcal{X}) \times \mathbb{P}(\mathcal{X}) \rightarrow [0, \infty)$  given by:

$$d_{TV}(P, Q) = \max_{A \subseteq \mathcal{X}} |P(A) - Q(A)| \quad \forall P, Q \in \mathbb{P}(\mathcal{X}).$$

## Proposition (Herron et al. '97)

The lower envelope of  $B_{d_{TV}}^\delta(P)$  is given by:

$$\underline{Q}_P(A) = \max\{P(A) - \delta, 0\} \quad \forall A \subset \mathcal{X}, \quad \underline{Q}_P(\mathcal{X}) = 1.$$

# Imprecise Total Variation (first approach)

## Definition (Imprecise Total Variation)

Let  $\underline{P}$  be a coherent lower probability and  $\delta > 0$ . The **imprecise total variation model** induced by  $(\underline{P}, \delta)$  is the lower probability  $\underline{Q}$  given by

$$\underline{Q}(A) = \max \{ \underline{P}(A) - \delta, 0 \} \quad \forall A \subset \mathcal{X}, \quad \underline{Q}(\mathcal{X}) = 1. \quad (\text{ITV1})$$



# Imprecise Total Variation (first approach)

## Definition (Imprecise Total Variation)

Let  $\underline{P}$  be a coherent lower probability and  $\delta > 0$ . The **imprecise total variation model** induced by  $(\underline{P}, \delta)$  is the lower probability  $\underline{Q}$  given by

$$\underline{Q}(A) = \max \{ \underline{P}(A) - \delta, 0 \} \quad \forall A \subset \mathcal{X}, \quad \underline{Q}(\mathcal{X}) = 1. \quad (\text{ITV1})$$

This generalises the eponymous model for the precise case, and has also appeared as the *strong*  $\delta$ -core in game theory (used to find a solution of a distorted game when  $\mathcal{M}(\underline{P}) = \emptyset$ ).

## Proposition (Expansion and Aggregation)

*The imprecise total variation (ITV1) satisfies expansion and aggregation.*

## Proposition (Expansion and Aggregation)

*The imprecise total variation (ITV1) satisfies expansion and aggregation.*

## Proposition (Structure preservation)

*Let  $\underline{P}$  be a lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the total variation model they induce. If  $\underline{P}$  has any of the following properties:*

- ❶ avoiding sure loss (i.e.  $\mathcal{M}(\underline{P}) \neq \emptyset$ ),*
- ❷ coherence,*
- ❸ 2-monotonicity,*
- ❹ minitivity,*

*so does  $\underline{Q}$ .*

## Proposition (Expansion and Aggregation)

*The imprecise total variation (ITV1) satisfies expansion and aggregation.*

## Proposition (Structure preservation)

*Let  $\underline{P}$  be a lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the total variation model they induce. If  $\underline{P}$  has any of the following properties:*

- i avoiding sure loss (i.e.  $\mathcal{M}(\underline{P}) \neq \emptyset$ ),*
- ii coherence,*
- iii 2-monotonicity,*
- iv minitivity,*

*so does  $\underline{Q}$ .*

However, k-monotonicity is not preserved in general.

# Imprecise Total Variation (second approach)

What about commutativity?

Consider:

$$\underline{Q}'(A) = \min \left\{ Q(A) : Q \in \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_P) \right\}, \quad \forall A \subseteq \mathcal{X}. \quad (\text{ITV2})$$

# Imprecise Total Variation (second approach)

What about commutativity?

Consider:

$$\underline{Q}'(A) = \min \left\{ Q(A) : Q \in \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_P) \right\}, \quad \forall A \subseteq \mathcal{X}. \quad (\text{ITV2})$$

## Proposition

*Let  $\underline{P}$  be a coherent lower probability,  $\delta > 0$  and  $\underline{Q}, \underline{Q}'$  the lower probabilities induced by Eqs. (ITV1), (ITV2). Then  $\underline{Q}' = \underline{Q}$ .*

Consider the premetric:

$$d_{\text{TV}}^{\min}(\underline{P}, \underline{Q}) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} d_{\text{TV}}(P, Q) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} \max_{A \subseteq \mathcal{X}} |P(A) - Q(A)|.$$

Consider the premetric:

$$d_{\text{TV}}^{\min}(\underline{P}, \underline{Q}) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} d_{\text{TV}}(P, Q) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} \max_{A \subseteq \mathcal{X}} |P(A) - Q(A)|.$$

## Proposition

Let  $\underline{P}$  be a coherent lower probability and  $\delta > 0$ . The following set is closed and convex:

$$B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P}) = \{Q \in \mathbb{P}(\mathcal{X}) : d_{\text{TV}}^{\min}(Q, \underline{P}) \leq \delta\}.$$



Consider the premetric:

$$d_{\text{TV}}^{\min}(\underline{P}, \underline{Q}) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} d_{\text{TV}}(P, Q) = \min_{\substack{P \in \mathcal{M}(\underline{P}) \\ Q \in \mathcal{M}(\underline{Q})}} \max_{A \subseteq \mathcal{X}} |P(A) - Q(A)|.$$

## Proposition

Let  $\underline{P}$  be a coherent lower probability and  $\delta > 0$ . The following set is closed and convex:

$$B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P}) = \{Q \in \mathbb{P}(\mathcal{X}) : d_{\text{TV}}^{\min}(Q, \underline{P}) \leq \delta\}.$$

It holds:

$$B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P}) = \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_P).$$

Is  $\mathcal{M}(\underline{Q})$  equal to  $B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P}) = \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_P)$  ?

Is  $\mathcal{M}(\underline{Q})$  equal to  $B_{d_{\text{TV}}}^{\delta}(\underline{P}) = \bigcup_{P \in \mathcal{M}(\underline{P})} \mathcal{M}(\underline{Q}_P)$  ?

## Example

$A$	$\{x_1\}$	$\{x_2\}$	$\{x_3\}$	$\{x_4\}$	$\{x_5\}$	$\{x_6\}$
$P_1(A)$	0.2	0.061	0.061	0.19	0.199	0.289
$P_2(A)$	0.161	0.1	0.1	0.238	0.141	0.26
$P_3(A)$	0.239	0.099	0.12	0.161	0.17	0.211
$P_4(A)$	0.161	0.177	0.13	0.161	0.102	0.269
$P_5(A)$	0.22	0.041	0.109	0.21	0.199	0.221
$P_6(A)$	0.22	0.041	0.041	0.23	0.199	0.269
$P_7(A)$	0.178	0.16	0.111	0.161	0.1	0.29
$P_8(A)$	0.161	0.157	0.14	0.181	0.073	0.288
$P_9(A)$	0.19	0.073	0.078	0.21	0.15	0.299

## Example (Continuation)

$A$	$\{x_1\}$	$\{x_2\}$	$\{x_3\}$	$\{x_4\}$	$\{x_5\}$	$\{x_6\}$
$P_{10}(A)$	0.161	0.158	0.159	0.161	0.112	0.249
$P_{11}(A)$	0.2	0.071	0.14	0.199	0.199	0.191
$P_{12}(A)$	0.161	0.177	0.138	0.161	0.073	0.29
$P_{13}(A)$	0.239	0.071	0.1	0.189	0.199	0.202
$P_{14}(A)$	0.22	0.041	0.081	0.238	0.199	0.221
$P_{15}(A)$	0.219	0.119	0.061	0.161	0.17	0.27
$P_{16}(A)$	0.218	0.043	0.062	0.238	0.168	0.271
$P_{17}(A)$	0.2	0.0705	0.1585	0.1805	0.199	0.1915

## Example (Continuation)

$A$	$\{x_1\}$	$\{x_2\}$	$\{x_3\}$	$\{x_4\}$	$\{x_5\}$	$\{x_6\}$
$P_{10}(A)$	0.161	0.158	0.159	0.161	0.112	0.249
$P_{11}(A)$	0.2	0.071	0.14	0.199	0.199	0.191
$P_{12}(A)$	0.161	0.177	0.138	0.161	0.073	0.29
$P_{13}(A)$	0.239	0.071	0.1	0.189	0.199	0.202
$P_{14}(A)$	0.22	0.041	0.081	0.238	0.199	0.221
$P_{15}(A)$	0.219	0.119	0.061	0.161	0.17	0.27
$P_{16}(A)$	0.218	0.043	0.062	0.238	0.168	0.271
$P_{17}(A)$	0.2	0.0705	0.1585	0.1805	0.199	0.1915

$$Q := (0.2, 0.05, 0.05, 0.201, 0.199, 0.3) \in \mathcal{M}(\underline{Q}) \setminus B_{d_{\text{TV}}}^{\delta}(\underline{P}),$$

where  $\underline{P}(A) = \min\{P_i(A) : i = 1, \dots, 17\}$  for any  $A \subseteq \mathcal{X}$ ,  $\delta = 0.011$ .

## Proposition

*Let  $\underline{P}$  be a 2-monotone lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the lower probability defined by (ITV1). Then,  $\mathcal{M}(\underline{Q}) = B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P})$ .*

## Proposition

Let  $\underline{P}$  be a 2-monotone lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the lower probability defined by (ITV1). Then,  $\mathcal{M}(\underline{Q}) = B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P})$ .

## Proof.

Let  $\mathcal{H} \subseteq \mathcal{L}(\mathcal{X})$  be the set of gambles that take values in  $[0, 1]$ .

## Proposition

Let  $\underline{P}$  be a 2-monotone lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the lower probability defined by (ITV1). Then,  $\mathcal{M}(\underline{Q}) = B_{d_{\text{TV}}^{\min}}^{\delta}(\underline{P})$ .

## Proof.

Let  $\mathcal{H} \subseteq \mathcal{L}(\mathcal{X})$  be the set of gambles that take values in  $[0, 1]$ . Define, for a given  $Q \in \mathbb{P}(\mathcal{X})$ ,  $f_Q(P, g) := P(g) - Q(g)$  over  $\mathcal{M}(\underline{P}) \times \mathcal{H}$ .



## Proposition

Let  $\underline{P}$  be a 2-monotone lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the lower probability defined by (ITV1). Then,  $\mathcal{M}(\underline{Q}) = B_{d_{\text{TV}}}^{\delta}(\underline{P})$ .

## Proof.

Let  $\mathcal{H} \subseteq \mathcal{L}(\mathcal{X})$  be the set of gambles that take values in  $[0, 1]$ . Define, for a given  $Q \in \mathbb{P}(\mathcal{X})$ ,  $f_Q(P, g) := P(g) - Q(g)$  over  $\mathcal{M}(\underline{P}) \times \mathcal{H}$ .

$$\max_{\mathcal{H}} \min_{\mathcal{M}(\underline{P})} f_Q(P, g) \leq \delta \Leftrightarrow Q \in \mathcal{M}(\underline{Q}) \quad \text{and}$$

$$\min_{\mathcal{M}(\underline{P})} \max_{\mathcal{H}} f_Q(P, g) \leq \delta \Leftrightarrow Q \in B_{d_{\text{TV}}}^{\delta}(\underline{P}).$$

## Proposition

Let  $\underline{P}$  be a 2-monotone lower probability,  $\delta > 0$  and let  $\underline{Q}$  be the lower probability defined by (ITV1). Then,  $\mathcal{M}(\underline{Q}) = B_{d_{\text{TV}}}^{\delta}(\underline{P})$ .

## Proof.

Let  $\mathcal{H} \subseteq \mathcal{L}(\mathcal{X})$  be the set of gambles that take values in  $[0, 1]$ . Define, for a given  $Q \in \mathbb{P}(\mathcal{X})$ ,  $f_Q(P, g) := P(g) - Q(g)$  over  $\mathcal{M}(\underline{P}) \times \mathcal{H}$ .

$$\max_{\mathcal{H}} \min_{\mathcal{M}(\underline{P})} f_Q(P, g) \leq \delta \Leftrightarrow Q \in \mathcal{M}(\underline{Q}) \quad \text{and}$$

$$\min_{\mathcal{M}(\underline{P})} \max_{\mathcal{H}} f_Q(P, g) \leq \delta \Leftrightarrow Q \in B_{d_{\text{TV}}}^{\delta}(\underline{P}).$$

Applying the min-max theorem,

$$\min_{\mathcal{M}(\underline{P})} \max_{\mathcal{H}} f_Q(P, g) = \max_{\mathcal{H}} \min_{\mathcal{M}(\underline{P})} f_Q(P, g). \quad \square$$

## ① Preliminaries

## ② Distortion of imprecise models

## ③ Imprecise total variation


## ④ Conclusions


# Synthesis

Properties	Imprecise TV
Expansion	✓
Aggregation	✓
Structure Preservation	

# Synthesis

Properties	Imprecise TV
Expansion	✓
Aggregation	✓
Structure Preservation (ASL, coherence, 2-monot., minitivity)	✓
Structure Preservation (k-monotonicity)	X
Commutativity	$\sim$ (✓ under 2-monot.)





The dashed line expresses the partial correspondence, in this case, under 2-monotonicity.

## Future lines

- Extensions of ‘classical’ distortion models to the imprecise setting.



# Future lines

- Extensions of ‘classical’ distortion models to the imprecise setting.
- Analysis of other desirable properties (e.g. Destercke, Moral) such as invariance under permutations on  $\mathcal{X}$  and under marginalization.

# Future lines

- Extensions of 'classical' distortion models to the imprecise setting.
- Analysis of other desirable properties (e.g. Destercke, Moral) such as invariance under permutations on  $\mathcal{X}$  and under marginalization.
- Distortion of other IP models (e.g. Moral), and also through different premetrics (e.g. maximum instead of minimum).

# Future lines

- Extensions of 'classical' distortion models to the imprecise setting.
- Analysis of other desirable properties (e.g. Destercke, Moral) such as invariance under permutations on  $\mathcal{X}$  and under marginalization.
- Distortion of other IP models (e.g. Moral), and also through different premetrics (e.g. maximum instead of minimum).
- Connections with game theory (e.g. weak  $\delta$ -core, through a penalised version of  $d_{TV}$ ).

# References



Destercke, S.: A new contextual discounting rule for lower probabilities. In: Hüllermeier, E., Kruse, R., Hoffmann, F. (eds.) Proceedings of IPMU'2010, Part II. pp. 198–207. Communications in Computer and Information Science, 81 (2010)



Herron, T., Seidenfeld, T., Wasserman, L.: Divisive conditioning: further results on dilation. *Philosophy of Science* **64**, 411–444 (1997)



Montes, I., Miranda, E., Destercke, S.: Unifying neighbourhood and distortion models: Part I- New results on old models. *International Journal of General Systems* **49**(6), 602–635 (2020)



Moral, S.: Discounting imprecise probabilities. In: Gil, E., Gil, E., Gil, J., Gil, M. (eds.) *The Mathematics of the Uncertain, Studies in Systems, Decision and Control*, vol. 142. Springer (2018)



Walley, P.: *Statistical Reasoning with Imprecise Probabilities*. Chapman and Hall, London (1991)

# Funding



Proyecto **PID2022-140585NB-I00** financiado por MICIU/AEI/10.13039/501100011033 y por FEDER, UE