

Infimal-convolution-type regularization for inverse problems in imaging

Kristian Bredies

Institute for Mathematics and Scientific Computing
University of Graz

IHP semester “The Mathematics of Imaging”
Workshop “Variational methods and optimization in imaging”
February 7th, 2019

Motivation

Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics (e.g. cartoon/texture)
- Apply broadly to imaging problems (denoising, inpainting, inverse problems/reconstruction)

Motivation


Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics
(e.g. cartoon/texture)
- Apply broadly to imaging
problems (denoising, inpainting,
inverse problems/reconstruction)

Motivation




Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics
(e.g. cartoon/texture)
- Apply broadly to imaging
problems (denoising, inpainting,
inverse problems/reconstruction)

Motivation





Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics (e.g. cartoon/textture)
- Apply broadly to imaging problems (denoising, inpainting, inverse problems/reconstruction)

Motivation





Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics (e.g. cartoon/textture)
- Apply broadly to imaging problems (denoising, inpainting, inverse problems/reconstruction)

Motivation



Goals:

- Find flexible models for stable solution of inverse problems in function spaces
- Establish meaningful combined models

In particular:

- Combine multiple smoothness orders
- Separate image characteristics (e.g. cartoon/textture)
- Apply broadly to imaging problems (denoising, inpainting, inverse problems/reconstruction)

Here: Infimal-convolution regularizers

Outline

- 1 Infimal convolution regularization
- 2 Total generalized variation
 - Definition and properties
 - Applications
- 3 Infimal convolution TGV
 - Accelerated dynamic MRI
- 4 Infimal convolution of oscillation TGV
 - Oscillation TGV and infimal convolution
 - Numerical realization
 - Applications
- 5 Summary

Outline

- 1 Infimal convolution regularization
- 2 Total generalized variation
 - Definition and properties
 - Applications
- 3 Infimal convolution TGV
 - Accelerated dynamic MRI
- 4 Infimal convolution of oscillation TGV
 - Oscillation TGV and infimal convolution
 - Numerical realization
 - Applications
- 5 Summary

The infimal convolution

Definition:

- $\Phi_1, \Phi_2 : X \rightarrow]-\infty, \infty]$ proper, convex, l.s.c.

$$(\Phi_1 \square \Phi_2)(u) = \inf_{u_1 + u_2 = u} \Phi_1(u_1) + \Phi_2(u_2)$$

- Infimal convolution of Φ_1 and Φ_2
- $\Phi_1 \square \Phi_2$ is exact if the infimum is always attained

The infimal convolution

Definition:

- $\Phi_1, \Phi_2 : X \rightarrow]-\infty, \infty]$ proper, convex, l.s.c.

$$(\Phi_1 \square \Phi_2)(u) = \inf_{u_1 + u_2 = u} \Phi_1(u_1) + \Phi_2(u_2)$$

- Infimal convolution of Φ_1 and Φ_2
- $\Phi_1 \square \Phi_2$ is exact if the infimum is always attained

m-fold infimal convolution:

$$(\Phi_1 \square \dots \square \Phi_m)(u) = \inf_{u_1 + \dots + u_m = u} \sum_{i=1}^m \Phi_i(u_i)$$

The infimal convolution

Definition:

- $\Phi_1, \Phi_2 : X \rightarrow]-\infty, \infty]$ proper, convex, l.s.c.

$$(\Phi_1 \square \Phi_2)(u) = \inf_{u_1 + u_2 = u} \Phi_1(u_1) + \Phi_2(u_2)$$


- Infimal convolution of Φ_1 and Φ_2
- $\Phi_1 \square \Phi_2$ is exact if the infimum is always attained

m-fold infimal convolution:

$$(\Phi_1 \square \dots \square \Phi_m)(u) = \inf_{u_1 + \dots + u_m = u} \sum_{i=1}^m \Phi_i(u_i)$$

- Optimal balancing of m competing regularization functionals

Basic variational models



Smoothness-based:


- Total variation

$$\Phi(u) = \int_{\Omega} d|\nabla u|$$

[Rudin/Osher/Fatemi '92]

- Accounts for edges
- Unaware of higher-order smoothness
 \rightsquigarrow staircasing effect

Basic variational models



Smoothness-based:

- Total variation

$$\Phi(u) = \int_{\Omega} d|\nabla u|$$

[Rudin/Osher/Fatemi '92]

- Accounts for edges
- Unaware of higher-order smoothness
~~> staircasing effect

- Higher-order TV


$$\Phi(u) = \int_{\Omega} d|\nabla^2 u|$$

[Lysaker/Lundervold/Tai '03]

[Hinterberger/Scherzer '04]

- Favors smooth solutions
- Edges are not preserved

Infimal-convolution models



Smoothness-based:


- TV-TV² infimal convolution

$$\Phi(u) = \min_{u=u_1+u_2} \int_{\Omega} d|\nabla u_1| + \beta \int_{\Omega} d|\nabla^2 u_2|$$

[Chambolle/Lions '97]

- Models piecewise smooth images
- Staircase effect dominates solutions

Infimal-convolution models



Smoothness-based:

- TV-TV² infimal convolution

$$\Phi(u) = \min_{u=u_1+u_2} \int_{\Omega} d|\nabla u_1| + \beta \int_{\Omega} d|\nabla^2 u_2|$$

[Chambolle/Lions '97]

- Models piecewise smooth images
- Staircase effect dominates solutions


- TV-L¹ \circ Δ infimal convolution

$$\Phi(u) = \min_{u=u_1+u_2} \int_{\Omega} d|\nabla u_1| + \beta \int_{\Omega} d|\Delta u_2|$$

[Chan/Esedoglu/Park '05]

- Denoising similar to TV-TV² infimal convolution

Infimal-convolution-type models



Variations:


- Total generalized variation of order 2

$$\begin{aligned}\Phi(u) &= \text{TGV}_\alpha^2(u) \\ &= \min_{\nabla u = u_1 + u_2} \alpha_1 \int_{\Omega} d|u_1| + \alpha_0 \int_{\Omega} d|\mathcal{E}u_2|\end{aligned}$$

[B./Kunisch/Pock '10]

- Models piecewise smooth images
- Favors smoothness where given
- Preserves relevant edges

Infimal-convolution-type models



Variations:

- Total generalized variation of order 2


$$\begin{aligned}\Phi(u) &= \text{TGV}_\alpha^2(u) \\ &= \min_{\nabla u = u_1 + u_2} \alpha_1 \int_{\Omega} d|u_1| + \alpha_0 \int_{\Omega} d|\mathcal{E}u_2|\end{aligned}$$

[B./Kunisch/Pock '10]

- Models piecewise smooth images
- Favors smoothness where given
- Preserves relevant edges
- Other variants by

[Setzer/Steidl/Teuber '11]

Cartoon/texture decomposition



Infimal convolution models:

- TV- G -norm decomposition

$$\Phi(u) = \min_{u_1+u_2=u} \int_{\Omega} d|\nabla u_1| + \beta \|u_2\|_{TV^*}$$

[Osher/Vese '03], [Aujol et al. '05]


- TV- H^{-1} -norm decomposition

$$\Phi(u) = \min_{u_1+u_2=u} \int_{\Omega} d|\nabla u_1| + \beta \|u_2\|_{H^{-1}}$$

[Osher/Sole/Vese '03]

- Separates piecewise smooth and oscillatory parts
- G -norm/ H^{-1} -norm part also captures non-oscillations

Cartoon/texture decomposition



Infimal convolution models:

- TV- G -norm decomposition

$$\Phi(u) = \min_{u_1+u_2=u} \int_{\Omega} d|\nabla u_1| + \beta \|u_2\|_{TV^*}$$

[Osher/Vese '03], [Aujol et al. '05]

- TV- H^{-1} -norm decomposition

$$\Phi(u) = \min_{u_1+u_2=u} \int_{\Omega} d|\nabla u_1| + \beta \|u_2\|_{H^{-1}}$$

[Osher/Sole/Vese '03]

- Separates piecewise smooth and oscillatory parts
- G -norm/ H^{-1} -norm part also captures non-oscillations

Infimal convolution for inverse problems

Tikhonov regularization:

Solve

$$\min_{u \in X} S(Ku, f) + \Phi(u)$$

- S discrepancy for $Ku = f$, Φ regularization functional

Infimal convolution for inverse problems

Tikhonov regularization:

Solve

$$\min_{u \in X} S(Ku, f) + \Phi(u)$$

- S discrepancy for $Ku = f$, Φ regularization functional

Two viewpoints:

- 1 Monolithic regularization

$$\min_{u \in X} S(Ku, f) + \Phi(u), \quad \Phi = \Phi_1 \square \dots \square \Phi_m$$

- 2 Vector-valued regularization

$$\min_{(u_1, \dots, u_m) \in X^m} S(K(u_1 + \dots + u_m), f) + \Phi_1(u_1) + \dots + \Phi_m(u_m)$$

Infimal convolution for inverse problems

Tikhonov regularization:

Solve

$$\min_{u \in X} S(Ku, f) + \Phi(u)$$

- S discrepancy for $Ku = f$, Φ regularization functional

Two viewpoints:

1 Monolithic regularization

$$\min_{u \in X} S(Ku, f) + \Phi(u), \quad \Phi = \Phi_1 \square \dots \square \Phi_m$$

2 Vector-valued regularization

$$\min_{(u_1, \dots, u_m) \in X^m} S(K(u_1 + \dots + u_m), f) + \Phi_1(u_1) + \dots + \Phi_m(u_m)$$

Monolithic regularization:

- Intrinsic decomposition & problem-adapted interpretation

General existence result

Theorem:

- X reflexive Banach space, Y normed space
- $K : X \rightarrow Y$ bounded linear
- $S(\cdot, f) : Y \rightarrow [0, \infty]$ proper, convex, l.s.c.
- $\Phi : X \rightarrow [0, \infty]$ proper, convex, l.s.c.
- Φ 1-homogeneous, $\dim \ker(\Phi) < \infty$, and coercive, i.e.,

$$\|u - Pu\|_X \leq C\Phi(u)$$

for $P : X \rightarrow \ker(\Phi)$ bounded linear projector

- Then:
$$\min_{u \in X} S(Ku, f) + \alpha\Phi(u)$$

has a solution for each $\alpha > 0$

General regularization properties

$$\min_{u \in X} S(Ku, f) + \alpha \Phi(u)$$

- Let: Y Hilbert space, $S(v, f) = \frac{1}{2} \|v - f\|_Y^2$

Then:

- (Subsequential) stability for varying f
- Noise level $\delta \rightarrow 0$ and $\alpha \rightarrow 0$, $\delta^2/\alpha \rightarrow 0$
 \Rightarrow (Subsequential) convergence
- Source condition $K^*w \in \partial\Phi(u^\dagger)$, $\alpha \sim \delta$
 \Rightarrow $\mathcal{O}(\delta)$ for the Bregman distance w.r.t. Φ

General regularization properties

$$\min_{u \in X} S(Ku, f) + \alpha \Phi(u)$$

- Let: Y Hilbert space, $S(v, f) = \frac{1}{2} \|v - f\|_Y^2$

Then:

- (Subsequential) stability for varying f
- Noise level $\delta \rightarrow 0$ and $\alpha \rightarrow 0$, $\delta^2/\alpha \rightarrow 0$
 \Rightarrow (Subsequential) convergence
- Source condition $K^*w \in \partial\Phi(u^\dagger)$, $\alpha \sim \delta$
 \Rightarrow $\mathcal{O}(\delta)$ for the Bregman distance w.r.t. Φ

Difficulties:

- $\dim \ker(\Phi) = \infty \Rightarrow K$ has to be stably invertible on $\ker(\Phi)$
- Φ not coercive $\Rightarrow K$ has to be stably invertible on X

Infimal convolution regularization

Lemma:

- $\Phi_1, \Phi_2 : X \rightarrow [0, \infty]$ proper, convex, l.s.c.
- Φ_i 1-homogeneous, $\dim \ker(\Phi) < \infty$ for $i = 1, 2$
- Then: $\Phi_1 \square \Phi_2 : X \rightarrow [0, \infty]$ is exact, proper, convex, l.s.c., 1-homogeneous, $\ker \Phi_1 \square \Phi_2 = \ker \Phi_1 + \ker \Phi_2$

Infimal convolution regularization

Lemma:

- $\Phi_1, \Phi_2 : X \rightarrow [0, \infty]$ proper, convex, l.s.c.
- Φ_i 1-homogeneous, $\dim \ker(\Phi) < \infty$ for $i = 1, 2$
- Then: $\Phi_1 \square \Phi_2 : X \rightarrow [0, \infty]$ is exact, proper, convex, l.s.c., 1-homogeneous, $\ker \Phi_1 \square \Phi_2 = \ker \Phi_1 + \ker \Phi_2$

Lemma:

- Additionally: $X \hookrightarrow Z$, Z Banach space and

$$\|u\|_X \leq C(\|u\|_Z + \Phi_i(u))$$

for $i = 1, 2$ and all $u \in X$

- Then: $\Phi_1 \square \Phi_2$ is coercive, i.e.,

$$\|u - Pu\|_X \leq C\Phi(u) \quad \text{for all } u \in X$$

Infimal convolution regularization

Lemma:

- $\Phi_1, \Phi_2 : X \rightarrow [0, \infty]$ proper, convex, l.s.c.
- Φ_i 1-homogeneous, $\dim \ker(\Phi) < \infty$ for $i = 1, 2$
- Then: $\Phi_1 \square \Phi_2 : X \rightarrow [0, \infty]$ is exact, proper, convex, l.s.c., 1-homogeneous, $\ker \Phi_1 \square \Phi_2 = \ker \Phi_1 + \ker \Phi_2$

Lemma:

- Additionally: $X \hookrightarrow Z$, Z Banach space and

$$\|u\|_X \leq C(\|u\|_Z + \Phi_i(u))$$

for $i = 1, 2$ and all $u \in X$

- Then: $\Phi_1 \square \Phi_2$ is coercive, i.e.,

$$\|u - Pu\|_X \leq C\Phi(u) \quad \text{for all } u \in X$$

~~~ sufficient conditions for a regularizer

# Examples

## TV-TV<sup>2</sup> infimal convolution:

- TV and TV<sup>2</sup> have finite-dimensional kernels + embedding:

$$\|u\|_{BV} \leq \|u\|_{BV^2} \leq C(\|u\|_1 + TV^2(u))$$

- Consequently: TV  $\square \beta$  TV<sup>2</sup> is a regularizer ( $\beta > 0$ )

# Examples

## TV-TV<sup>2</sup> infimal convolution:

- TV and TV<sup>2</sup> have finite-dimensional kernels + embedding:

$$\|u\|_{BV} \leq \|u\|_{BV^2} \leq C(\|u\|_1 + TV^2(u))$$

- Consequently: TV  $\square \beta$  TV<sup>2</sup> is a regularizer ( $\beta > 0$ )

## TV- $L^1 \circ \Delta$ infimal convolution:

- $\ker(TV \square \|\cdot\|_{\mathcal{M}} \circ \Delta) = \{u \in BV \mid u \text{ harmonic}\}$
- Infinite-dimensional kernel  $\Rightarrow$  not a regularizer

# Examples

## TV-TV<sup>2</sup> infimal convolution:

- TV and TV<sup>2</sup> have finite-dimensional kernels + embedding:

$$\|u\|_{BV} \leq \|u\|_{BV^2} \leq C(\|u\|_1 + TV^2(u))$$

- Consequently: TV  $\square \beta$  TV<sup>2</sup> is a regularizer ( $\beta > 0$ )

## TV- $L^1 \circ \Delta$ infimal convolution:

- $\ker(TV \square \|\cdot\|_{\mathcal{M}} \circ \Delta) = \{u \in BV \mid u \text{ harmonic}\}$
- Infinite-dimensional kernel  $\Rightarrow$  not a regularizer

## TV-G-norm/TV- $H^{-1}$ infimal convolution:

- $\|\cdot\|_{TV^*}/\|\cdot\|_{H^{-1}}$  not coercive in  $L^p$ -spaces
- Infimal convolution not coercive  $\Rightarrow$  not a regularizer

# Examples

## TV-TV<sup>2</sup> infimal convolution:

- TV and TV<sup>2</sup> have finite-dimensional kernels + embedding:

$$\|u\|_{BV} \leq \|u\|_{BV^2} \leq C(\|u\|_1 + TV^2(u))$$

- Consequently: TV  $\square \beta$  TV<sup>2</sup> is a regularizer ( $\beta > 0$ )

## TV- $L^1 \circ \Delta$ infimal convolution:

- $\ker(TV \square \|\cdot\|_{\mathcal{M}} \circ \Delta) = \{u \in BV \mid u \text{ harmonic}\}$
- Infinite-dimensional kernel  $\Rightarrow$  not a regularizer

## TV- $G$ -norm/TV- $H^{-1}$ infimal convolution:

- $\|\cdot\|_{TV^*}/\|\cdot\|_{H^{-1}}$  not coercive in  $L^p$ -spaces
- Infimal convolution not coercive  $\Rightarrow$  not a regularizer

↝ Develop regularizing cartoon/texture models  
+ complex smoothness-based regularizers

# Outline

1 Infimal convolution regularization

2 Total generalized variation

- Definition and properties
- Applications

3 Infimal convolution TGV


- Accelerated dynamic MRI

4 Infimal convolution of oscillation TGV


- Oscillation TGV and infimal convolution
- Numerical realization
- Applications

5 Summary

# Total generalized variation



noisy image



TV-regularization

## Motivation:

- TV-based first-order regularization favors certain *artifacts*

## Total generalized variation:

[B./Kunisch/Pock '10]


$$\text{TGV}_\alpha^k(u) = \sup \left\{ \int_{\Omega} u \operatorname{div}^k v \, dx \mid v \in \mathcal{C}_c^k(\Omega, \operatorname{Sym}^k(\mathbb{R}^d)), \|\operatorname{div}^l v\|_\infty \leq \alpha_l, l = 0, \dots, k-1 \right\}$$

- $\alpha = (\alpha_0, \dots, \alpha_{k-1}) > 0$  weights


## Second-order version:

$$\text{TGV}_\alpha^2(u) = \min_{w \in \text{BD}(\Omega)} \alpha_1 \|\nabla u - w\|_{\mathcal{M}} + \alpha_0 \|\mathcal{E}w\|_{\mathcal{M}}$$

# Total generalized variation



noisy image

 $\text{TGV}_\alpha^2$ -regularization

## Motivation:

- TV-based first-order regularization favors certain *artifacts*

## Total generalized variation:

[B./Kunisch/Pock '10]

$$\text{TGV}_\alpha^k(u) = \sup \left\{ \int_{\Omega} u \operatorname{div}^k v \, dx \mid v \in \mathcal{C}_c^k(\Omega, \operatorname{Sym}^k(\mathbb{R}^d)), \|\operatorname{div}^l v\|_\infty \leq \alpha_l, l = 0, \dots, k-1 \right\}$$

- $\alpha = (\alpha_0, \dots, \alpha_{k-1}) > 0$  weights

## Second-order version:

$$\text{TGV}_\alpha^2(u) = \min_{w \in \text{BD}(\Omega)} \alpha_1 \|\nabla u - w\|_{\mathcal{M}} + \alpha_0 \|\mathcal{E}w\|_{\mathcal{M}}$$

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

**Advanced properties:** [B./Valkonen '11]

- $\text{BGV}_\alpha^2(\Omega) = \text{BV}(\Omega)$  in the sense of equivalent norms
- $\text{TGV}_\alpha^2$  is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{TGV}_\alpha^2(u)$$

for  $P : L^1(\Omega) \rightarrow \Pi^1$  continuous projection

# Properties of $\text{TGV}_\alpha^2$

**Basic properties:** [B./Kunisch/Pock '10]

- $\text{TGV}_\alpha^2$  is proper, convex, lower semi-continuous
- $\text{TGV}_\alpha^2$  is translation and rotation invariant
- $\text{TGV}_\alpha^2 + \|\cdot\|_1$  gives the Banach space  $\text{BGV}_\alpha^2(\Omega)$
- $\ker(\text{TGV}_\alpha^2) = \Pi^1$  affine functions
- $\text{TGV}_\alpha^2$  measures piecewise affine only at the interfaces

**Advanced properties:** [B./Valkonen '11]

- $\text{BGV}_\alpha^2(\Omega) = \text{BV}(\Omega)$  in the sense of equivalent norms
- $\text{TGV}_\alpha^2$  is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{TGV}_\alpha^2(u)$$

for  $P : L^1(\Omega) \rightarrow \Pi^1$  continuous projection

# Existence and stability

**Theorem:**

[B./Valkonen '11]

$$\left. \begin{array}{l} \blacksquare 1 < p \leq d/(d-1) \\ \blacksquare K : L^p(\Omega) \rightarrow H \text{ linear and continuous,} \\ \quad H \text{ Hilbert space} \\ \blacksquare K \text{ injective on } \Pi^1 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Optimization problem} \\ \min_{u \in L^p(\Omega)} \frac{1}{2} \|Ku - f\|^2 \\ \quad + \text{TGV}_\alpha^2(u) \\ \text{possesses a solution} \end{array} \right.$$

# Existence and stability

**Theorem:**

[B./Valkonen '11]


$$\left. \begin{array}{l} \blacksquare 1 < p \leq d/(d-1) \\ \blacksquare K : L^p(\Omega) \rightarrow H \text{ linear and continuous,} \\ \quad H \text{ Hilbert space} \\ \blacksquare K \text{ injective on } \Pi^1 \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Optimization problem} \\ \min_{u \in L^p(\Omega)} \frac{1}{2} \|Ku - f\|^2 \\ \quad + \text{TGV}_\alpha^2(u) \\ \text{possesses a solution} \end{array} \right.$$

**Stability:**  $f^n \rightarrow f$  in  $H \Rightarrow \begin{cases} u^n \rightharpoonup u \text{ in } L^p(\Omega) \text{ (subseq.)} \\ \text{TGV}_\alpha^2(u^n) \rightarrow \text{TGV}_\alpha^2(u) \end{cases}$


# Example: Denoising a “cartoon” image

Solve:

$$\min_{u \in L^2(\Omega)} \frac{\|u - f\|^2}{2} + \text{TGV}_{\alpha}^2(u)$$




noisy image



# Example: Denoising a “cartoon” image

Solve:

$$\min_{u \in L^2(\Omega)} \frac{\|u - f\|^2}{2} + \text{TGV}_{\alpha}^2(u)$$




TV regularization

# Example: Denoising a “cartoon” image

Solve:

$$\min_{u \in L^2(\Omega)} \frac{\|u - f\|^2}{2} + \text{TGV}_{\alpha}^2(u)$$




$\text{TGV}_{\alpha}^2$  regularization

# Electron tomography

Joint work with Georg Haberfehlner and Richard Huber




## Scanning Transmission Electron Microscopy (STEM):

- Focused electron beam
  - High Angle Annular Dark Field (HAADF)
    - ~~ non Bragg-scattered electrons
    - ~~ proportional to mass-thickness
    - ~~ Reconstruction via Radon inversion
- 

# Regularized reconstruction


Reconstruction of one slice:



Projections


Sinogram

# Regularized reconstruction




Projections

Reconstruction of one slice:



Filtered back-projection

# Regularized reconstruction




Projections

Reconstruction of one slice:




TV-regularization

# Regularized reconstruction



Projections

Reconstruction of one slice:



TGV-regularization

# Outline

- 1 Infimal convolution regularization
- 2 Total generalized variation
  - Definition and properties
  - Applications
- 3 Infimal convolution TGV
  - Accelerated dynamic MRI
- 4 Infimal convolution of oscillation TGV
  - Oscillation TGV and infimal convolution
  - Numerical realization
  - Applications
- 5 Summary

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_x}(u) = \int_{[0, T] \times \Omega} \sqrt{\epsilon |\partial_{x_1} u|^2 + \epsilon |\partial_{x_2} u|^2 + |\partial_t u|^2} dt dx$$

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_x}(u) = \int_{[0, T] \times \Omega} \sqrt{\epsilon |\partial_{x_1} u|^2 + \epsilon |\partial_{x_2} u|^2 + |\partial_t u|^2} dt dx$$

Compressed

$\text{TV}_{\epsilon_x}$

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

Compressed

$\text{TV}_{\epsilon_t}$


# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$



$\text{TV}_{\epsilon_t}$



# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

**Approach:** Optimal balancing with **infimal convolution (IC)**

$$\text{ICTV}_\epsilon = \text{TV}_{\epsilon_x} \square \text{TV}_{\epsilon_t}$$

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

**Approach:** Optimal balancing with **infimal convolution (IC)**

$$\text{ICTV}_\epsilon = \text{TV}_{\epsilon_x} \square \text{TV}_{\epsilon_t}$$

Compressed

ICTV<sub>ε</sub>

# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization


$$\text{TV}_{\epsilon_t}(u) = \int_{[0,T] \times \Omega} \sqrt{| \partial_{x_1} u |^2 + | \partial_{x_2} u |^2 + \epsilon | \partial_t u |^2} \, dt \, dx$$

**Approach:** Optimal balancing with infimal convolution (IC)

$$\text{ICTV}_{\epsilon} = \text{TV}_{\epsilon_x} \square \text{TV}_{\epsilon_t}$$



ICTV<sub>€</sub>



# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

**Approach:** Optimal balancing with **infimal convolution (IC)**

$$\text{ICTV}_\epsilon = \text{TV}_{\epsilon_x} \square \text{TV}_{\epsilon_t}$$

 $u_1$  $u_2$


# Regularization for image sequences

**Motivation:** Straightforward spatio-temporal regularization

$$\text{TV}_{\epsilon_t}(u) = \int_{[0, T] \times \Omega} \sqrt{|\partial_{x_1} u|^2 + |\partial_{x_2} u|^2 + \epsilon |\partial_t u|^2} dt dx$$

**Approach:** Optimal balancing with **infimal convolution (IC)**

$$\text{ICTV}_\epsilon = \text{TV}_{\epsilon_x} \square \text{TV}_{\epsilon_t}$$



↷ Use infimal convolution of TGV

# Infimal convolution TGV

**Definition:** Anisotropic Total Generalized Variation

$$\text{TGV}_{\beta}^k(u) = \sup \left\{ \int_{\Omega} u \operatorname{div}^k v \, dx \mid v \in \mathcal{C}_c^k(\Omega, \operatorname{Sym}^k(\mathbb{R}^d)), \right.$$
$$\left. \| \operatorname{div}^l v \|_{\infty, \beta_l^*} \leq 1, l=0, \dots, k-1 \right\}$$

- $\beta = (|\cdot|_{\beta_0}, \dots, |\cdot|_{\beta_{k-1}})$  tensor norms
- $\|v_l\|_{\infty, \beta_l^*} = \sup_{x \in \Omega} |v_l(x)|_{\beta_l^*}$ ,  $|\cdot|_{\beta_l^*}$  dual norm

# Infimal convolution TGV

**Definition:** Anisotropic Total Generalized Variation

$$\text{TGV}_\beta^k(u) = \sup \left\{ \int_{\Omega} u \operatorname{div}^k v \, dx \mid v \in \mathcal{C}_c^k(\Omega, \operatorname{Sym}^k(\mathbb{R}^d)), \right. \\ \left. \| \operatorname{div}^l v \|_{\infty, \beta_l^*} \leq 1, l=0, \dots, k-1 \right\}$$

- $\beta = (|\cdot|_{\beta_0}, \dots, |\cdot|_{\beta_{k-1}})$  tensor norms
- $\|v_l\|_{\infty, \beta_l^*} = \sup_{x \in \Omega} |v_l(x)|_{\beta_l^*}$ ,  $|\cdot|_{\beta_l^*}$  dual norm

**Infimal convolution TGV:** [Holler/Kunisch '14]

$$\text{ICTGV}_\beta^k = \text{TGV}_{\beta^1}^{k_1} \square \dots \square \text{TGV}_{\beta^m}^{k_m}$$

# Infimal convolution TGV

**Definition:** Anisotropic Total Generalized Variation

$$\text{TGV}_{\beta}^k(u) = \sup \left\{ \int_{\Omega} u \operatorname{div}^k v \, dx \mid v \in \mathcal{C}_c^k(\Omega, \operatorname{Sym}^k(\mathbb{R}^d)), \right. \\ \left. \| \operatorname{div}^l v \|_{\infty, \beta_l^*} \leq 1, l=0, \dots, k-1 \right\}$$

- $\beta = (|\cdot|_{\beta_0}, \dots, |\cdot|_{\beta_{k-1}})$  tensor norms
- $\|v_l\|_{\infty, \beta_l^*} = \sup_{x \in \Omega} |v_l(x)|_{\beta_l^*}$ ,  $|\cdot|_{\beta_l^*}$  dual norm

**Infimal convolution TGV:** [Holler/Kunisch '14]

$$\text{ICTGV}_{\beta}^k = \text{TGV}_{\beta^1}^{k_1} \square \dots \square \text{TGV}_{\beta^m}^{k_m}$$

**Regularization properties:**


- Each  $\text{TGV}_{\beta^i}^{k_i}$  has finite-dimensional kernel + embedding

$$\|u\|_{BV} \leq C(\|u\|_1 + \text{TGV}_{\beta^i}^{k_i}(u))$$

↪ **ICTGV is a regularizer**

# Accelerated dynamic MRI

*Joint work with Martin Holler and Matthias Schlögl*



sum of squares

## Dynamic MRI:

- Acquire image sequences of moving objects
- Due to limited acquisition speed, only low-resolution images are obtainable

# Accelerated dynamic MRI

*Joint work with Martin Holler and Matthias Schlögl*


sum of squares

## Dynamic MRI:

- Acquire image sequences of moving objects
- Due to limited acquisition speed, only low-resolution images are obtainable

# Accelerated dynamic MRI

Joint work with Martin Holler and Matthias Schlögl



sum of squares

## Dynamic MRI:


- Acquire image sequences of moving objects
- Due to limited acquisition speed, only low-resolution images are obtainable

## Goal:

- Improve spatio-temporal resolution by undersampling reconstruction

# Accelerated dynamic MRI

*Joint work with Martin Holler and Matthias Schlögl*



sum of squares

regularized

## Dynamic MRI:

- Acquire image sequences of moving objects
- Due to limited acquisition speed, only low-resolution images are obtainable

## Goal:

- Improve spatio-temporal resolution by undersampling reconstruction

~~~ **Apply ICTGV  
regularization**

The variational model

Minimization problem:

$$\min_u \sum_{t,c} \frac{\lambda}{2} \|K_{t,c}(u_t) - d_{t,c}\|_2^2 + \text{ICTGV}_\beta^2(u)$$

- $K_{t,c}(u_t) = M_t \mathcal{F}(u_t \sigma_c)$ masked Fourier transform
(σ_c complex coil sensitivities)
- $\text{ICTGV}_\beta^2 = \text{TGV}_{\beta^1}^2 \square \text{TGV}_{\beta^2}^2$

The variational model

Minimization problem:

$$\min_u \sum_{t,c} \frac{\lambda}{2} \|K_{t,c}(u_t) - d_{t,c}\|_2^2 + \text{ICTGV}_{\beta}^2(u)$$


- $K_{t,c}(u_t) = M_t \mathcal{F}(u_t \sigma_c)$ masked Fourier transform
(σ_c complex coil sensitivities)
- $\text{ICTGV}_{\beta}^2 = \text{TGV}_{\beta^1}^2 \square \text{TGV}_{\beta^2}^2$

Primal-dual algorithm:


- Guaranteed convergence, duality-based stopping criterion
- GPU-optimized version:
 ≈ 160 seconds including coil-sensitivity estimation
(NVidia GeForce GTX770 with AGILE library)

Numerical test

Acceleration factor 8:



Reference data



Unregularized reconstruction

Numerical test

Acceleration factor 8:

Reference data

Unregularized reconstruction

Numerical test

Acceleration factor 8:

Low rank + sparse model

Difference

Numerical test

Acceleration factor 8:

ICTGV model

Difference

Numerical test

Acceleration factor 16:

ICTGV model

Difference

Numerical test


Acceleration factor 8:

Slow component


Fast component

Numerical test

Acceleration factor 8:



Slow component




Fast component

- Favorable quantitative comparison
- 2nd place at the ISMRM 2013 reconstruction challenge

Outline

- 1 Infimal convolution regularization
- 2 Total generalized variation
 - Definition and properties
 - Applications
- 3 Infimal convolution TGV
 - Accelerated dynamic MRI
- 4 Infimal convolution of oscillation TGV
 - Oscillation TGV and infimal convolution
 - Numerical realization
 - Applications
- 5 Summary

Motivation: Denoising “barbara”



noisy image



Motivation: Denoising “barbara”



TGV_α^2 regularization

Motivation: Denoising “barbara”



TGV_{α}^2 regularization

~ Capture oscillatory structures ~ $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Oscillation TGV


Joint work with Yiming Gao

Idea:

- Choose differential equation with oscillatory functions as solutions

Oscillation TGV

Joint work with Yiming Gao



Idea:

- Choose differential equation with oscillatory functions as solutions

$$\nabla^2 u + \mathbf{c}u = 0, \quad \mathbf{c} = \omega \otimes \omega, \quad \omega \in \mathbb{R}^d, \omega \neq 0$$


- Solutions: $u(x) = C_1 \cos(\omega \cdot x) + C_2 \sin(\omega \cdot x)$, $C_1, C_2 \in \mathbb{R}$

Oscillation TGV

Joint work with Yiming Gao

Idea:

- Choose differential equation with oscillatory functions as solutions



$$\nabla^2 u + \mathbf{c}u = 0, \quad \mathbf{c} = \omega \otimes \omega, \quad \omega \in \mathbb{R}^d, \omega \neq 0$$

- Solutions: $u(x) = C_1 \cos(\omega \cdot x) + C_2 \sin(\omega \cdot x)$, $C_1, C_2 \in \mathbb{R}$
- Adapt second-order TGV to the corresponding operator


$$\nabla^2 u + \mathbf{c}u = 0$$

Oscillation TGV

Joint work with Yiming Gao

Idea:

- Choose differential equation with oscillatory functions as solutions



$$\nabla^2 u + \mathbf{c}u = 0, \quad \mathbf{c} = \omega \otimes \omega, \quad \omega \in \mathbb{R}^d, \omega \neq 0$$

- Solutions: $u(x) = C_1 \cos(\omega \cdot x) + C_2 \sin(\omega \cdot x)$, $C_1, C_2 \in \mathbb{R}$
- Adapt second-order TGV to the corresponding operator

$$\nabla u - w = 0, \quad \mathcal{E}w + \mathbf{c}u = 0$$

Oscillation TGV

Joint work with Yiming Gao


Idea:

- Choose differential equation with oscillatory functions as solutions

$$\nabla^2 u + \mathbf{c}u = 0, \quad \mathbf{c} = \omega \otimes \omega, \quad \omega \in \mathbb{R}^d, \omega \neq 0$$

- Solutions: $u(x) = C_1 \cos(\omega \cdot x) + C_2 \sin(\omega \cdot x)$, $C_1, C_2 \in \mathbb{R}$
- Adapt second-order TGV to the corresponding operator

$$\nabla u - w = 0, \quad \mathcal{E}w + \mathbf{c}u = 0$$



Total generalized variation (second order):

$$\text{TGV}_{\alpha,\beta}^2(u) = \min_{w \in \text{BD}(\Omega)} \alpha \int_{\Omega} d|\nabla u - w| + \beta \int_{\Omega} d|\mathcal{E}w|$$

- $\alpha > 0, \beta > 0$

Oscillation TGV

Joint work with Yiming Gao


Idea:

- Choose differential equation with oscillatory functions as solutions

$$\nabla^2 u + \mathbf{c}u = 0, \quad \mathbf{c} = \omega \otimes \omega, \quad \omega \in \mathbb{R}^d, \omega \neq 0$$

- Solutions: $u(x) = C_1 \cos(\omega \cdot x) + C_2 \sin(\omega \cdot x)$, $C_1, C_2 \in \mathbb{R}$
- Adapt second-order TGV to the corresponding operator

$$\nabla u - w = 0, \quad \mathcal{E}w + \mathbf{c}u = 0$$



Oscillation total generalized variation:

$$\text{TGV}_{\alpha, \beta, \mathbf{c}}^{\text{osci}}(u) = \min_{w \in \text{BD}(\Omega)} \alpha \int_{\Omega} d|\nabla u - w| + \beta \int_{\Omega} d|\mathcal{E}w + \mathbf{c}u|$$

- $\alpha > 0, \beta > 0, \mathbf{c} = \omega \otimes \omega, \omega \in \mathbb{R}^d, \omega \neq 0$

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Advanced properties:

- $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega) = \text{BV}(\Omega)$ in the sense of equivalent norms
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{TGV}_{\alpha,\beta,c}^{\text{osci}}(u)$$

for $P : L^1(\Omega) \rightarrow \ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}})$ continuous projection

Properties of $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$

Basic properties:

[Gao/B. '17]

- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is proper, convex, lower semi-continuous
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is translation and rotation invariant
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}} + \|\cdot\|_1$ gives the Banach space $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega)$
- $\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ measures piecewise oscillations only at interfaces

Advanced properties:

- $\text{BGV}_{\alpha,\beta,c}^{\text{osci}}(\Omega) = \text{BV}(\Omega)$ in the sense of equivalent norms
- $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{TGV}_{\alpha,\beta,c}^{\text{osci}}(u)$$

for $P : L^1(\Omega) \rightarrow \ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}})$ continuous projection

Infimal convolution of TGV^{osci}

Next steps:

- Separate cartoon components from oscillatory components
 - Allow for multiple directions and frequencies
- $\rightsquigarrow m\text{-fold infimal convolution}$

Infimal convolution of TGV^{osci}

Next steps:

- Separate cartoon components from oscillatory components
- Allow for multiple directions and frequencies
 - ~~~ *m-fold infimal convolution*

Infimal convolution of TGV^{osci} :

$$\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}}^{\text{osci}}(u) = (\text{TGV}_{\alpha_1, \beta_1, \mathbf{c}_1}^{\text{osci}} \square \dots \square \text{TGV}_{\alpha_m, \beta_m, \mathbf{c}_m}^{\text{osci}})(u)$$

- $\alpha_i > 0, \beta_i > 0, \mathbf{c}_i = \omega_i \otimes \omega_i, \omega_i \in \mathbb{R}^d$

Infimal convolution of TGV^{osci}

Infimal convolution of TGV^{osci} :

$$\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}}^{\text{osci}}(u) = (\text{TGV}_{\alpha_1, \beta_1, \mathbf{c}_1}^{\text{osci}} \square \dots \square \text{TGV}_{\alpha_m, \beta_m, \mathbf{c}_m}^{\text{osci}})(u)$$


- $\alpha_i > 0, \beta_i > 0, \mathbf{c}_i = \omega_i \otimes \omega_i, \omega_i \in \mathbb{R}^d$

Choice of parameters:

- $\alpha_i, \beta_i \rightsquigarrow$ regularization parameters, similar to TGV^2
- $\omega_i \rightsquigarrow$ directions and frequencies, e.g., in 2D:

$$\omega_i = f \begin{pmatrix} \sin\left(\frac{i\pi}{k}\right) \\ \cos\left(\frac{i\pi}{k}\right) \end{pmatrix}, \quad k > 0$$

$f > 0$ frequency



Infimal convolution of TGV^{osci}

Infimal convolution of TGV^{osci} :

$$\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}}^{\text{osci}}(u) = (\text{TGV}_{\alpha_1, \beta_1, \mathbf{c}_1}^{\text{osci}} \square \dots \square \text{TGV}_{\alpha_m, \beta_m, \mathbf{c}_m}^{\text{osci}})(u)$$


- $\alpha_i > 0, \beta_i > 0, \mathbf{c}_i = \omega_i \otimes \omega_i, \omega_i \in \mathbb{R}^d$

Choice of parameters:

- $\alpha_i, \beta_i \rightsquigarrow$ regularization parameters, similar to TGV^2
- $\omega_i \rightsquigarrow$ directions and frequencies, e.g., in 2D:

$$\omega_i = f \begin{pmatrix} \sin\left(\frac{i\pi}{k}\right) \\ \cos\left(\frac{i\pi}{k}\right) \end{pmatrix}, \quad k > 0$$

$f > 0$ frequency



Infimal convolution of TGV^{osci}

Infimal convolution of TGV^{osci} :

$$\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}}^{\text{osci}}(u) = (\text{TGV}_{\alpha_1, \beta_1, \mathbf{c}_1}^{\text{osci}} \square \dots \square \text{TGV}_{\alpha_m, \beta_m, \mathbf{c}_m}^{\text{osci}})(u)$$


- $\alpha_i > 0, \beta_i > 0, \mathbf{c}_i = \omega_i \otimes \omega_i, \omega_i \in \mathbb{R}^d$

Choice of parameters:

- $\alpha_i, \beta_i \rightsquigarrow$ regularization parameters, similar to TGV^2
- $\omega_i \rightsquigarrow$ directions and frequencies, e.g., in 2D:

$$\omega_i = f \begin{pmatrix} \sin\left(\frac{i\pi}{k}\right) \\ \cos\left(\frac{i\pi}{k}\right) \end{pmatrix}, \quad k > 0$$

$f > 0$ frequency



Remark:

- For $\omega = 0 \Rightarrow \text{TGV}^{\text{osci}} = \text{TGV}^2 \rightsquigarrow$ cartoon component

Infimal convolution of TGV^{osci}

Typical choice for cartoon/texture models:

- $\omega_1 = 0 \rightsquigarrow$ cartoon component
- $\omega_2, \dots, \omega_9 \rightsquigarrow$ 8 texture directions, fixed frequency

Optionally:

- $\omega_{10}, \dots, \omega_{17} \rightsquigarrow$ 8 texture directions, higher frequency

Infimal convolution of TGV^{osci}

Typical choice for cartoon/texture models:

- $\omega_1 = 0 \rightsquigarrow$ cartoon component
- $\omega_2, \dots, \omega_9 \rightsquigarrow$ 8 texture directions, fixed frequency

Optionally:

- $\omega_{10}, \dots, \omega_{17} \rightsquigarrow$ 8 texture directions, higher frequency

Variant of $\text{ICTGV}^{\text{osci}}$:

$$\begin{aligned}\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}(u) = & ((\gamma_1 \| \cdot \|_1 + \text{TGV}_{\alpha_1, \beta_1, c_1}^{\text{osci}}) \square \\ & \dots \square (\gamma_m \| \cdot \|_1 + \text{TGV}_{\alpha_m, \beta_m, c_m}^{\text{osci}}))(u)\end{aligned}$$

- $\gamma_i \geq 0$ sparsifying parameter \rightsquigarrow sparser textures

Properties of $\text{ICTGV}^{\text{osci}}$

Basic properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is proper and convex
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is translation and rotation invariant
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ gives a Banach space
- $\ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}) = \bigoplus_{\gamma_i=0} \ker(\text{TGV}_{\alpha_i, \beta_i, \mathbf{c}_i}^{\text{osci}})$

Properties of $\text{ICTGV}^{\text{osci}}$

Basic properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}$ is proper and convex
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}$ is translation and rotation invariant
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ gives a Banach space
- $\ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}) = \bigoplus_{\gamma_i=0} \ker(\text{TGV}_{\alpha_i, \beta_i, \mathbf{c}_i}^{\text{osci}})$

Properties of $\text{ICTGV}^{\text{osci}}$

Basic properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}$ is proper and convex
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}$ is translation and rotation invariant
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ gives a Banach space
- $\ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\mathbf{c}}, \vec{\gamma}}^{\text{osci}}) = \bigoplus_{\gamma_i=0} \ker(\text{TGV}_{\alpha_i, \beta_i, \mathbf{c}_i}^{\text{osci}})$

Properties of $\text{ICTGV}^{\text{osci}}$

Basic properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}$ is proper and convex
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}$ is translation and rotation invariant
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ gives a Banach space
- $\ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}) = \bigoplus_{\gamma_i=0} \ker(\text{TGV}_{\alpha_i, \beta_i, \epsilon_i}^{\text{osci}})$

Properties of ICTGV^{osci}

Advanced properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is lower semi-continuous and exact, i.e.,
the infimum in the definition is attained
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ is an equivalent norm on $\text{BV}(\Omega)$
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}(u)$$

for $P : L^1(\Omega) \rightarrow \ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}})$ continuous projection

Properties of ICTGV^{osci}

Advanced properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is lower semi-continuous and exact, i.e., the infimum in the definition is attained
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ is an equivalent norm on $\text{BV}(\Omega)$
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}(u)$$

for $P : L^1(\Omega) \rightarrow \ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}})$ continuous projection

Properties of ICTGV^{osci}

Advanced properties:

- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is lower semi-continuous and exact, i.e., the infimum in the definition is attained
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}} + \|\cdot\|_1$ is an equivalent norm on $\text{BV}(\Omega)$
- $\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}$ is coercive in the sense

$$\|u - Pu\|_{\text{BV}} \leq C \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}}(u)$$

for $P : L^1(\Omega) \rightarrow \ker(\text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{c}, \vec{\gamma}}^{\text{osci}})$ continuous projection

Solution of imaging problems

Theorem:

$$\left. \begin{array}{l} \blacksquare 1 < p \leq d/(d-1) \\ \blacksquare K : L^p(\Omega) \rightarrow Y \text{ linear and continuous,} \\ \quad Y \text{ Banach space} \\ \blacksquare f \in Y, 1 \leq q < \infty \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Optimization problem} \\ \min_{u \in L^p(\Omega)} \frac{1}{q} \|Ku - f\|_Y^q \\ \quad + \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}(u) \\ \text{possesses a solution} \end{array} \right.$$

Solution of imaging problems

Theorem:

$$\left. \begin{array}{l} \blacksquare 1 < p \leq d/(d-1) \\ \blacksquare K : L^p(\Omega) \rightarrow Y \text{ linear and continuous,} \\ \quad Y \text{ Banach space} \\ \blacksquare f \in Y, 1 \leq q < \infty \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Optimization problem} \\ \min_{u \in L^p(\Omega)} \frac{1}{q} \|Ku - f\|_Y^q \\ \quad + \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}(u) \\ \text{possesses a solution} \end{array} \right.$$

Remarks:

- Ensures applicability to a variety of imaging problems (denoising, deblurring, image reconstruction (CT, MRI), ...)
- Intrinsic cartoon/texture decomposition for these problems

Solution of imaging problems

Theorem:

$$\left. \begin{array}{l} \blacksquare 1 < p \leq d/(d-1) \\ \blacksquare K : L^p(\Omega) \rightarrow Y \text{ linear and continuous,} \\ \quad Y \text{ Banach space} \\ \blacksquare f \in Y, 1 \leq q < \infty \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Optimization problem} \\ \min_{u \in L^p(\Omega)} \frac{1}{q} \|Ku - f\|_Y^q \\ \quad + \text{ICTGV}_{\vec{\alpha}, \vec{\beta}, \vec{\epsilon}, \vec{\gamma}}^{\text{osci}}(u) \\ \text{possesses a solution} \end{array} \right.$$

Remarks:

- Ensures applicability to a variety of imaging problems (denoising, deblurring, image reconstruction (CT, MRI), ...)
- Intrinsic cartoon/texture decomposition for these problems

Numerical realization

Discretization in 2D and for $Y = L^2(\Omega')$:

- Discretize $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ with forward/backward differences

- Maintain kernel, i.e.,

$$\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$$

$$\rightsquigarrow \mathbf{c} = \begin{bmatrix} 2 - 2\cos(\omega_1) & 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) \\ 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) & 2 - 2\cos(\omega_2) \end{bmatrix}$$

$$\text{for } \omega = (\omega_1, \omega_2) \in \mathbb{R}^2 \setminus \pi\mathbb{Z}^2$$

- Solve non-smooth discrete optimization problems with first-order methods, such as a primal-dual iteration

[Chambolle/Pock '11]

Numerical realization

Discretization in 2D and for $Y = L^2(\Omega')$:

- Discretize $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ with forward/backward differences

- Maintain kernel, i.e.,

$$\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$$

$$\rightsquigarrow \mathbf{c} = \begin{bmatrix} 2 - 2\cos(\omega_1) & 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) \\ 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) & 2 - 2\cos(\omega_2) \end{bmatrix}$$

$$\text{for } \omega = (\omega_1, \omega_2) \in \mathbb{R}^2 \setminus \pi\mathbb{Z}^2$$

- Solve non-smooth discrete optimization problems with first-order methods, such as a primal-dual iteration

[Chambolle/Pock '11]

Numerical realization

Discretization in 2D and for $Y = L^2(\Omega')$:

- Discretize $\text{TGV}_{\alpha,\beta,c}^{\text{osci}}$ with forward/backward differences

- Maintain kernel, i.e.,

$$\ker(\text{TGV}_{\alpha,\beta,c}^{\text{osci}}) = \text{span}\{x \mapsto \cos(\omega \cdot x), x \mapsto \sin(\omega \cdot x)\}$$

$$\rightsquigarrow \mathbf{C} = \begin{bmatrix} 2 - 2\cos(\omega_1) & 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) \\ 1 + \cos(\omega_1 - \omega_2) - \cos(\omega_1) - \cos(\omega_2) & 2 - 2\cos(\omega_2) \end{bmatrix}$$

$$\text{for } \omega = (\omega_1, \omega_2) \in \mathbb{R}^2 \setminus \pi\mathbb{Z}^2$$

- Solve non-smooth discrete optimization problems with first-order methods, such as a primal-dual iteration


[Chambolle/Pock '11]

Optimization algorithm

Iteration:

$$\left\{ \begin{array}{l} \lambda^{n+1} = (\lambda^n + \sigma(K \sum_{i=1}^m \bar{u}_i^n - f)) / (1 + \sigma) \\ \text{for } i = 1, \dots, m \text{ do} \\ \quad p_i^{n+1} = \mathcal{P}_{\alpha_i}(p_i^n + \sigma(\nabla \bar{u}_i^n - \bar{w}_i^n)) \\ \quad q_i^{n+1} = \mathcal{P}_{\beta_i}(q_i^n + \sigma(\mathcal{E} \bar{w}_i^n + \mathbf{c}_i \bar{u}_i^n)) \\ \quad \tilde{u}_i^{n+1} = u_i^n - \tau(K^* \lambda^{n+1} - \operatorname{div}_1 p_i^{n+1} + \mathbf{c}_i q_i^{n+1}) \\ \quad u_i^{n+1} = \operatorname{Shrink}_{\tau \gamma_i}(\tilde{u}_i^{n+1}) \\ \quad w_i^{n+1} = w_i^n + \tau(p_i^{n+1} + \operatorname{div}_2 q_i^{n+1}) \\ \quad \bar{u}_i^{n+1} = 2u_i^{n+1} - u_i^n \\ \quad \bar{w}_i^{n+1} = 2w_i^{n+1} - w_i^n \\ \text{end for} \end{array} \right.$$

Example: Denoising “barbara”



noisy image



Example: Denoising “barbara”



TGV² regularization

Example: Denoising “barbara”



$\text{ICTGV}^{\text{osci}}$ regularization


Example: Denoising “barbara”



$\text{ICTGV}^{\text{osci}}$ regularization

~ Better recovery of oscillatory structures


Cartoon/texture decomposition



Tested algorithms:


- Total variation/ G -norm decomposition [Aujol et al. '05]
- Total variation/ H^{-1} -norm decomposition [Osher/Sole/Vese '03]
- Framelet/Local discrete cosine transform model [Cai/Osher/Shen '09]
- Infimal convolution of oscillation TGV

Cartoon/texture decomposition




TV- G -norm decomposition

Cartoon/texture decomposition




TV- H^{-1} -norm decomposition

Cartoon/texture decomposition



Framelet+LDCT model

Cartoon/texture decomposition




ICTGV^{osci} model

Example: Image denoising

Tested models/algorithms:

- Total generalized variation of second order [B./Kunisch/Pock '10]
- Nonlocal total variation [Gilboa/Osher '08]
- Infimal convolution of total generalized variation [Holler/Kunisch '14]
- Framelet + local discrete cosine transform [Cai/Osher/Shen '09]
- Infimal convolution of oscillation TGV
- Block matching and 3D filtering [Dabov et al. '07]


Example: Denoising “barbara”



noisy image



Example: Denoising “barbara”




TGV² regularization (27.41 dB)

Example: Denoising “barbara”



Nonlocal TV regularization (32.05 dB)

Example: Denoising “barbara”



ICTGV regularization (31.15 dB)

Example: Denoising “barbara”




Framelet + LDCT model (31.70 dB)

Example: Denoising “barbara”



$\text{ICTGV}^{\text{osci}}$ regularization (32.21 dB)

Example: Denoising “barbara”



BM3D (34.43 dB)

Example: Image inpainting

Setup:


- Recover an image from 50% randomly deleted pixels



Tested algorithms:

- Total generalized variation
[B./Kunisch/Pock '10]
- Framelet/Local discrete cosine transform model
[Cai/Osher/Shen '09]
- Infimal convolution of oscillation TGV

Example: Inpainting “barbara”



corrupted image

Example: Inpainting “barbara”




TGV² regularization (27.49 dB)

Example: Inpainting “barbara”




Framelet + LDCT model (32.75 dB)

Example: Inpainting “barbara”



$\text{ICTGV}^{\text{osci}}$ regularization (34.03 dB)

Example: Undersampled MRI



Setup:

- Recover an image from a few radially sampled Fourier coefficients

Tested algorithms:


- Total generalized variation + shearlet model [Guo/Qin/Yin '14]
- Infimal convolution of oscillation TGV

Example: MRI reconstruction



TGV + shearlet model (70 radial lines)

Example: MRI reconstruction



TGV + shearlet model (60 radial lines)

Example: MRI reconstruction




TGV + shearlet model (50 radial lines)

Example: MRI reconstruction



$\text{ICTGV}^{\text{osci}}$ model (70 radial lines)

Example: MRI reconstruction



$\text{ICTGV}^{\text{osci}}$ model (60 radial lines)

Example: MRI reconstruction



$\text{ICTGV}^{\text{osci}}$ model (50 radial lines)

Example: MRI reconstruction



ground truth



Example: MRI reconstruction

Quantitative comparison:

| Model | 50 lines | 60 lines | 70 lines |
|-----------------------|----------|----------|----------|
| TGV + shearlet | 28.77 dB | 29.59 dB | 30.10 dB |
| ICTGV ^{osci} | 29.16 dB | 29.98 dB | 30.56 dB |

Outline

- 1 Infimal convolution regularization
- 2 Total generalized variation
 - Definition and properties
 - Applications
- 3 Infimal convolution TGV
 - Accelerated dynamic MRI
- 4 Infimal convolution of oscillation TGV
 - Oscillation TGV and infimal convolution
 - Numerical realization
 - Applications
- 5 Summary

Summary

- Infimal convolution is a flexible tool to combine variational models, covering several known approaches
- It provides a regularizer for linear inverse problems in many cases
- It allows to construct stabilizing and regularizing cartoon/texture decomposition models
- Numerical experiments show a promising efficiency



Yiming Gao and Kristian Bredies.

Infimal convolution of oscillation total generalized variation for the recovery of images with structured texture.

SIAM Journal on Imaging Sciences, 11(3):2021–2063, 2018.



Supported by the Austrian Science Fund (FWF)
Project P29192 “Regularization graphs for variational imaging”