FILTERMATE 5 FOR QGIS

Five business questions. One panel.

Vector, raster and LiDAR point cloud layers filtered by one another in QGIS, with no SQL and no script. Five scenarios recorded on Estavar (French Pyrenees): BD TOPO, cadastre, 1 m DTM, 25 cm CIR orthophoto, LiDAR HD.

1 **Buildings along the main roads**
vector → vector

2 **The survey package around the church**
vector → raster + point cloud

3 **Parcels on a slope**
raster → vector

4 **Vegetated parcels**
raster → vector

5 **Roads to prune**
point cloud → vector


EXPLORING selects. FILTERING relates. EXPORTING writes.

imagodata.github.io/filter_mate · official QGIS plugin repository

Free and open source · QGIS 3.22 to 4.2

Buildings along the main roads

vector → vector



WHAT YOU DO

Source: the road segments with `"importance" = '4'`, 15 segments. Intersect predicate, 50 m buffer. Targets: buildings and parcels.


WHAT YOU GET

39 buildings and 11 parcels, filtered in place, ready to export as GeoPackage.

EXPLORING selects. FILTERING relates. EXPORTING writes.

The survey package around the church

vector → raster + point cloud



WHAT YOU DO

Source: the church (single selection), 50 m buffer. Targets: the 1 m DTM and the LiDAR HD tile.


WHAT YOU GET

A clipped DTM (VRT) and a derived cloud (COPC) that keep their style; Undo brings the source back.

EXPLORING selects. FILTERING relates. EXPORTING writes.

Parcels on a slope

raster → vector



WHAT YOU DO

Source: the DTM, "slope (%)" computed on the fly, threshold ≥ 20 %. Target: parcels, minimum coverage 50 %.


WHAT YOU GET

2 parcels out of 42 have more than half of their area on the slope.

EXPLORING selects. FILTERING relates. EXPORTING writes.

Vegetated parcels

raster → vector



WHAT YOU DO

Source: the CIR orthophoto, band roles NIR-R-G, NDVI vegetation preset (> 0.3). Target: parcels, minimum coverage 50 %.

WHAT YOU GET


7 mostly vegetated parcels; the selection is read at full resolution, tile by tile.

EXPLORING selects. FILTERING relates. EXPORTING writes.

FILTERMATE 5 FOR QGIS

Roads to prune

point cloud → vector



WHAT YOU DO

Source: the LiDAR HD, class 5 (high vegetation). Target: road segments, 3 m buffer around the footprint of the points.

WHAT YOU GET

100 segments out of 131 touch high vegetation within 3 m.

EXPLORING selects. FILTERING relates. EXPORTING writes.

FILTERMATE 5 FOR QGIS

New this week

Three features merged on the main branch, on their way to 5.0.1.

Clicked-object pipette

the patch under the click (pool, roof, field), not every similar pixel of the raster

Similar features

one click on a feature selects every feature with the same value

MCP server

the filtering driven from an AI assistant, one tool per REST endpoint

EXPLORING selects. FILTERING relates. EXPORTING writes.

imagodata.github.io/filter_mate · official QGIS plugin repository

Free and open source · QGIS 3.22 to 4.2