

Fit of catch data


Total catch




Total catch by fleet




Catch residuals by fleet




Absolute residual catch by fleet



Residuals catch-at-age by fleet



Residuals catch-at-length by fleet



Age fits N_Chile


Observed



Predicted



Proportion at age



Age

Age fits SC_Chile_PS


Observed



Predicted



Proportion at age



Age

Age fits Offshore_Trawl


Observed



Predicted



Proportion at age



Length fits FarNorth

Observed




Predicted



10 20 30 40 50

10 20 30 40 50

Proportion at length



Predicted and observed catches by fleet

Observed



Predicted



1970

1980

1990

2000

2010

Offshore_Trawl

SC_Chile_PS

4000

3000

2000

1000

0

FarNorth

N_Chile

4000

3000

2000

1000

0

1970

1980


1990

2000

2010

Years

Thousand tonnes




Fit of survey data

Predicted and observed indices

Observed



Predicted



Standardized survey residuals


-4.99
-3.08
-1.17




0.74
2.64
4.55



Log residuals




SD per input series




Fleet & Survey sel & F


Selectivity of the survey by Pentad




F at age




F proportion at age



N at age



Proportion at age



Fishery mean age

Observed



Modelled



SC_Chile_PS

10

8

6

4

2

N_Chile

Offshore_Trawl

10

8


6

4

2

Years

Age



Fishery mean length

Observed



Modelled



FarNorth

Length (cm)

Years

35

30


25

1980

1990

2000

2010



Survey mean age


Observed




Modelled




Age



Chile_AcousCS




Chile_AcousN




Years

Stock summary

Summary sheet



Uncertainty of key parameters



Mature – Immature fish

Mature

Immature —

Biomass in kt

20000

15000

10000

5000

0

1970


1980

1990

2000


2010

Years



Stock Recruitment

Observed ● Modelled - - -



Fished vs. unfished biomass

Fished



Unfished



Total biomass

40000
30000
20000
10000

1970

1980

1990

2000

2010

Years

