

MAT454 Notes

Jad Elkhaleq Ghalayini

January 17 2020

Zeros and Poles

Definition 1 (Zero). If f is holomorphic in a neighborhood of $z_0 \in \mathbb{C}$ and $f(z_0) = 0$, we can write, for some $k \in \mathbb{N}$,

$$f(z) = (z - z_0)^k f_1(z)$$

where $f_1(z)$ is nonvanishing near z_0 . In this case k is called the **order** or **multiplicity** of the **zero** z_0

Zeros of holomorphic functions form a discrete set. We want to study, however, not only holomorphic functions, but also quotients of holomorphic functions

Definition 2 (Meromorphic). A function f is **meromorphic** on an open $\Omega \subseteq \mathbb{C}$ if it is defined and holomorphic in the complement of a discrete set such that in some neighborhood of every point of Ω we can write $f(z) = g(z)/h(z)$ where g, h are holomorphic and h is not identically zero.

Why is it interesting to work with meromorphic and not just holomorphic functions? Essentially, it's because meromorphic functions in a domain Ω form a field (whereas holomorphic functions only form a ring). Note that, in this course, when we say "domain", what we mean is a connected open set. If $f(z), g(z)$ are holomorphic near z_0 , like before, we can write

$$f(z) = (z - z_0)^k f_1(z), \quad g(z) = (z - z_0)^\ell g_1(z)$$

where $f_1(z_0), g_1(z_0) \neq 0$. Near z_0 , then, the quotient looks like

$$\frac{f(z)}{g(z)} = (z - z_0)^{k-\ell} \frac{f_1(z)}{g_1(z)}$$

So what are the different possibilities? If $k \geq \ell$, then this function extends to be holomorphic at z_0 . On the other hand, if $k < \ell$, then, of course,

$$\lim_{z \rightarrow z_0} \left| \frac{f(z)}{g(z)} \right| = \infty$$

Note: *not* undefined, but ∞ . In this case, we say that z_0 is a pole of order $\ell - k$. Holomorphic functions in an annulus $r < |z| < R$ have a convergent Laurent expansion in an annulus

$$\sum_{n=-\infty}^{\infty} a_n z^n = \sum_{n<0} a_n z^n + \sum_{n \geq 0} a_n z^n$$

Note that the LHS converges when $r < |z|$, whereas the RHS converges when $|z| < R$. This actually comes from Cauchy's theorem, just in the case of a convergent power series expansion of a holomorphic function. So this is from Cauchy's integral formula:

$$f(z) = \frac{1}{2\pi i} \int_{\gamma_1} \frac{f(\xi)}{\xi - z} d\xi - \frac{1}{2\pi i} \int_{\gamma_2} \frac{f(\xi)}{\xi - z} d\xi = \sum_{n=-\infty}^{\infty} a_n z^n$$

for z between γ_1, γ_2 . So

$$a_n = \frac{1}{2\pi i} \int_{\gamma_\gamma} \frac{f(\xi)}{\xi^{n+1}} d\xi$$

So of course, this should be just like before for the positive part. Note that this is the integral over γ_1 if $n \geq 0$ and over γ_2 if $n < 0$.