

MAT454 Notes

Jad Elkhaleq Ghalayini

February 5 2020

Definition 1 (n -dimensional complex projective space). *We define*

$$\mathbb{P}^n(\mathbb{C}) = \mathbb{C}^{n+1} \setminus \{0\} / \sim$$

where

$$(x_0, \dots, x_n) \sim (x'_0, \dots, x'_n) \iff \exists \lambda \in \mathbb{C}, (x'_0, \dots, x'_n) = (\lambda x_0, \dots, \lambda x_n)$$

We denote the equivalence class of (x_0, \dots, x_n) by $[x_0, \dots, x_n]$.

Definition 2 (Homogeneous coordinates). *We define coordinate charts $U_i = \{[x_0, \dots, x_n] \in \mathbb{P}^n(\mathbb{C}) : x_i \neq 0\}$ with affine coordinates $U_i \rightarrow \mathbb{C}^n$,*

$$[x_0, \dots, x_n] \mapsto \left(\frac{x_0}{x_i}, \dots, \frac{x_{i-1}}{x_i}, \frac{x_{i+1}}{x_i}, \dots, \frac{x_n}{x_i} \right)$$

with inverse

$$(g_1, \dots, g_n) \mapsto [g_1, \dots, g_{i-1}, 1, g_{i+1}, \dots, g_n]$$

Using these coordinates, we have that $\mathbb{P}^n(\mathbb{C})$ has the structure of an n -dimensional complex manifold, as the transition mappings are rational. Let's take one of the charts here, say U_0 , to be \mathbb{C}^n . So

$$\mathbb{P}^n(\mathbb{C}) = U_0 \cup \text{everything else}$$

But what's everything else? So U_0 is all the points where $x_0 \neq 0$, so everything else is the set of points

$$\{x_0 = 0\} = \{[0, x_1, \dots, x_n]\} \simeq \mathbb{P}^{n-1}(\mathbb{C}) \implies \mathbb{P}^n(\mathbb{C}) = U_0 \cup \mathbb{P}^{n-1}(\mathbb{C})$$

We call this copy of $\mathbb{P}^{n-1}(\mathbb{C}) \simeq \{x_0 = 0\}$ the **hyperplane at infinity**. This is like a generation of the Riemann sphere which we saw before, which we saw was given by $S^2 = \mathbb{P}^1(\mathbb{C})$. So when we talk about $\mathbb{P}^2(\mathbb{C})$, that's like having 2-complex coordinates with a line at infinity. Specifically, we can write it as

$$\mathbb{P}^2(\mathbb{C}) = \{[x, y, t]\} = \mathbb{C}_{(x,y)}^2 \cup \{t = 0\}$$

the **projective line at infinity**. Now assume we have a curve $X \subset \mathbb{C}^2$ generated by the equation

$$y^2 = 4x^3 - 20a_2x - 28a_4$$

where the RHS has three distinct roots. We want to compute the **compactification of X in $\mathbb{P}^2(\mathbb{C})$** .