

# MAT454 Notes

Jad Elkhaleq Ghalayini

January 17 2020

## Zeros and Poles

**Definition 1** (Zero). *If  $f$  is holomorphic in a neighborhood of  $z_0 \in \mathbb{C}$  and  $f(z_0) = 0$ , we can write, for some  $k \in \mathbb{N}$ ,*

$$f(z) = (z - z_0)^k f_1(z)$$

*where  $f_1(z)$  is nonvanishing near  $z_0$ . In this case  $k$  is called the **order** or **multiplicity** of the **zero**  $z_0$*

Zeros of holomorphic functions form a discrete set. We want to study, however, not only holomorphic functions, but also quotients of holomorphic functions

**Definition 2** (Meromorphic). *A function  $f$  is **meromorphic** on an open  $\Omega \subseteq \mathbb{C}$  if it is defined and holomorphic in the complement of a discrete set such that in some neighborhood of every point of  $\Omega$  we can write  $f(z) = g(z)/h(z)$  where  $g, h$  are holomorphic and  $h$  is not identically zero.*

Why is it interesting to work with meromorphic and not just holomorphic functions? Essentially, it's because meromorphic functions in a domain  $\Omega$  form a field (whereas holomorphic functions only form a ring). Note that, in this course, when we say “domain”, what we mean is a connected open set. If  $f(z), g(z)$  are holomorphic near  $z_0$ , like before, we can write

$$f(z) = (z - z_0)^k f_1(z), \quad g(z) = (z - z_0)^\ell g_1(z)$$

where  $f_1(z_0), g_1(z_0) \neq 0$ . Near  $z_0$ , then, the quotient looks like

$$\frac{f(z)}{g(z)} = (z - z_0)^{k-\ell} \frac{f_1(z)}{g_1(z)}$$

So what are the different possibilities? If  $k \geq \ell$ , then this function extends to be holomorphic at  $z_0$ . On the other hand, if  $k < \ell$ , then, of course,

$$\lim_{z \rightarrow z_0} \left| \frac{f(z)}{g(z)} \right| = \infty$$

Note: *not* undefined, but  $\infty$ . In this case, we say that  $z_0$  is a pole of order  $\ell - k$ .