



Section 1: Hospital Pricing and Selection on Observables

Ian McCarthy | Emory University
Econ 470 & HLTH 470


Table of contents

1. Hospital Pricing
2. HCRIS Data
3. Causal Inference and Potential Outcomes
4. Average Treatment Effects
5. Selection Bias
6. Matching and Weighting
7. Pricing and Profit Status

Background on Hospital Pricing

What is a hospital price?

Defining characteristic of hospital services: *it's complicated!*



1/04/11	1	0416826	CLEARLINK DUO-VENT	17.00
1/04/11	1	0406462	TUBE CONNECTING STERIL 6FT	27.00
1/04/11	1	0406462	TUBE CONNECTING STERIL 6FT	27.00
1/05/11	1	3005741	ACCU-CHEK CCRV	18.00
1/05/11	1	3019692	SURGICEL 2X14 STRIP EACH	451.00
1/05/11	1	3005741	ACCU-CHEK CCRV	18.00
1/05/11	1	3005741	ACCU-CHEK CCRV	18.00
1/05/11	1-	3019692	SURGICEL 2X14 STRIP EACH	451.00-
1/05/11	1	3005741	ACCU-CHEK CCRV	18.00
1/05/11	1	3005741	ACCU-CHEK CCRV	18.00
1/05/11	10	2900025	OXYGEN HOURLY	560.00
1/05/11	1	0402230	LEUKINS TUBE SPECIM TRAP	77.00
1/05/11	1	0416826	SET EXTENSION 1-VALVE	12.00
1/05/11	1	0406793	SUCTION YANKAER	44.00
1/05/11	1	0416826	SECOND SET LUER LOCK	5.00
		0416826		12.00

Brill, Steven. 2013. "Bitter Pill: Why Medical Bills are Killing Us." *Time Magazine*.


What is a hospital price?

Lots of different payers paying lots of different prices:

- Medicare fee-for-service prices
- Medicaid payments
- Private insurance negotiations (including Medicare Advantage)
- But what about the price to patients?

Price \neq charge \neq cost \neq patient out-of-pocket spending

What is a hospital price?



Source: [Health Care Pricing Project](#)

What is a hospital price?

Not clear what exactly is negotiated...

Fee-for-service

- price per procedure
- percentage of charges
- markup over Medicare rates

Capitation

- payment per patient
- pay-for-performance
- shared savings


Hospital prices in real life

A few empirical facts:

1. Hospital services are expensive
2. Prices vary dramatically across different areas
3. Lack of competition is a major reason for high prices

Hospital prices in real life


Hospital Prices for Hip Replacement
Atlanta, GA HRR, 2008-2011



Note: Each column captures a hospital's negotiated transaction price and Medicare reimbursement. Prices are averaged from 2008-2011 and presented in 2011 dollars. CoV captures the coefficient of variation of hospital negotiated transaction prices within the HRR. Max/Min captures the max/min ratio of hospital's negotiated transaction prices within the HRR. Horizontal lines indicate average rates and prices within the region.

© Health Care Pricing Project

Hospital Prices for Lower Limb MRI
Atlanta, GA HRR, 2008-2011



Note: Each column captures a hospital's negotiated transaction price and Medicare reimbursement. Prices are averaged from 2008-2011 and presented in 2011 dollars. CoV captures the coefficient of variation of hospital negotiated transaction prices within the HRR. Max/Min captures the max/min ratio of hospital's negotiated transaction prices within the HRR. Horizontal lines indicate average rates and prices within the region.

© Health Care Pricing Project

Understanding HCRIS Data

What is HCRIS?

Healthcare Cost Report Information System ('cost reports')

- Nursing Homes (SNFs)
- Hospice
- Home Health Agencies
- Hospitals

Hospital Cost Reports

10-12

FORM CMS-2552-10

4090 (Cont.)

STATEMENT OF PATIENT REVENUES
AND OPERATING EXPENSES

PROVIDER CCN:

PERIOD:

FROM _____
TO _____

WORKSHEET G-2,
PARTS I & II

PART I - PATIENT REVENUES

REVENUE CENTER		INPATIENT	OUTPATIENT	TOTAL	
		1	2	3	
GENERAL INPATIENT ROUTINE CARE SERVICES					
1	Hospital				1
2	Subprovider IPF				2
3	Subprovider IRF				3
4	Subprovider (Other)				4
5	Swing bed - SNF				5
6	Swing bed - NF				6
7	Skilled nursing facility				7
8	Nursing facility				8
9	Other long term care				9
10	Total general inpatient care services (sum of lines 1-9)				10
INTENSIVE CARE TYPE INPATIENT HOSPITAL SERVICES					
11	Intensive care unit				11
12	Coronary care unit				12
13	Burn intensive care unit				13
14	Surgical intensive care unit				14
15	Other special care (specify)				15
16	Total intensive care type inpatient hospital services (sum of lines 11-15)				16
17	Total inpatient routine care services (sum of lines 10 and 16)				17
18	Ancillary services				18
19	Outpatient services				19
20	Rural Health Clinic (RHC)				20
21	Federally Qualified Health Center (FQHC)				21
22	Home health agency				22


The Data

Let's work with the [HCRIS GitHub repository](#). But forming the dataset is up to you this time.




The Data

```
hcris.data %>%  
  ggplot(aes(x=as.factor(year))) +  
  geom_bar() +  
  labs(  
    x="Year",  
    y="Number of Hospitals",  
    title="Number of Hospitals per Year"  
  ) + theme_bw() +  
  theme(axis.text.x = element_text(angle = 90, hjust=1))
```



Number of hospitals




Estimating hospital prices

```
hcris.data <- hcris.data %>%  
  mutate( discount_factor = 1-tot_discounts/tot_charges,  
          price_num = (ip_charges + icu_charges + ancillary_charges)*discount_factor - tot_mcare_payment,  
          price_denom = tot_discharges - mcare_discharges,  
          price = price_num/price_denom)
```



Estimating hospital prices

```
hcris.data %>% group_by(year) %>%  
  filter(price_denom>10, !is.na(price_denom),  
         price_num>0, !is.na(price_num)) %>%  
  select(price, year) %>%  
  summarize(mean_price=mean(price, na.rm=TRUE))  
ggplot(aes(x=as.factor(year), y=mean_price))  
  geom_line(aes(group=1)) +  
  labs(  
    x="Year",  
    y="Average Hospital Price",  
    title="Hospital Prices per Year"  
  ) + scale_y_continuous(labels=comma) +  
  theme_bw() + theme(axis.text.x = element_te
```



Estimating hospital prices

```
hcris.data %>% group_by(year) %>%  
  filter(price_denom>100, !is.na(price_denom),  
         price_num>0, !is.na(price_num),  
         price<1000000) %>%  
  select(price, year) %>%  
  summarize(mean_price=mean(price, na.rm=TRUE))  
ggplot(aes(x=as.factor(year), y=mean_price))  
  geom_line(aes(group=1)) +  
  labs(  
    x="Year",  
    y="Average Hospital Price",  
    title="Hospital Prices per Year"  
  ) + scale_y_continuous(labels=comma) +  
  theme_bw() + theme(axis.text.x = element_te
```



Causal Inference and Potential Outcomes

Why causal inference?

Another example: **What price should we charge for a night in a hotel?**

Machine Learning

- Focuses on prediction
- High prices are strongly correlated with higher sales
- Increase prices to attract more people?

Causal Inference

- Focuses on **counterfactuals**
- What would sales look like if prices were higher?

Goal of Causal Inference

- **Goal:** Estimate effect of some policy or program
- Key building block for causal inference is the idea of **potential outcomes**

Some notation

Observed outcome

$$Y_i = Y_{1i} \times D_i + Y_{0i} \times (1 - D_i)$$

or

$$Y_i = \begin{cases} Y_{1i} & \text{if } D_i = 1 \\ Y_{0i} & \text{if } D_i = 0 \end{cases}$$

Assumes **SUTVA** (stable unit treatment value assumption)...no interference across units

Example of "Potential Outcomes"



$$Y_1 = \$75,000$$

$$\text{Earnings due to Emory} = Y_1 - Y_0 = \$15,000$$



$$Y_0 = \$60,000$$

Example of "Potential Outcomes"



$$Y_1 = \$75,000$$

$$\text{Earnings due to Emory} = Y_1 - Y_0 = ?$$



$$Y_0 = ?$$

Do we ever observe the potential outcomes?



Without a time machine...not possible to get *individual* effects.

Fundamental Problem of Causal Inference

- We don't observe the counterfactual outcome...what would have happened if a treated unit was actually untreated.
- *ALL* attempts at causal inference represent some attempt at estimating the counterfactual outcome. We need an estimate for Y_0 among those that were treated, and vice versa for Y_1 .

Average Treatment Effects

Different treatment effects

Tend to focus on **averages**¹:

- **ATE**: $\delta_{ATE} = E[Y_1 - Y_0]$
- **ATT**: $\delta_{ATT} = E[Y_1 - Y_0 | D = 1]$
- **ATU**: $\delta_{ATU} = E[Y_1 - Y_0 | D = 0]$

¹ or similar measures such as medians or quantiles

Average Treatment Effects

- **Estimand:**

$$\delta_{ATE} = E[Y_1 - Y_0] = E[Y|D = 1] - E[Y|D = 0]$$

- **Estimate:**

$$\hat{\delta}_{ATE} = \frac{1}{N_1} \sum_{D_i=1} Y_i - \frac{1}{N_0} \sum_{D_i=0} Y_i,$$

where N_1 is number of treated and N_0 is number untreated (control)

- With random assignment and equal groups, inference/hypothesis testing with standard two-sample t-test

Selection Bias

Selection bias

- Without random assignment, there's a high probability that

$$E[Y_{0i}|D_i = 1] \neq E[Y_{0i}|D_i = 0]$$

- i.e., outcomes without treatment are different for the treated group

Omitted variables bias

- In a regression setting, selection bias is the same problem as omitted variables bias (OVB)
- Quick review: Goal of OLS is to find $\hat{\beta}$ to "best fit" the linear equation $y_i = \alpha + x_i\beta + \epsilon_i$

Regression review

$$\min_{\beta} \sum_{i=1}^N (y_i - \alpha - x_i \beta)^2 = \min_{\beta} \sum_{i=1}^N (y_i - (\bar{y} - \bar{x} \beta) - x_i \beta)^2$$

$$0 = \sum_{i=1}^N \left(y_i - \bar{y} - (x_i - \bar{x}) \hat{\beta} \right) (x_i - \bar{x})$$

$$0 = \sum_{i=1}^N (y_i - \bar{y})(x_i - \bar{x}) - \hat{\beta} \sum_{i=1}^N (x_i - \bar{x})^2$$

$$\hat{\beta} = \frac{\sum_{i=1}^N (y_i - \bar{y})(x_i - \bar{x})}{\sum_{i=1}^N (x_i - \bar{x})^2} = \frac{Cov(y, x)}{Var(x)}$$

Omitted variables bias

$$\begin{aligned}\hat{\beta} &= \frac{Cov(Y_i, s_i)}{Var(s_i)} \\&= \frac{Cov(\alpha + \beta s_i + \gamma A_i + \epsilon_i, s_i)}{Var(s_i)} \\&= \frac{\beta Cov(s_i, s_i) + \gamma Cov(A_i, s_i) + Cov(\epsilon_i, s_i)}{Var(s_i)} \\&= \beta \frac{Var(s_i)}{Var(s_i)} + \gamma \frac{Cov(A_i, s_i)}{Var(s_i)} + 0 \\&= \beta + \gamma \times \theta_{as}\end{aligned}$$

Removing selection bias without RCT

- The field of causal inference is all about different strategies to remove selection bias
- The first strategy (really, assumption) in this class: **selection on observables** or **conditional independence**

Intuition

Some assumptions:

- $Y_{0i} = \alpha + \eta_i$
- $Y_{1i} - Y_{0i} = \delta$
- There is some set of "controls", x_i , such that $\eta_i = \beta x_i + u_i$ and $E[u_i | x_i] = 0$ (conditional independence assumption, or CIA)

$$\begin{aligned} Y_i &= Y_{1i} \times D_i + Y_{0i} \times (1 - D_i) \\ &= \delta D_i + Y_{0i} D_i + Y_{0i} - Y_{0i} D_i \\ &= \delta D_i + \alpha + \eta_i \\ &= \delta D_i + \alpha + \beta x_i + u_i \end{aligned}$$

ATEs vs regression coefficients

What does this mean?

- OLS puts more weight on observations with treatment D_i "unexplained" by X_i
- "Reverse" weighting such that the proportion of treated units are used to weight the ATU while the proportion of untreated units enter the weights of the ATT
- This is *an* average effect, but probably not the average we want

Matching and Weighting

Goal

Find covariates \mathbf{X}_i such that the following assumptions are plausible:

1. Selection on observables:

$$Y_{0i}, Y_{1i} \perp\!\!\!\perp D_i | \mathbf{X}_i$$

2. Common support:

$$0 < \Pr(D_i = 1 | \mathbf{X}_i) < 1$$

Then we can use \mathbf{X}_i to group observations and use expectations for control as the predicted counterfactuals among treated, and vice versa.

Assumption 1: Selection on Observables

$$E[Y_1|D, X] = E[Y_1|X]$$

In words...nothing unobserved that determines treatment selection and affects your outcome of interest.

Assumption 2: Common Support

Someone of each type must be in both the treated and untreated groups

$$0 < \Pr(D = 1|X) < 1$$

Causal inference with observational data

With selection on observables and common support:

1. Matching estimators
 2. Regression estimators
- What's the difference?

Estimation options

- Matching
- Weighting
- Regression
- Doubly-robust weighting + regression (won't cover)

Matching: The process

1. For each observation i , find the m "nearest" neighbors, $J_m(i)$.
2. Impute $\hat{Y}_i(0)$ and $\hat{Y}_i(1)$ for each observation:

$$\hat{Y}_i(0) = \begin{cases} Y_i & \text{if } W_i = 0 \\ \frac{1}{m} \sum_{j \in J_m(i)} Y_j & \text{if } W_i = 1 \end{cases}$$

$$\hat{Y}_i(1) = \begin{cases} Y_i & \text{if } W_i = 1 \\ \frac{1}{m} \sum_{j \in J_m(i)} Y_j & \text{if } W_i = 0 \end{cases}$$

3. Form "matched" ATE:

$$\hat{\delta}^{\text{match}} = \frac{1}{N} \sum_{i=1}^N \left(\hat{Y}_i(1) - \hat{Y}_i(0) \right)$$

Matching: Defining "nearest"

1. Euclidean distance:

$$\sum_{k=1}^K (X_{ik} - X_{jk})^2$$

2. Scaled Euclidean distance:


$$\sum_{k=1}^K \frac{1}{\sigma_{X_k}^2} (X_{ik} - X_{jk})^2$$

3. Mahalanobis distance:

$$(X_i - X_j)' \Sigma_X^{-1} (X_i - X_j)$$

Animation for matching

The Effect of Treatment on Y while Matching on X (with a caliper)
1. Start with raw data.



Weighting

1. Estimate propensity score `ps <- glm(W~X, family=binomial, data)`, denoted $\hat{\pi}(X_i)$
2. Weight by inverse of propensity score

$$\hat{\mu}_1 = \frac{\sum_{i=1}^N \frac{Y_i W_i}{\hat{\pi}(X_i)}}{\sum_{i=1}^N \frac{W_i}{\hat{\pi}(X_i)}} \text{ and } \hat{\mu}_0 = \frac{\sum_{i=1}^N \frac{Y_i (1-W_i)}{1-\hat{\pi}(X_i)}}{\sum_{i=1}^N \frac{1-W_i}{1-\hat{\pi}(X_i)}}$$

3. Form "inverse-propensity weighted" ATE:

$$\hat{\delta}^{IPW} = \hat{\mu}_1 - \hat{\mu}_0$$

Regression

1. Regress Y_i on X_i among $W_i = 1$ to form $\hat{\mu}_1(X_i)$
2. Regress Y_i on X_i among $W_i = 0$ to form $\hat{\mu}_0(X_i)$
3. Form difference in predictions:

$$\hat{\delta}^{reg} = \frac{1}{N} \sum_{i=1}^N (\hat{\mu}_1(X_i) - \hat{\mu}_0(X_i))$$


Regression

Or estimate in one step,

$$Y_i = \delta W_i + \beta X_i + W_i \times (X_i - \bar{X}) \gamma + \varepsilon_i$$

Animation for regression

The Relationship between Y and X, Controlling for a Binary Variable W
1. Start with raw data. Correlation between X and Y: 0.403



Simulated data

Now let's do some matching, re-weighting, and regression with simulated data:

```
n <- 5000
select.dat <- tibble(
  x = runif(n, 0, 1),
  z = rnorm(n, 0, 1),
  w = (x > 0.65),
  y = -2.5 + 4*w + 1.5*x + rnorm(n, 0, 1),
  w_alt = (x + z > 0.35),
  y_alt = -2.5 + 4*w_alt + 1.5*x + 2.25*z + rnorm(n, 0, 1)
)
```

Simulation: nearest neighbor matching

```
nn.est1 ← Matching::Match(Y=select.dat$y,  
                           Tr=select.dat$w,  
                           X=select.dat$x,  
                           M=1,  
                           Weight=1,  
                           estimand="ATE")
```

```
summary(nn.est1)
```

```
##  
## Estimate ...    4.0175  
## AI SE.....   0.52954  
## T-stat.....   7.5869  
## p.val.....    3.2863e-14  
##  
## Original number of observations..... 5000  
## Original number of treated obs..... 1732  
## Matched number of observations..... 5000  
## Matched number of observations (unweighted). 5016
```

Simulation: nearest neighbor matching

```
nn.est2 ← Matching::Match(Y=select.dat$y,  
                           Tr=select.dat$w,  
                           X=select.dat$x,  
                           M=1,  
                           Weight=2,  
                           estimand="ATE")
```

```
summary(nn.est2)
```

```
##  
## Estimate ...    4.0175  
## AI SE.....   0.52954  
## T-stat.....   7.5869  
## p.val.....    3.2863e-14  
##  
## Original number of observations..... 5000  
## Original number of treated obs..... 1732  
## Matched number of observations..... 5000  
## Matched number of observations (unweighted). 5016
```

Simulation: regression

```
reg1.dat ← select.dat %>% filter(w==1)
reg1 ← lm(y ~ x, data=reg1.dat)

reg0.dat ← select.dat %>% filter(w==0)
reg0 ← lm(y ~ x, data=reg0.dat)
pred1 ← predict(reg1,new=select.dat)
pred0 ← predict(reg0,new=select.dat)
mean(pred1-pred0)
```

```
## [1] 4.076999
```

Violation of selection on observables

NN Matching

```
nn.est3 ← Matching::Match(Y=select.dat$y_alt,  
                           Tr=select.dat$w_alt,  
                           X=select.dat$x,  
                           M=1,  
                           Weight=2,  
                           estimand="ATE")  
  
summary(nn.est3)
```

```
##  
## Estimate...    7.6642  
## AI SE.....  0.052903  
## T-stat..... 144.87  
## p.val.....  < 2.22e-16  
##  
## Original number of observations..... 5000  
## Original number of treated obs..... 2748  
## Matched number of observations..... 5000  
## Matched number of observations (unweighted). 23014
```

Regression

```
reg1.dat ← select.dat %>% filter(w_alt=1)  
reg1 ← lm(y_alt ~ x, data=reg1.dat)  
  
reg0.dat ← select.dat %>% filter(w_alt=0)  
reg0 ← lm(y_alt ~ x, data=reg0.dat)  
pred1_alt ← predict(reg1,new=select.dat)  
pred0_alt ← predict(reg0,new=select.dat)  
mean(pred1_alt-pred0_alt)
```

```
## [1] 7.646532
```

What covariates to use?

- There are such things as "bad controls"

Pricing and Hospital Profit Status

Penalized hospitals

```
final.hcris <- hcris.data %>% ungroup() %>%  
  filter(price_denom>100, !is.na(price_denom),  
         price_num>0, !is.na(price_num),  
         price<1000000,  
         beds>30, year=2012) %>%  
  mutate( hvbp_payment = ifelse(is.na(hvbp_payment),0,hvbp_payment),  
         hrrp_payment = ifelse(is.na(hrrp_payment),0,abs(hrrp_payment)),  
         penalty = (hvbp_payment-hrrp_payment<0))
```

Summary stats

Always important to look at your data before doing any formal analysis. Ask yourself a few questions:


1. Are the magnitudes reasonable?
2. Are there lots of missing values?
3. Are there clear examples of misreporting?

Summary stats

```
summary(hcris.data$price)
```

##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
##	-123697	4783	7113	Inf	10230	Inf	63662


```
plot(density(hcris.data$price, na.rm=TRUE))
```



```
summary(final.hcris$price)
```

##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
##	340.8	6129.9	8705.4	9646.9	11905.4	97688.8

```
plot(density(final.hcris$price))
```



Dealing with problems

We've adopted a very brute force way to deal with outlier prices. Other approaches include:

1. Investigate very closely the hospitals with extreme values
2. Winsorize at certain thresholds (replace extreme values with pre-determined thresholds)
3. Impute prices for extreme hospitals

Differences among penalized hospitals

- Mean price among penalized hospitals: 9,896.31
- Mean price among non-penalized hospitals: 9,560.41
- Mean difference: 335.9


Comparison of hospitals

Are penalized hospitals sufficiently similar to non-penalized hospitals?


Let's look at covariate balance using a love plot, part of the `library(cobalt)` package.

Love plots without adjustment

```
love.plot(bal.tab(lp.covs,treat=lp.vars$penalty), colors="black", shapes="circle", threshold=0.1) +  
  theme_bw() + theme(legend.position="none")
```



Love plots without adjustment



Using matching to improve balance

Some things to think about:

- exact versus nearest neighbor
- with or without ties (and how to break ties)
- measure of distance

1. Exact Matching

```
m.exact ← Matching::Match(Y=lp.vars$price,  
                          Tr=lp.vars$penalty,  
                          X=lp.covs,  
                          M=1,  
                          exact=TRUE)  
print(m.exact)
```


```
## [1] NA  
## attr(,"class")  
## [1] "Match"
```

1. Exact Matching (on a subset)


```
lp.covs2 ← lp.covs %>% select(beds, mcaid_discharges)
m.exact ← Matching::Match(Y=lp.vars$price,
                          Tr=lp.vars$penalty,
                          X=lp.covs2,
                          M=1,
                          exact=TRUE,
                          estimand="ATE")
```

1. Exact Matching (on a subset)

```
love.plot(bal.tab(m.exact, covs = lp.covs2, treat = lp.vars$penalty),  
          threshold=0.1,  
          grid=FALSE, sample.names=c("Unmatched", "Matched"),  
          position="top", shapes=c("circle", "triangle"),  
          colors=c("black", "blue")) +  
theme_bw()
```



1. Exact Matching (on a subset)




2. Nearest neighbor matching (inverse variance)


```
m.nn.var <- Matching::Match(Y=lp.vars$price,  
                           Tr=lp.vars$penalty,  
                           X=lp.covs,  
                           M=4,  
                           Weight=1,  
                           estimand="ATE")  
  
v.name=data.frame(new=c("Beds", "Medicaid Discharges", "Inpatient Charges",  
                        "Medicare Discharges", "Medicare Payments"))
```

2. Nearest neighbor matching (inverse variance)

```
love.plot(bal.tab(m.nn.var, covs = lp.covs, treat = lp.vars$penalty),
          threshold=0.1,
          var.names=v.name,
          grid=FALSE, sample.names=c("Unmatched", "Matched"),
          position="top", shapes=c("circle", "triangle"),
          colors=c("black", "blue")) +
theme_bw()
```



2. Nearest neighbor matching (inverse variance)




2. Nearest neighbor matching (inverse variance)


```
m.nn.var2 ← Matching::Match(Y=lp.vars$price,  
                             Tr=lp.vars$penalty,  
                             X=lp.covs,  
                             M=1,  
                             Weight=1,  
                             estimand="ATE")
```

2. Nearest neighbor matching (inverse variance)

```
love.plot(bal.tab(m.nn.var2, covs = lp.covs, treat = lp.vars$penalty),
  threshold=0.1,
  var.names=v.name,
  grid=FALSE, sample.names=c("Unmatched", "Matched"),
  position="top", shapes=c("circle","triangle"),
  colors=c("black","blue")) +
  theme_bw()
```



2. Nearest neighbor matching (inverse variance)




2. Nearest neighbor matching (Mahalanobis)


```
m.nn.md ← Matching::Match(Y=lp.vars$price,  
                          Tr=lp.vars$penalty,  
                          X=lp.covs,  
                          M=1,  
                          Weight=2,  
                          estimand="ATE")
```

2. Nearest neighbor matching (Mahalanobis)

```
love.plot(bal.tab(m.nn.md, covs = lp.covs, treat = lp.vars$penalty),
          threshold=0.1,
          var.names=v.name,
          grid=FALSE, sample.names=c("Unmatched", "Matched"),
          position="top", shapes=c("circle","triangle"),
          colors=c("black","blue")) +
theme_bw()
```



2. Nearest neighbor matching (Mahalanobis)




2. Nearest neighbor matching (propensity score)


```
logit.model ← glm(penalty ~ beds + mcaid_discharges + ip_charges + mcare_discharges +  
                  tot_mcare_payment, family=binomial, data=lp.vars)  
ps ← fitted(logit.model)  
m.nn.ps ← Matching::Match(Y=lp.vars$price,  
                          Tr=lp.vars$penalty,  
                          X=ps,  
                          M=1,  
                          estimand="ATE")
```


2. Nearest neighbor matching (propensity score)


```
love.plot(bal.tab(m.nn.ps, covs = lp.covs, treat = lp.vars$penalty),  
          threshold=0.1,  
          var.names=v.name,  
          grid=FALSE, sample.names=c("Unmatched", "Matched"),  
          position="top", shapes=c("circle", "triangle"),  
          colors=c("black", "blue")) +  
theme_bw()
```



2. Nearest neighbor matching (propensity score)



3. Weighting



Results: Exact matching

```
##  
## Estimate ... 1777.6  
## AI SE..... 34.725  
## T-stat..... 51.191  
## p.val..... < 2.22e-16  
##  
## Original number of observations..... 2707  
## Original number of treated obs..... 698  
## Matched number of observations..... 12  
## Matched number of observations (unweighted). 12  
##  
## Number of obs dropped by 'exact' or 'caliper' 2695
```

Results: Nearest neighbor

- Inverse variance

```
##  
## Estimate ... -526.95  
## AI SE..... 223.06  
## T-stat..... -2.3623  
## p.val..... 0.01816  
##  
## Original number of observations..... 2707  
## Original number of treated obs..... 698  
## Matched number of observations..... 2707  
## Matched number of observations (unweighted). 2711
```

Results: Nearest neighbor

- Mahalanobis

```
##
## Estimate ...    -492.82
## AI SE.....   223.55
## T-stat.....   -2.2046
## p.val.....    0.027485
##
## Original number of observations..... 2707
## Original number of treated obs..... 698
## Matched number of observations..... 2707
## Matched number of observations (unweighted). 2708
```

Results: Nearest neighbor

- Propensity score

```
##  
## Estimate ... -201.03  
## AI SE..... 275.76  
## T-stat..... -0.72898  
## p.val..... 0.46601  
##  
## Original number of observations..... 2707  
## Original number of treated obs..... 698  
## Matched number of observations..... 2707  
## Matched number of observations (unweighted). 14795
```

Results: IPW weighting

```
lp.vars <- lp.vars %>%  
  mutate(ipw = case_when(  
    penalty=1 ~ 1/ps,  
    penalty=0 ~ 1/(1-ps),  
    TRUE ~ NA_real_  
  ))  
mean.t1 <- lp.vars %>% filter(penalty=1) %>%  
  select(price, ipw) %>% summarize(mean_p=weighted.mean(price,w=ipw))  
mean.t0 <- lp.vars %>% filter(penalty=0) %>%  
  select(price, ipw) %>% summarize(mean_p=weighted.mean(price,w=ipw))  
mean.t1$mean_p - mean.t0$mean_p
```

```
## [1] -196.8922
```


Results: IPW weighting with regression

```
ipw.reg <- lm(price ~ penalty, data=lp.vars, weights=ipw)
summary(ipw.reg)

##
## Call:
## lm(formula = price ~ penalty, data = lp.vars, weights = ipw)
##
## Weighted Residuals:
##      Min       1Q   Median       3Q      Max
## -18691  -4802  -1422    2651   94137
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   9876.4      147.8   66.808  <2e-16 ***
## penaltyTRUE   -196.9      211.2   -0.932    0.351
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7829 on 2705 degrees of freedom
## Multiple R-squared:  0.0003211,    Adjusted R-squared:  -4.85e-05
## F-statistic: 0.8688 on 1 and 2705 DF,  p-value: 0.3514
```

Results: Regression

```
reg1.dat <- lp.vars %>% filter(penalty=1, complete.cases(.))
reg1 <- lm(price ~ beds+ mcaid_discharges + ip_charges + mcare_discharges +
           tot_mcare_payment, data=reg1.dat)

reg0.dat <- lp.vars %>% filter(penalty=0, complete.cases(.))
reg0 <- lm(price ~ beds + mcaid_discharges + ip_charges + mcare_discharges +
           tot_mcare_payment, data=reg0.dat)
pred1 <- predict(reg1,new=lp.vars)
pred0 <- predict(reg0,new=lp.vars)
mean(pred1-pred0)

## [1] -5.845761
```

Results: Regression in one step

```
reg.dat <- lp.vars %>% ungroup() %>% filter(complete.cases(.)) %>%  
  mutate(beds_diff = penalty*(beds - mean(beds)),  
         mcaid_diff = penalty*(mcaid_discharges - mean(mcaid_discharges)),  
         ip_diff = penalty*(ip_charges - mean(ip_charges)),  
         mcare_diff = penalty*(mcare_discharges - mean(mcare_discharges)),  
         mpay_diff = penalty*(tot_mcare_payment - mean(tot_mcare_payment)))  
reg <- lm(price ~ penalty + beds + mcaid_discharges + ip_charges + mcare_discharges + tot_mcare_payment +  
         beds_diff + mcaid_diff + ip_diff + mcare_diff + mpay_diff,  
         data=reg.dat)
```

Results: Regression in one step

```
##
## Call:
## lm(formula = price ~ penalty + beds + mcaid_discharges + ip_charges +
##      mcare_discharges + tot_mcare_payment + beds_diff + mcaid_diff +
##      ip_diff + mcare_diff + mpay_diff, data = reg.dat)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -38175  -2900   -597    2105   67409
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    8.466e+03  1.711e+02  49.482  < 2e-16 ***
## penaltyTRUE    -5.846e+00  2.124e+02  -0.028  0.97804
## beds           1.107e+00  1.421e+00   0.779  0.43618
## mcaid_discharges -4.714e-01  7.296e-02  -6.462  1.23e-10 ***
## ip_charges       6.426e-06  1.285e-06   5.002  6.04e-07 ***
## mcare_discharges -8.122e-01  9.257e-02  -8.774  < 2e-16 ***
## tot_mcare_payment 9.502e-05  6.858e-06  13.857  < 2e-16 ***
## beds_diff       2.517e+00  2.986e+00   0.843  0.39931
## mcaid_diff       1.058e-01  1.570e-01   0.674  0.50050
## ip_diff         -4.534e-06  2.027e-06  -2.237  0.02539 *
## mcare_diff       4.806e-01  1.809e-01   2.657  0.00793 **
## mpay_diff       -5.452e-05  1.321e-05  -4.128  3.78e-05 ***
## ---
```

Summary of ATEs

1. Exact matching: 1777.63
2. NN matching, inverse variance: -526.95
3. NN matching, mahalanobis: -492.82
4. NN matching, pscore: -201.03
5. Inverse pscore weighting: -196.89
6. IPW regression: -196.89
7. Regression: -5.85
8. Regression 1-step: -5.85

So what have we learned?

Key assumptions for causal inference

1. Selection on observables
2. Common support

These become more nuanced but the intuition is the same in almost all questions of causal inference.

Causal effect assuming selection on observables

If we assume selection on observables holds, then we only need to condition on the relevant covariates to identify a causal effect. But we still need to ensure common support...

1. Matching
2. Reweighting
3. Regression