Physician Behaviors and Hospital Influence

Haizhen Lin & Ian McCarthy & Michael Richards

University of Nevada, Las Vegas

December 7, 2018

Background

Physician with decision-making authority for treatment

- Information asymmetry
- Regulatory restrictions

Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use

Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use

 \longrightarrow Incentives for hospitals to influence physicians


Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use

 \longrightarrow Incentives for hospitals to influence physicians


Most direct way (arguably) is to purchase physician practice

Changing Physician Relationships



Richards et al., Medical Care, 2016

Changing Physician Relationships



Baker, Bundorf, and Kessler, Health Affairs, 2014

What do we expect from integration?

- Hospitals claim efficiency gains via reduced fragmentation and increased coordination
- Increased bargaining power
- Financial incentives for cost increases and decreases
 - Lower costs with fixed payment versus higher costs/payments from private insurance
 - Change location of care
 - More resources due to pay-for-performance

In context

- Physician agency (Clemens & Gottlieb 2014, AER; Afendulis & Kessler 2007, AER; Gruber & Owings 1996, RAND; Iizuka 2012, AER)
- Vertical integration (Cuellar & Gertler 2006, JHE; Ciliberto & Dranove 2006, JHE; Baker et al. 2016, JHE; Koch et al. 2017, JHE)

1. Motivestimation

- 1. Motivestimation
- 2. Initial Results

- 1. Motivestimation
- 2. Initial Results
- 3. Event Study

- 1. Motivestimation
- 2. Initial Results
- 3. Event Study
- 4. Instrumental Variables

- 1. Motivestimation
- 2. Initial Results
- 3. Event Study
- 4. Instrumental Variables
- 5. Other Outcomes

Theoretical Framework

Observed care at time t is

$$y_{ijk} = \arg\max_{y} \theta_{u} \tilde{u}\left(y; \Gamma_{j}, \kappa_{i}\right) + \theta_{\pi} \pi\left(y; \Gamma_{k}, \Gamma_{j}\right).$$

Observed care at time t is

$$y_{ijk} = \arg \max_{y} \theta_u \tilde{u}(y; \Gamma_j, \kappa_i) + \theta_{\pi} \pi(y; \Gamma_k, \Gamma_j).$$


With assumptions on linearity and separability in patient preferences:

$$y_{ijk} = \boxed{\alpha_i + x_i \beta} + \boxed{\Gamma_{jk}} + \epsilon_{ijk}$$

Observed care at time t is

$$y_{ijk} = \arg \max_{y} \theta_{u} \tilde{u} (y; \Gamma_{j}, \kappa_{i}) + \theta_{\pi} \pi (y; \Gamma_{k}, \Gamma_{j}).$$

With assumptions on linearity and separability in patient preferences:




Patient Preferences

Observed care at time t is

$$y_{ijk} = \arg\max_{y} \theta_{u} \tilde{u}\left(y; \Gamma_{j}, \kappa_{i}\right) + \theta_{\pi} \pi\left(y; \Gamma_{k}, \Gamma_{j}\right).$$

With assumptions on linearity and separability in patient preferences:



6

Suggests two-step estimation strategy:

Suggests two-step estimation strategy:

1. Estimate $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$ at patient level (separately by year). This isolates variation in care to physicians and hospitals (not patients).

7

Suggests two-step estimation strategy:


- 1. Estimate $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$ at patient level (separately by year). This isolates variation in care to physicians and hospitals (not patients).
- 2. Estimate $\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt}$ with physician-hospital panel. This further isolates variation to physician-hospital interaction.

- Draws from "match values" in labor literature (Abowd et al., 2002; Card et al., 2013, QJE)
- Exploits variation across inpatient stays and splits the separation of match value into two steps
- Identifies effects on match value from within-physician variation across hospitals (e.g., patient movers in Finkelstein et al., 2016, QJE)

Traditional "match value" approach:


$$y_{ijk} = \alpha_i + x_i \beta + \boxed{\Gamma_j} + \boxed{\Gamma_k} + \boxed{\Gamma_{jk}} + \epsilon_{ijk}$$

Traditional "match value" approach:




Physician effect

Traditional "match value" approach:



7

Traditional "match value" approach:




7

Our approach:

$$y_{ijk} = \alpha_i + x_i \beta + \underbrace{\Gamma_{jk}^t}_{\Gamma_j} + \epsilon_{ijk}$$


$$\Gamma_j + \underbrace{\Gamma_k}_{\Gamma_k} + \underbrace{z_{jkt} \delta}$$

Our approach:



7


Our approach:




Physician, hospital, and match effect (jointly)

Physician effect -

Our approach:



Our approach:



7

Intuition

- Hospital influence on physicians is an interaction effect
- Physician agency is net of patient preference

Data

Data Sources

- CMS: 100% inpatient and institutional outpatient Medicare claims data (2008-2015)
- SK&A: Hospital ownership of physician practices
- AHA, HCRIS, POS: Hospital characteristics
- Annual IPPS Impact Files: Hospital cost-to-charge ratios (CCR)
- ACS: County-level demographics, education, income, and employment

 Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
- Drop physicians with NPIs not matched in the SK&A data

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
- Drop physicians with NPIs not matched in the SK&A data
- Drop lowest/highest 1% of charges and patients < 65 years old

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
- Drop physicians with NPIs not matched in the SK&A data
- $\bullet\,$ Drop lowest/highest 1% of charges and patients < 65 years old
- → 518,398 unique observations at the physician/hospital/year
- \longrightarrow 7.5mm inpatient stays (47% of total) and 24mm outpatient procedures

Estimation of Match Values

Specification

Two-step estimation strategy:

- 1. Estimate $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$ at patient level (separately by year)
- 2. Estimate $\hat{\Gamma}_{jkt}=\gamma_j+\gamma_k+\tau_t+z_{jkt}\delta+\eta_{jkt}$ with physician-hospital panel

Specification

$$y_{ijk} = \alpha_i + x_i \beta + \Gamma_{jk} + \epsilon_{ijk},$$

Outcomes

$$\mathbf{y_{ijk}} = \alpha_i + x_i \beta + \Gamma_{jk} + \epsilon_{ijk},$$

- Total inpatient and outpatient Medicare payments
- Total inpatient and outpatient hospital costs (from cost-to-charge ratios)
- Inpatient hospital costs
- Inpatient length of stay
- Outpatient hospital costs

$$y_{ijk} = \frac{\alpha_i}{\epsilon} + x_i \beta + \Gamma_{jk} + \epsilon_{ijk},$$

- Quartiles of total "other" Medicare payments and procedures
- Covers 2008 through 2015 period
- Beneficiary-specific measure of "utilization"

$$y_{ijk} = \alpha_i + \frac{\mathbf{x_i}}{\beta} + \Gamma_{jk} + \epsilon_{ijk},$$

- Age, gender, race
- Indicators for ICD9 diagnosis code groups (18 diagnosis groups per variable plus missing group)
- Indicators for primary DRGs (with at least 1000 observations in a given year)
- Minor differences between total, inpatient, and outpatient specifications

Summary of Match Values

1. Calculate Cost Differential

Apply minimum cost physician-hospital combination to all of physician j's patients:


$$\begin{split} \Delta_k y_{ij} &= \hat{y}_{ijk} - \hat{y}_{ij\underline{\mathbf{k}}} \\ &= \hat{\alpha}_i + x_i \hat{\beta} + \hat{\Gamma}_{jk} - \hat{\alpha}_i - x_i \hat{\beta} - \min\left\{\Gamma_{j1}, ..., \Gamma_{jK}\right\} \\ &= \hat{\Gamma}_{jk} - \min\left\{\Gamma_{j1}, ..., \Gamma_{jK}\right\}. \end{split}$$

Summary of Match Values


2. Summarize

- Total cost differential for each physician
- Limit to pairs with 5 or more procedures
- Limit to physicians with 2 or more hospitals in a year
- Present interquartile range and mean


Within-physician Variation in Payments



Within-physician Variation in Payments



Within-physician Variation in Payments



Estimation of Hospital Influence

Specification

Two-step estimation strategy:

- 1. Estimate $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$ at patient level (separately by year)
- 2. Estimate $\hat{\Gamma}_{jkt}=\gamma_j+\gamma_k+\tau_t+z_{jkt}\delta+\eta_{jkt}$ with physician-hospital panel

Specification

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152	8,171	8,501	8,941	9,169	8,094
	(7,595)	(8,472)	(8,290)	(8,724)	(8,755)	(8,228)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152	8,171	8,501	8,941	9,169	8,094
	(7,595)	(8,472)	(8,290)	(8,724)	(8,755)	(8,228)
Total Costs	9,387	11,323	11,756	12,237	12,736	10,965
	(9,632)	(10,954)	(10,906)	(11,549)	(11,728)	(10,626)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152	8,171	8,501	8,941	9,169	8,094
	(7,595)	(8,472)	(8,290)	(8,724)	(8,755)	(8,228)
Total Costs	9,387	11,323	11,756	12,237	12,736	10,965
	(9,632)	(10,954)	(10,906)	(11,549)	(11,728)	(10,626)
Inpatient Costs	13,655	16,958	17,711	18,367	19,081	16,294
	(7,752)	(9,407)	(9,612)	(9,997)	(10,184)	(9,256)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152	8,171	8,501	8,941	9,169	8,094
	(7,595)	(8,472)	(8,290)	(8,724)	(8,755)	(8,228)
Total Costs	9,387	11,323	11,756	12,237	12,736	10,965
	(9,632)	(10,954)	(10,906)	(11,549)	(11,728)	(10,626)
Inpatient Costs	13,655	16,958	17,711	18,367	19,081	16,294
	(7,752)	(9,407)	(9,612)	(9,997)	(10,184)	(9,256)
Inpatient LOS	5.984	6.021	6.002	6.062	6.029	5.960
	(2.427)	(2.493)	(2.494)	(2.513)	(2.492)	(2.436)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152	8,171	8,501	8,941	9,169	8,094
	(7,595)	(8,472)	(8,290)	(8,724)	(8,755)	(8,228)
Total Costs	9,387	11,323	11,756	12,237	12,736	10,965
	(9,632)	(10,954)	(10,906)	(11,549)	(11,728)	(10,626)
Inpatient Costs	13,655	16,958	17,711	18,367	19,081	16,294
	(7,752)	(9,407)	(9,612)	(9,997)	(10,184)	(9,256)
Inpatient LOS	5.984	6.021	6.002	6.062	6.029	5.960
	(2.427)	(2.493)	(2.494)	(2.513)	(2.492)	(2.436)
Outpatient Costs	3,007	3,806	4,014	4,190	4,361	3,693
	(2,135)	(2,782)	(2,925)	(3,096)	(3,195)	(2,749)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130	0.206	0.233	0.255	0.332	0.196
	(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130	0.206	0.233	0.255	0.332	0.196
	(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)
Physician FTE	24.23	28.59	31.14	31.74	33.13	28.43
	(99.28)	(109.8)	(120.5)	(120.0)	(119.5)	(110.9)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

2008	2012	2013	2014	2015	Overall
0.130	0.206	0.233	0.255	0.332	0.196
(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)
24.23	28.59	31.14	31.74	33.13	28.43
(99.28)	(109.8)	(120.5)	(120.0)	(119.5)	(110.9)
25.77	28.45	29.13	30.69	30.97	28.08
(108.2)	(120.4)	(121.4)	(125.9)	(127.8)	(117.8)
	0.130 (0.336) 24.23 (99.28) 25.77	0.130 0.206 (0.336) (0.404) 24.23 28.59 (99.28) (109.8) 25.77 28.45	0.130 0.206 0.233 (0.336) (0.404) (0.422) 24.23 28.59 31.14 (99.28) (109.8) (120.5) 25.77 28.45 29.13	0.130 0.206 0.233 0.255 (0.336) (0.404) (0.422) (0.436) 24.23 28.59 31.14 31.74 (99.28) (109.8) (120.5) (120.0) 25.77 28.45 29.13 30.69	0.130 0.206 0.233 0.255 0.332 (0.336) (0.404) (0.422) (0.436) (0.471) 24.23 28.59 31.14 31.74 33.13 (99.28) (109.8) (120.5) (120.0) (119.5) 25.77 28.45 29.13 30.69 30.97

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130	0.206	0.233	0.255	0.332	0.196
	(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)
Physician FTE	24.23	28.59	31.14	31.74	33.13	28.43
	(99.28)	(109.8)	(120.5)	(120.0)	(119.5)	(110.9)
Resident FTE	25.77	28.45	29.13	30.69	30.97	28.08
	(108.2)	(120.4)	(121.4)	(125.9)	(127.8)	(117.8)
Nurse FTE	340.8	365.7	369.1	384.9	402.7	364.8
	(446.8)	(487.8)	(494.8)	(519.1)	(550.7)	(487.3)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130	0.206	0.233	0.255	0.332	0.196
	(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)
Physician FTE	24.23	28.59	31.14	31.74	33.13	28.43
	(99.28)	(109.8)	(120.5)	(120.0)	(119.5)	(110.9)
Resident FTE	25.77	28.45	29.13	30.69	30.97	28.08
	(108.2)	(120.4)	(121.4)	(125.9)	(127.8)	(117.8)
Nurse FTE	340.8	365.7	369.1	384.9	402.7	364.8
	(446.8)	(487.8)	(494.8)	(519.1)	(550.7)	(487.3)
Other FTE	749.9	763.0	761.8	776.4	806.0	762.8
	(975.5)	(1032.4)	(1076.2)	(1101.5)	(1157.2)	(1037.4)
	'					'

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130	0.206	0.233	0.255	0.332	0.196
	(0.336)	(0.404)	(0.422)	(0.436)	(0.471)	(0.397)
Physician FTE	24.23	28.59	31.14	31.74	33.13	28.43
	(99.28)	(109.8)	(120.5)	(120.0)	(119.5)	(110.9)
Resident FTE	25.77	28.45	29.13	30.69	30.97	28.08
	(108.2)	(120.4)	(121.4)	(125.9)	(127.8)	(117.8)
Nurse FTE	340.8	365.7	369.1	384.9	402.7	364.8
	(446.8)	(487.8)	(494.8)	(519.1)	(550.7)	(487.3)
Other FTE	749.9	763.0	761.8	776.4	806.0	762.8
	(975.5)	(1032.4)	(1076.2)	(1101.5)	(1157.2)	(1037.4)
Beds (100s)	1.980	1.967	1.958	1.982	2.009	1.976
	(2.160)	(2.142)	(2.137)	(2.172)	(2.235)	(2.154)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Practice Size	13.73	17.31	17.31	17.82	18.41	16.10
	(32.10)	(30.70)	(29.28)	(28.46)	(28.02)	(30.05)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Practice Size	13.73	17.31	17.31	17.82	18.41	16.10
	(32.10)	(30.70)	(29.28)	(28.46)	(28.02)	(30.05)
Experience	22.55		23.94			23.17
	(6.496)	(6.703)	(6.950)	(6.902)	(6.989)	(6.746)

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \mathbf{z}_{jkt}\delta + \eta_{jkt},$$

2008	2012	2013	2014	2015	Overall
13.73	17.31	17.31	17.82	18.41	16.10
(32.10)	(30.70)	(29.28)	(28.46)	(28.02)	(30.05)
22.55	23.00	23.94	23.65	24.77	23.17
(6.496)	(6.703)	(6.950)	(6.902)	(6.989)	(6.746)
0.249	0.248	0.266	0.284	0.344	0.264
0.452	0.501	0.507	0.508	0.454	0.480
	13.73 (32.10) 22.55 (6.496) 0.249	13.73 17.31 (32.10) (30.70) 22.55 23.00 (6.496) (6.703) 0.249 0.248	13.73 17.31 17.31 (32.10) (30.70) (29.28) 22.55 23.00 23.94 (6.496) (6.703) (6.950) 0.249 0.248 0.266	13.73 17.31 17.31 17.82 (32.10) (30.70) (29.28) (28.46) 22.55 23.00 23.94 23.65 (6.496) (6.703) (6.950) (6.902) 0.249 0.248 0.266 0.284	13.73 17.31 17.31 17.82 18.41 (32.10) (30.70) (29.28) (28.46) (28.02) 22.55 23.00 23.94 23.65 24.77 (6.496) (6.703) (6.950) (6.902) (6.989) 0.249 0.248 0.266 0.284 0.344

Estimated Effects of Vertical Integration

Outcome Estimate St. Error

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments	108.293**	(43.335)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs	108.293** 235.404***	(43.335) (60.717)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs Inpatient Hospital Costs	108.293** 235.404*** 157.018***	(43.335) (60.717) (49.787)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs Inpatient Hospital Costs Inpatient Length of Stay	108.293** 235.404*** 157.018*** -0.014	(43.335) (60.717) (49.787) (0.017)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments	108.293**	(43.335)
Total Hospital Costs	235.404***	(60.717)
Inpatient Hospital Costs	157.018***	(49.787)
Inpatient Length of Stay	-0.014	(0.017)
Outpatient Hospital Costs	-47.940***	(18.892)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Effects on Components of Inpatient Stay

Outcome	Estimate	St. Error
Charges for:		
Medical Supplies	32.969	(30.076)
Operating Room	-7.492	(23.023)
Anesthesia	5.715	(5.044)
Labs	10.324	(8.767)
Radiology	-3.843	(6.035)
MRI	-0.207	(1.355)

^{*} p-value $<\!0.1,$ ** p-value $<\!0.05,$ *** p-value $<\!0.01$

Effects on Components of Inpatient Stay

Outcome	Estimate	St. Error
Counts of:		
Private Days	-0.003	(0.012)
ICU Days	0.023*	(0.013)
Radiology	-0.00	(0.00)
Procedures	0.028***	(0.009)
* 1 01	k-t- 1	0.05 *** 0.04

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Threats to Identification and Interpretation

Estimator is effectively a two-way fixed effects DD with time varying treatment


Threats to Identification and Interpretation

Estimator is effectively a two-way fixed effects DD with time varying treatment


Potential Problems

- 1. Vertical integration due to time-varying unobservables & outcomes (standard DD concern)
- 2. Weighted average of all 2×2 DD estimates, with some potentially negative weights


Event Study: Total Medicare Payments



Event Study: Total Hospital Costs



Event Study: Inpatient Hospital Costs



Takeaways

- Evidence of increase in payments and costs
- Evidence consistent with common trends assumption for total payments and costs
- Some concern about common trends for inpatient costs

Integration could be driven by:

- Unobserved, time-varying practice characteristics
- Existing costs and treatment patterns

1. Set of possible physician-hospital pairs

Form set of all hospitals where physician operates from 2008-2015

2. Estimate probability of integration

$$\Pr(I_{jk} = 1) = \frac{\exp(\lambda z_{jk})}{1 + \exp(\lambda z_{jk})}$$

- Hospital and practice characteristics
- Average differential distance (relative to nearest hospital in patient choice set)
- Differential distance interacted with hospital and practice characteristics

2. Estimate probability of integration

$$\hat{\Pr}\left(I_{jk}=1\right) = \frac{\exp\left(\hat{\lambda}z_{jk}\right)}{1 + \exp\left(\hat{\lambda}z_{jk}\right)}$$

Intuition: Physicians less likely to seek/allow acquisition if patients live further away

2. Estimate probability of integration

$$\hat{\Pr}\left(I_{jk}=1\right) = \frac{\exp\left(\hat{\lambda}z_{jk}\right)}{1 + \exp\left(\hat{\lambda}z_{jk}\right)}$$

Intuition: Physicians less likely to seek/allow acquisition if patients live further away

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \underbrace{I_{jkt}}_{\hat{I}_{jkt}} \delta_1 + \tilde{z}_{jkt} \delta_2 + \eta_{jkt},$$

$$\hat{I}_{jkt} = \hat{\Pr}(I_{jkt} = 1)$$

Outcome Estimate St. Error

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Total Medicare Payments	1028.994**	(498.917)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs	1028.994** 3238.716***	(498.917) (697.512)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs Inpatient Hospital Costs	1028.994** 3238.716*** 2922.754***	(498.917) (697.512) (532.646)

^{*} p-value $<\!0.1,$ ** p-value $<\!0.05,$ *** p-value $<\!0.01$

Outcome	Estimate	St. Error
Total Medicare Payments Total Hospital Costs Inpatient Hospital Costs Inpatient Length of Stay	1028.994** 3238.716*** 2922.754*** 0.322*	(498.917) (697.512) (532.646) (0.181)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Total Medicare Payments	1028.994**	(498.917)
Total Hospital Costs	3238.716***	(697.512)
Inpatient Hospital Costs	2922.754***	(532.646)
Inpatient Length of Stay	0.322*	(0.181)
Outpatient Hospital Costs	-240.470	(200.800)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

IV Results: Components of Inpatient Stay

Outcome	Estimate	St. Error
Charges for:		
Medical Supplies	88.308	(310.707)
Operating Room	914.093***	(237.412)
Anesthesia	126.673***	(48.727)
Labs	-5.867	(110.491)
Radiology	-143.188**	(66.940)
MRI	-41.617***	(15.833)
-t- 1 0 1 -tt-		1 0 0 1

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

IV Results: Components of Inpatient Stay

Outcome	Estimate	St. Error
Counts of:		
Private Days	-0.343**	(0.141)
ICU Days	-0.060	(0.155)
Radiology	-0.054***	(0.016)
Procedures	0.783***	(0.096)
*	**	*** 0.01

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Allocation of Procedures and

Patients

Other Effects

Other ways integration posited to affect physician behavior:

- More procedures overall (not per patient)
- Reallocating procedures from other hospitals
- Reallocating procedures across inpatient and outpatient settings
- Changing patient profile

Outcome Estimate St. Error	Outcome	Estimate	St. Error
----------------------------	---------	----------	-----------

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share	0.081***	(0.003)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share	0.081***	(0.003)
Physician's outpatient share	0.057***	(0.003)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share	0.081***	(0.003)
Physician's outpatient share	0.057***	(0.003)
Total patients	7.079***	(0.505)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share Physician's outpatient share Total patients Inpatient procedures	0.081*** 0.057*** 7.079*** 1.102***	(0.003) (0.003) (0.505) (0.161)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share Physician's outpatient share	0.081*** 0.057***	(0.003)
Total patients	7.079***	(0.505)
Inpatient procedures	1.102***	(0.161)
Outpatient procedures	10.069***	(1.015)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share	0.081***	(0.003)
Physician's outpatient share	0.057***	(0.003)
Total patients	7.079***	(0.505)
Inpatient procedures	1.102***	(0.161)
Outpatient procedures	10.069***	(1.015)
Patient Procedures	0.056	(0.179)

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Outcome	Estimate	St. Error
Physician's inpatient share	0.081***	(0.003)
Physician's outpatient share	0.057***	(0.003)
Total patients	7.079***	(0.505)
Inpatient procedures	1.102***	(0.161)
Outpatient procedures	10.069***	(1.015)
Patient Procedures	0.056	(0.179)
Patient Payments	-373.713	(293.133)
		` ,

^{*} p-value <0.1, ** p-value <0.05, *** p-value <0.01

Effects per Patient

- Increase in Medicare payments (\$110 to \$300) and hospital costs (\$235-\$500)
- Extrapolates to between \$77 and \$210 million in added Medicare payments from vertical integration
- Components of inpatient stays suggest more procedures per patient

Sensitivity

- Event study consistent with common trends for Medicare payments and total hospital costs
- Calculation of 2×2 DD weights suggests relatively small portion of negative weights (70% positive weights)
- No improvement in quality (mortality)
- As falsification test, no effects on payments or DRG weights per inpatient stay

Effects on Total Patients and Allocation of Procedures

- More procedures going to acquiring hospital
- New procedures predominantly coming from outpatient side (10 to 1)
- Perhaps higher usage, lower intensity patient selection (but insignificant)

Interpreting Main Results

- Total within-physician variation in Medicare payments of around \$140,000 per physician per year
- Increases due to vertical integration of between \$110 and \$300 per patient per year
- 5-13% of within-physician variation explained by vertical integration

Thank You