

# Physician Behaviors and Hospital Influence

---

Haizhen Lin & **Ian McCarthy** & Michael Richards

April 20, 2019

## Background

---

# Physician Agency

Physician with decision-making authority for treatment

- Information asymmetry
- Regulatory restrictions

## Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use
- Hospital as residual claimant on billable physician services

Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use
- Hospital as residual claimant on billable physician services

→ Incentives for hospitals to influence physicians


Differential financial incentives between physician and hospital

- More procedures = more revenue, but location of procedure may matter to hospital
- Hospital wants less cost with fixed payment, but physician dictates resource use
- Hospital as residual claimant on billable physician services

—→ Incentives for hospitals to influence physicians


Most direct way (arguably) is to purchase physician practice

# Changing Physician Relationships



Richards *et al.*, Medical Care, 2016

# Changing Physician Relationships



Baker, Bundorf, and Kessler, Health Affairs, 2014



- Physician agency (Clemens & Gottlieb 2014, AER; Afendulis & Kessler 2007, AER; Gruber & Owings 1996, RAND; Iizuka 2012, AER)
- Supply-side variation (Finkelstein *et al.* 2016, QJE; Molitor 2018, AEJ: Policy)
- Vertical integration (Cuellar & Gertler 2006, JHE; Ciliberto & Dranove 2006, JHE; Baker *et al.* 2016, JHE; Koch *et al.* 2017, JHE)

# Theoretical Framework

---

Observed care at time  $t$  is

$$y_{ijk} = \arg \max_y \theta_u \tilde{u}(y; \Gamma_k, \Gamma_j, \kappa_i) + \theta_\pi \pi(y; \Gamma_k, \Gamma_j, \kappa_i).$$

Observed care at time  $t$  is

$$y_{ijk} = \arg \max_y \theta_u \tilde{u}(y; \Gamma_k, \Gamma_j, \kappa_i) + \theta_\pi \pi(y; \Gamma_k, \Gamma_j, \kappa_i).$$

With assumptions on linearity and separability in patient preferences:

$$y_{ijk} = \alpha_i + x_i \beta + \Gamma_{jk} + \epsilon_{ijk}$$


Observed care at time  $t$  is

$$y_{ijk} = \arg \max_y \theta_u \tilde{u}(y; \Gamma_k, \Gamma_j, \kappa_i) + \theta_\pi \pi(y; \Gamma_k, \Gamma_j, \kappa_i).$$

With assumptions on linearity and separability in patient preferences:

$$y_{ijk} = \alpha_i + x_i \beta + \Gamma_{jk} + \epsilon_{ijk}$$

Patient Preferences



Observed care at time  $t$  is

$$y_{ijk} = \arg \max_y \theta_u \tilde{u}(y; \Gamma_k, \Gamma_j, \kappa_i) + \theta_\pi \pi(y; \Gamma_k, \Gamma_j, \kappa_i).$$

With assumptions on linearity and separability in patient preferences:

$$y_{ijk} = \alpha_i + x_i \beta + \Gamma_{jk} + \epsilon_{ijk}$$

Patient Preferences

Physician and hospital characteristics

# Estimation Strategy

Suggests two-step estimation strategy:

# Estimation Strategy

Suggests two-step estimation strategy:

1. Estimate  $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$  at patient level (separately by year). This isolates variation in care to physicians and hospitals (not patients).



# Estimation Strategy

Suggests two-step estimation strategy:

1. Estimate  $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$  at patient level (separately by year). This isolates variation in care to physicians and hospitals (not patients).
2. Estimate  $\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt}$  with physician-hospital panel. This further isolates variation to physician-hospital interaction.

## Estimation Strategy

- Draws from “match values” in labor literature (Abowd *et al.*, 2002; Card *et al.*, 2013, QJE )
- Exploits variation across inpatient stays and splits the separation of match value into two steps
- Identifies effects on match value from within-physician variation across hospitals (e.g., patient movers in Finkelstein *et al.*, 2016, QJE)


# Estimation Strategy

Traditional “match value” approach:

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_j + \Gamma_k + \Gamma_{jk} + \epsilon_{ijk}$$

# Estimation Strategy

Traditional “match value” approach:

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_j + \Gamma_k + \Gamma_{jk} + \epsilon_{ijk}$$


Physician effect

# Estimation Strategy

Traditional “match value” approach:


$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_j + \Gamma_k + \Gamma_{jk} + \epsilon_{ijk}$$

Physician effect

Hospital effect

# Estimation Strategy

Traditional “match value” approach:

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_j + \Gamma_k + \Gamma_{jk} + \epsilon_{ijk}$$


Physician effect

Hospital effect

Physician-hospital match value


# Estimation Strategy

Our approach:

$$y_{ijk} = \alpha_i + x_i\beta + \underbrace{\Gamma_{jk}^t}_{\Gamma_j + \Gamma_k + z_{jkt}\delta} + \epsilon_{ijk}$$

# Estimation Strategy

Our approach:

$$y_{ijk} = \alpha_i + x_i\beta + \underbrace{\Gamma_{jk}^t}_{\Gamma_j + \Gamma_k + z_{jkt}\delta} + \epsilon_{ijk}$$


Physician, hospital, and match effect (jointly)



# Estimation Strategy

Our approach:

$$y_{ijk} = \alpha_i + x_i\beta + \underbrace{\Gamma_{jk}^t}_{\Gamma_j + \Gamma_k + z_{jkt}\delta} + \epsilon_{ijk}$$

The diagram illustrates the decomposition of the treatment effect term  $\Gamma_{jk}^t$  in the equation above. A bracket under  $\Gamma_{jk}^t$  points to a sum of three components:  $\Gamma_j$  (in a green box),  $\Gamma_k$  (in a red box), and  $z_{jkt}\delta$  (in an orange oval). Arrows from the text labels below point to these components: 'Physician effect' points to  $\Gamma_j$ , and 'Physician, hospital, and match effect (jointly)' points to the entire sum  $\Gamma_j + \Gamma_k + z_{jkt}\delta$ .


Physician, hospital, and match effect (jointly)

Physician effect

# Estimation Strategy

Our approach:

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk}^t + \epsilon_{ijk}$$



Physician, hospital, and match effect (jointly)


Physician effect

Hospital effect

# Estimation Strategy

Our approach:

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk}^t + \epsilon_{ijk}$$



Physician, hospital, and match effect (jointly)

Physician effect

Hospital effect

Physician-hospital integration

- Hospital influence on physicians is an interaction effect
- Potential influence should be net of patient preference

# Data

---

## Data Sources

- CMS: 100% inpatient and institutional outpatient Medicare claims data (2008-2015)
- SK&A: Hospital ownership of physician practices and practice characteristics
- AHA, HCRIS, POS: Hospital characteristics
- Annual IPPS Impact Files: Hospital cost-to-charge ratios (CCR)
- ACS: County-level demographics, education, income, and employment

## Sample Construction

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician

## Sample Construction

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.



## Sample Construction

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
- Drop physicians with NPIs not matched in the SK&A data

## Sample Construction

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
- Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
- Drop physicians with NPIs not matched in the SK&A data
- Drop lowest/highest 1% of charges and patients  $< 65$  years old

## Sample Construction

- Planned inpatient stays (elective admissions initiated by a physician, clinic, or HMO referral) and outpatient procedures with observed NPI for the operating physician
  - Drop physicians operating in hospitals more than 120 miles from primary office or outside of contiguous U.S.
  - Drop physicians with NPIs not matched in the SK&A data
  - Drop lowest/highest 1% of charges and patients  $< 65$  years old
- 518,398 unique observations at the physician/hospital/year
- 7.5mm inpatient stays (47% of total) and 24mm outpatient procedures

## **Preliminary Evidence**


---

# Total Spending by Integration Status

Estimate and plot residual from:

$$y_{jkt} = \beta x_{jt} + \delta z_{kt} + \lambda_k + \lambda_j + \lambda_t + \varepsilon_{jkt}$$

# Total Spending by Integration Status



# Total Spending by Integration Status

Components of aggregate effect:

1. Total number of patients for physician  $k$
2. Change in patient profile
3. Reallocation of patients across hospitals
4. Change in treatment for observationally equivalent patients

## Estimation of Match Values

---



Two-step estimation strategy:

1. Estimate  $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$  at patient level (separately by year)
2. Estimate  $\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt}$  with physician-hospital panel

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk},$$

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk},$$

- Total inpatient and outpatient Medicare payments
- Total inpatient and outpatient hospital costs (from cost-to-charge ratios)

$$y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk},$$

- Quartiles of total “other” Medicare payments and procedures
- Covers 2008 through 2015 period
- Beneficiary-specific ranking of health care utilization

# Independent Variables

$$y_{ijk} = \alpha_i + \mathbf{x}_i\beta + \Gamma_{jk} + \epsilon_{ijk},$$

- Age, gender, race
- Indicators for ICD9 diagnosis code groups (18 diagnosis groups per variable plus missing group)
- Indicators for primary DRGs (with at least 1000 observations in a given year)

# Estimation of Hospital Influence

---

Two-step estimation strategy:

1. Estimate  $y_{ijk} = \alpha_i + x_i\beta + \Gamma_{jk} + \epsilon_{ijk}$  at patient level (separately by year)
2. Estimate  $\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt}$  with physician-hospital panel

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$



# Main Outcomes

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152 (7,595)	8,171 (8,472)	8,501 (8,290)	8,941 (8,724)	9,169 (8,755)	8,094 (8,228)

# Main Outcomes

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Total Payments	7,152 (7,595)	8,171 (8,472)	8,501 (8,290)	8,941 (8,724)	9,169 (8,755)	8,094 (8,228)
Total Costs	9,387 (9,632)	11,323 (10,954)	11,756 (10,906)	12,237 (11,549)	12,736 (11,728)	10,965 (10,626)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)
Physician FTE	24.23 (99.28)	28.59 (109.8)	31.14 (120.5)	31.74 (120.0)	33.13 (119.5)	28.43 (110.9)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)
Physician FTE	24.23 (99.28)	28.59 (109.8)	31.14 (120.5)	31.74 (120.0)	33.13 (119.5)	28.43 (110.9)
Resident FTE	25.77 (108.2)	28.45 (120.4)	29.13 (121.4)	30.69 (125.9)	30.97 (127.8)	28.08 (117.8)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)
Physician FTE	24.23 (99.28)	28.59 (109.8)	31.14 (120.5)	31.74 (120.0)	33.13 (119.5)	28.43 (110.9)
Resident FTE	25.77 (108.2)	28.45 (120.4)	29.13 (121.4)	30.69 (125.9)	30.97 (127.8)	28.08 (117.8)
Nurse FTE	340.8 (446.8)	365.7 (487.8)	369.1 (494.8)	384.9 (519.1)	402.7 (550.7)	364.8 (487.3)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)
Physician FTE	24.23 (99.28)	28.59 (109.8)	31.14 (120.5)	31.74 (120.0)	33.13 (119.5)	28.43 (110.9)
Resident FTE	25.77 (108.2)	28.45 (120.4)	29.13 (121.4)	30.69 (125.9)	30.97 (127.8)	28.08 (117.8)
Nurse FTE	340.8 (446.8)	365.7 (487.8)	369.1 (494.8)	384.9 (519.1)	402.7 (550.7)	364.8 (487.3)
Other FTE	749.9 (975.5)	763.0 (1032.4)	761.8 (1076.2)	776.4 (1101.5)	806.0 (1157.2)	762.8 (1037.4)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Integrated	0.130 (0.336)	0.206 (0.404)	0.233 (0.422)	0.255 (0.436)	0.332 (0.471)	0.196 (0.397)
Physician FTE	24.23 (99.28)	28.59 (109.8)	31.14 (120.5)	31.74 (120.0)	33.13 (119.5)	28.43 (110.9)
Resident FTE	25.77 (108.2)	28.45 (120.4)	29.13 (121.4)	30.69 (125.9)	30.97 (127.8)	28.08 (117.8)
Nurse FTE	340.8 (446.8)	365.7 (487.8)	369.1 (494.8)	384.9 (519.1)	402.7 (550.7)	364.8 (487.3)
Other FTE	749.9 (975.5)	763.0 (1032.4)	761.8 (1076.2)	776.4 (1101.5)	806.0 (1157.2)	762.8 (1037.4)
Beds (100s)	1.980 (2.160)	1.967 (2.142)	1.958 (2.137)	1.982 (2.172)	2.009 (2.235)	1.976 (2.154)



# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Practice Size	13.73 (32.10)	17.31 (30.70)	17.31 (29.28)	17.82 (28.46)	18.41 (28.02)	16.10 (30.05)

# Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Practice Size	13.73 (32.10)	17.31 (30.70)	17.31 (29.28)	17.82 (28.46)	18.41 (28.02)	16.10 (30.05)
Experience	22.55 (6.496)	23.00 (6.703)	23.94 (6.950)	23.65 (6.902)	24.77 (6.989)	23.17 (6.746)

## Independent Variables

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + z_{jkt}\delta + \eta_{jkt},$$

	2008	2012	2013	2014	2015	Overall
Practice Size	13.73 (32.10)	17.31 (30.70)	17.31 (29.28)	17.82 (28.46)	18.41 (28.02)	16.10 (30.05)
Experience	22.55 (6.496)	23.00 (6.703)	23.94 (6.950)	23.65 (6.902)	24.77 (6.989)	23.17 (6.746)
% Multi-Specialty	0.249	0.248	0.266	0.284	0.344	0.264
% Surgery Center	0.452	0.501	0.507	0.508	0.454	0.480

# Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
---------	----------	-----------

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments	75.121**	(30.902)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments	75.121**	(30.902)
Total Hospital Costs	132.466***	(42.026)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## Estimated Effects of Vertical Integration

Outcome	Estimate	St. Error
Total Medicare Payments	75.121**	(30.902)
Total Hospital Costs	132.466***	(42.026)
Total Stays	0.015***	(0.004)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## Threats to Identification and Interpretation

Estimator is effectively a two-way fixed effects DD with time varying treatment




# Threats to Identification and Interpretation

Estimator is effectively a two-way fixed effects DD with time varying treatment


## Potential Problems

1. Vertical integration due to time-varying unobservables & outcomes (standard DD concern)
2. Weighted average of all  $2 \times 2$  DD estimates, with some potentially negative weights

## Event Study: Total Medicare Payments



## Event Study: Total Hospital (IP & OP) Costs



## Takeaways

- Increase in payments and costs
- Evidence consistent with common trends assumption for total payments and costs
- Concerns about limited pre-period data

# Endogeneity of physician-hospital integration

Integration could be driven by:

- Unobserved, time-varying practice characteristics
- Existing costs and treatment patterns

# Endogeneity of physician-hospital integration

## 1. Set of possible physician-hospital pairs

Form set of all hospitals where physician operates from 2008-2015

## 2. Estimate probability of integration

$$\Pr(I_{jk} = 1) = \frac{\exp(\lambda z_{jk})}{1 + \exp(\lambda z_{jk})}$$

- Hospital and practice characteristics
- Average differential distance (relative to nearest hospital in patient choice set)
- Differential distance interacted with hospital and practice characteristics

## 2. Estimate probability of integration

$$\hat{\Pr}(I_{jk} = 1) = \frac{\exp(\hat{\lambda}z_{jk})}{1 + \exp(\hat{\lambda}z_{jk})}$$

Intuition: Physicians less likely to seek/allow acquisition if patients live further away



## 2. Estimate probability of integration

$$\hat{\Pr}(I_{jk} = 1) = \frac{\exp(\hat{\lambda} z_{jk})}{1 + \exp(\hat{\lambda} z_{jk})}$$

Intuition: Physicians less likely to seek/allow acquisition if patients live further away

$$\hat{\Gamma}_{jkt} = \gamma_j + \gamma_k + \tau_t + \underbrace{I_{jkt}}_{\hat{I}_{jkt} = \hat{\Pr}(I_{jkt}=1)} \delta_1 + \tilde{z}_{jkt} \delta_2 + \eta_{jkt},$$

## IV Results: Aggregate Outcomes

Outcome	Estimate	St. Error
---------	----------	-----------

---

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## IV Results: Aggregate Outcomes

Outcome	Estimate	St. Error
Total Medicare Payments	870.384**	(340.409)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## IV Results: Aggregate Outcomes

Outcome	Estimate	St. Error
Total Medicare Payments	870.384**	(340.409)
Total Hospital Costs	2,545.815***	(454.697)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## IV Results: Aggregate Outcomes

Outcome	Estimate	St. Error
Total Medicare Payments	870.384**	(340.409)
Total Hospital Costs	2,545.815***	(454.697)
Total Stays	0.271***	(0.042)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

**Does this Reflect Hospital  
Influence?**

---

# Reallocation of Patients

Outcome	Estimate	St. Error
Total Medicare Payments	75.121**	(30.902)
Total Hospital Costs	132.466***	(42.026)
Total Stays	0.015***	(0.004)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## Reallocation of Patients

Outcome	Estimate	St. Error
Total Medicare Payments	75.121**	(30.902)
Total Hospital Costs	132.466***	(42.026)
Total Stays	0.015***	(0.004)
Total Medicare Payments	63.291**	(30.853)
Total Hospital Costs	124.830***	(42.073)
Total Stays	0.014**	(0.004)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01



## Areas with most incentives...

If hospital is residual claimant on billable procedures, should see more procedures within inpatient stays

## Areas with most incentives...

If hospital is residual claimant on billable procedures, should see more procedures within inpatient stays

Outcome	Estimate	St. Error
Inpatient Costs	165.441***	(50.165)
Procedure Count	0.030***	(0.009)

\* p-value <0.1, \*\* p-value <0.05, \*\*\* p-value <0.01

## **Effects on Total Procedures and Patients**

---

# Aggregate Effects

Other ways integration posited to affect physician behavior:

- More procedures overall (largely coming from outpatient)
- Reallocating procedures (increased share to hospital)
- Changing patient profile (no evidence)

# Summary of Results

## Main Findings

- Increase in Medicare payments (\$75 to \$200) and hospital costs (\$130-\$350)
- Extrapolates to between \$55 and \$146 million in added Medicare payments from vertical integration
- Explains 4% to 10% of within-physician variation in Medicare payments

# Summary of Results

## Sensitivity

- Event study consistent with common pre-trends but limited pre-period data
- IV results suggest conservative estimates
- No improvement in quality (mortality)
- As falsification test, no effects on payments or DRG weights per inpatient stay

**Thank You!**

---