

Quantum Mechanics

Stefan Aeschbacher

July 13, 2021

Chapter 1

Dirac notation

A bra $\langle v|$ is an element in a complex vector space. The corresponding ket $|v\rangle$ is an element in its dual space. The usual rules of linear algebra are valid:

$$|u\rangle + |v\rangle = |w\rangle \quad (1.1)$$

$$c|v\rangle = |u\rangle; c \in \mathbb{C} \quad (1.2)$$

Convert between bras and kets:

$$c_1|v_1\rangle + c_2|v_2\rangle \iff c_1^*\langle v_1| + c_2^*\langle v_2| \quad (1.3)$$

1.1 Inner product

$$\langle u|v\rangle = \langle v|u\rangle^* \quad (1.4)$$

$$\langle v|v\rangle \geq 0 \quad (1.5)$$

$$\langle v|v\rangle = 0 \iff v = 0 \quad (1.6)$$

Linearity in the second argument and antilinear in the first:

$$\langle u|c_1v_1 + c_2v_2\rangle = c_1\langle u|v_1\rangle + c_2\langle u|v_2\rangle \quad (1.7)$$

$$\langle c_1u_1 + c_2u_2|v\rangle = c_1^*\langle u_1|v\rangle + c_2^*\langle u_2|v\rangle \quad (1.8)$$

$$(1.9)$$

1.2 Outer product

A bra and a ket can be combined in the outer product to create an operator

$$X = |v\rangle\langle u| \quad (1.10)$$

$$X|\Psi\rangle = |v\rangle\langle u|\Psi\rangle = |v\rangle\langle u|\Psi\rangle \quad (1.11)$$

$$(1.12)$$

Chapter 2

Postulates

2.1 Postulate 1: state

Postulate 1 *The state of a physical system is described by a state vector that belongs to a complex vector space V , called the state space of the system.*

2.2 Postulate X: time evolution

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H}(t) |\Psi(t)\rangle \quad (2.1)$$

Chapter 3

State space

$$|\Psi_1\rangle + |\Psi_2\rangle = |\Psi_3\rangle \tag{3.1}$$

$$|\Psi_1\rangle + |\Psi_2\rangle = |\Psi_2\rangle + |\Psi_1\rangle \tag{3.2}$$

Chapter 4

Operators

4.1 Basic Properties

An operator acting on a ket creates a new ket:

$$\hat{A}|\Psi\rangle = |\Psi'\rangle \quad (4.1)$$

$$|\Psi\rangle, |\Psi'\rangle \in V \quad (4.2)$$

Operators are linear

$$\hat{A}(a_1|\Psi_1\rangle + a_2|\Psi_2\rangle) = (a_1\hat{A}|\Psi_1\rangle + a_2\hat{A}|\Psi_2\rangle) \quad (4.3)$$

$$|\Psi\rangle, |\Psi'\rangle \in V; a_1, a_2 \in \mathbb{C} \quad (4.4)$$

Operators are associative and commutative under addition

$$\hat{A} + (\hat{B} + \hat{C}) = (\hat{A} + \hat{B}) + \hat{C} \quad (4.5)$$

$$\hat{A} + \hat{B} = \hat{B} + \hat{A} \quad (4.6)$$

Multiplying operators is interpreted as applying them to kets. It is associative but NOT (in general) commutative.

$$\hat{A}\hat{B}|\Psi\rangle = \hat{A}(\hat{B}|\Psi\rangle) = \hat{A}|\Psi'\rangle \quad (4.7)$$

$$\hat{A}(\hat{B}\hat{C}) = (\hat{A}\hat{B})\hat{C} \quad (4.8)$$

$$\hat{A}\hat{B} \neq \hat{B}\hat{A} \quad (4.9)$$

The lack of commutativity makes the "commutator" useful

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} \quad (4.10)$$

$$(4.11)$$

4.2 Hermitian Operators

An operator is called Hermitian (or self-adjoint) if it is its own conjugate $\hat{A} = \hat{A}^\dagger$

$$\hat{A} |A\rangle = |B\rangle \rightarrow \langle A| \hat{A}^\dagger = \langle B| \quad (4.12)$$

$$\hat{A} |A\rangle = |B\rangle \rightarrow \langle A| \hat{A} = \langle B| \quad (4.13)$$

$$(4.14)$$

A hermitian operator \hat{A} has the following properties

1. $\hat{A} |\lambda\rangle = \lambda |\lambda\rangle \rightarrow \lambda \in \mathbb{R}$
2. $\langle \hat{A} \rangle = \langle \Psi | \hat{A} | \Psi \rangle \in \mathbb{R}$
3. All eigenvectors with different eigenvalues are orthogonal

4.3 Projection Operators

A projection operator is defined

$$[\hat{A}, \hat{B}] = \hat{A} \hat{B} - \hat{B} \hat{A} \quad (4.15)$$

$$(4.16)$$

Chapter 5

Eigenvectors and Eigenvalues

5.1 Degenerate Eigenvectors

If two eigenvectors have the same eigenvalue:

$$\hat{H} |\lambda_1\rangle = \lambda |\lambda_1\rangle \quad (5.1)$$

$$\hat{H} |\lambda_2\rangle = \lambda |\lambda_2\rangle \quad (5.2)$$

their linear combination is an eigenvector as well:

$$\alpha \hat{H} |\lambda_1\rangle = \lambda \alpha |\lambda_1\rangle \quad (5.3)$$

$$\beta \hat{H} |\lambda_1\rangle = \lambda \beta |\lambda_1\rangle \quad (5.4)$$

$$\hat{H}[\alpha |\lambda_1\rangle + \beta |\lambda_2\rangle] = \lambda[\alpha |\lambda_1\rangle + \beta |\lambda_2\rangle] \quad (5.5)$$

Therefore it is possible to create two orthogonal eigenvectors for this eigenvalue.

The probability in the degenerate case is the sum of the probabilities for each eigenvector $|\langle \hat{H} | \lambda_1 \rangle|^2 + |\langle \hat{H} | \lambda_2 \rangle|^2$