How many sets were met?/ Set Logic Type	All	At least one, but not all	None
OR	Yes. The target dose can be skipped. The target dose status is 'Skipped'.	Yes. The target dose can be skipped. The target dose status is 'Skipped'.	No. The target dose cannot be skipped.

6.3 Evaluate for Inadvertent Vaccine

Evaluate for inadvertent vaccine determines if the vaccine type of a vaccine dose administered was an inadvertent administration due to the vaccine type that was administered.

The following process model, attribute table, Business Rule table and decision table, are used to evaluate for an unallowable vaccine.



FIGURE 6-4 EVALUATE FOR AN INADVERTENT VACCINE PROCESS MODEL

TABLE 6-12 INADVERTENT VACCINE ATTRIBUTES

Attribute Type	Attribute Name	Assumed Value if Empty
Vaccine dose administered	Vaccine Type	-
Supporting Data (inadvertent Vaccine)	Vaccine Type	-

TABLE 6-13 WAS THE VACCINE DOSE ADMINISTERED AN INADVERTENT ADMINISTRATION FOR THE TARGET DOSE?

CONDITIONS	RULES	
Is the vaccine type of the vaccine dose administered one of the vaccine types of an inadvertent vaccine for the target dose?	Yes	No
OUTCOMES	Yes. The vaccine dose administered was an inadvertent administration for the target dose. Target Dose Status is 'Not Satisfied'. Evaluation Status is 'Not Valid'. Evaluation Reason is 'Inadvertent Administration'.	No. The vaccine dose administered was not an inadvertent administration for the target dose.

6.4 Evaluate Age

Evaluate age validates the age at administration of a vaccine dose administered against a defined age range of a target dose. In cases where a target dose does not specify age attributes, the age at administration is considered "valid."



FIGURE 6-5 EVALUATE AGE TIMELINE

The following process model, attribute table and decision table are used to evaluate age at administration.

FIGURE 6-6 EVALUATE AGE PROCESS MODEL

TABLE 6-14 AGE ATTRIBUTES

Attribute Type	Attribute Name	Assumed Value if Empty
Patient	Date of Birth	-
Vaccine dose administered	Date Administered	-
Calculated date (CALCDTAGE-1)	Maximum Age Date	12/31/2999
Calculated date (CALCDTAGE-4)	Minimum Age Date	01/01/1900
Calculated date (CALCDTAGE-5)	Absolute Minimum Age Date	01/01/1900

TABLE 6-15 WAS THE VACCINE DOSE ADMINISTERED AT A VALID AGE?

CONDITIONS	RULES			
Is the date administered < absolute minimum age date?	Yes	No	No	No
Is the absolute minimum age date ≤ date administered < minimum age date?	No	Yes	No	No
Is the minimum age date ≤ date administered < maximum age date?	No	No	Yes	No
Is the date administered ≥ maximum age date?	No	No	No	Yes
OUTCOMES	No. The vaccine dose administered was not administered at a valid age for the target dose. Evaluation reason is 'Too young'.	Yes. The vaccine dose administered was administered at a valid age for the target dose. Evaluation reason is 'Grace period'.	Yes. The vaccine dose administered was administered at a valid age for the target dose.	No. The vaccine dose administered was not administered at a valid age for the target dose. It is extraneous. Evaluation reason is 'Too old'.

TABLE 6-16 EVALUATE AGE BUSINESS RULES

Business Rule ID	Business Rule
CALCDTAGE-1	A patient's maximum age date must be calculated as the patient's date of birth plus the maximum age.
CALCDTAGE-4	A patient's minimum age date must be calculated as the patient's date of birth plus the minimum age.
CALCDTAGE-5	A patient's absolute minimum age date must be calculated as the patient's date of birth plus the absolute minimum age.

6.5 Evaluate Preferable Interval

Evaluate preferable interval validates the date administered of a vaccine dose administered against defined preferable interval(s) from previous vaccine dose(s) administered or other events. In cases where a target dose does not specify preferable interval attributes, the interval is considered “valid.”

Preferable intervals can be measures in four different ways:

- “From Immediate Previous Dose Administered Flag” requires the interval to be evaluated from the immediate previous vaccine dose administered and is used in the majority of cases.
- “From Target Dose # in Series” requires the interval to be evaluated from the date of the specified dose.
- “From Most Recent Vaccine Type” requires the interval to be evaluated from the date of the most recently administered dose of any of the specific vaccine types listed (e.g., this is used in Pneumococcal to ensure proper spacing between the different intervals between PCV13 and PPSV23).