

Biostatistics

Applications in Genetics, Genomics, and other 'omics data

Nuno Sepúlveda, 08.01.2024

Syllabus

1. General review

- a. What is Biostatistics?
- b. Population/Sample/Sample size
- c. Type of Data – quantitative and qualitative variables
- d. Common probability distributions
- e. Work example – Malaria in Tanzania

2. Applications in Medicine

- a. Construction and analysis of diagnostic tools – Binomial distribution, sensitivity, specificity, ROC curve, Rogal-Gladen estimator
- b. Estimation of treatment effects - generalized linear models
- c. Survival analysis - Kaplan-Meier curve, log-rank test, Cox's proportional hazards model


3. Applications in Genetics, Genomics, and other 'omics data

- a. Genetic association studies – Hardy-Weinberg test, homozygosity, minor allele frequencies, additive model, multiple testing correction
- b. **Methylation association studies – M versus beta values, estimation of biological age**
- c. Gene expression studies based on RNA-seq experiments – Tests based on Poisson and Negative-Binomial

4. Other Topics

- a. Estimation of Species diversity – Diversity indexes, Poisson mixture models
- b. Serological analysis – Gaussian (skew-normal) mixture models
- c. Advanced sample size and power calculations

Genotypic-phenotype mapping



Single nucleotide
polymorphisms
Copy number variation
Inversion/deletion

Post-translation modification
DNA methylation
microRNA

Different
symptoms for the
same disease

Genetic association studies

Genetic
variation



Disease

Single nucleotide
polymorphisms

Copy number variation

Inversion/deletion


A C T T A **C** G C C T T A A T C C G

T G A A T **G** C C G A A T T A G G C

A C T T A **A** G C C T T A A T C C G

T G A A T **T** C C G A A T T A G G C

Genotypic-phenotype mapping



Single nucleotide
polymorphisms
Copy number variation
Inversion/deletion

Post-translation modification
DNA methylation
microRNA


Different
symptoms for the
same disease

DNA methylation



Gene might be expressed

Production of the protein



Gene might not be expressed

No production of the protein

Epigenome-wide association studies


Aim:

Search the whole genome for methylation modifications associated with the phenotype (i.e., presence of disease)

Example: Case-control study



n=48
75% Women
Mean age of 37 years old
Mean BMI of 27 kg/m²




n=61
79% Women
Mean age of 41 years old
Mean BMI of 27 kg/m²


Human Methylation 450K Array

DNA methylation array

Array




Green
light




Red
light

$$\frac{\text{Intensity of red light}}{(\text{Intensity of red light} + \text{Intensity of green light})}$$

Example of data for a single probe




Simple statistical analysis of a single probe



Which statistical tests can we use to check whether patients differ from healthy controls in terms of the methylation level for this probe?

Simple statistical analysis of a single probe



T test using original or transformed data

Mann-Whitney test

Exercise: Data_lecture_12_ACE_ACE2.csv

Heliyon 7 (2021) e07665



Contents lists available at [ScienceDirect](#)

Heliyon

journal homepage: www.cell.com/heliyon



Research article

The SARS-CoV-2 receptor angiotensin-converting enzyme 2 (ACE2) in myalgic encephalomyelitis/chronic fatigue syndrome: A meta-analysis of public DNA methylation and gene expression data



João Malato^{a,b}, Franziska Sotzny^{c,**}, Sandra Bauer^c, Helma Freitag^c, André Fonseca^d, Anna D. Grabowska^e, Luís Graça^a, Clara Cordeiro^{b,d}, Luís Nacul^{f,g}, Eliana M. Lacerda^f, Jesus Castro-Marrero^h, Carmen Scheibenbogen^c, Francisco Westermeier^{i,j}, Nuno Sepúlveda^{b,c,k,*}

^a Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal

^b CEAUL – Centro de Estatística e Aplicações, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal

^c Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt Universität zu Berlin and Berlin Institute of Health, Institute of Medical Immunology, Berlin, Germany

^d Faculdade de Ciências e Tecnologia, Universidade do Algarve, Faro, Portugal

^e Department of Biophysics, Physiology, and Pathophysiology, Medical University of Warsaw, Warsaw, Poland

^f Department of Clinical Research, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom

^g Complex Chronic Diseases Program, British Columbia Women's Hospital and Health Centre, Vancouver, British Columbia, Canada

^h Vall d'Hebron Hospital Research Institute, Division of Rheumatology, ME/CFS Unit, Universitat Autònoma de Barcelona, Barcelona, Spain

ⁱ Institute of Biomedical Science, Department of Health Studies, FH Joanneum University of Applied Sciences, Graz, Austria

^j Centro Integrativo de Biología y Química Aplicada (CIBQA), Universidad Bernardo O'Higgins, Santiago, Chile

^k Department of Infection Biology, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, United Kingdom

Exercise: Data_lecture_12_ACE_ACE2.csv

Information about the Probes: ACE_ACE2_merged_annotation_heliyon.csv

Compare the methylation levels of 27 CpG probes located in ACE and ACE2 genes between patients with chronic fatigue syndrome and healthy controls using T test and Mann-Whitney test.


Which test is preferable to analyse data of each probe?

Which probes are differentially methylated when comparing patients and healthy controls after adjusting for multiple testing?

Are these probes hypo or hypermethylated in patients?

Statistical analysis of a single probe adjusting for covariates

Linear regression (as in GWAS for quantitative traits)



Y_{ij} = methylation levels of probe j in individual i

$x_{group,i}$ = group of individual i

$x_{k,i}$ = value of covariate k for the individual i

$$Y_{ij} = \beta_{0j} + \beta_{1j}x_{group,i} + \sum_{k=2}^p \beta_{kj}x_{k,i} + \underbrace{\epsilon_{ij}}_{\text{residuals}}$$

$$\epsilon_{ij} \rightsquigarrow N(\mu = 0; \sigma = \sigma_0)$$

$$H_0 : \beta_{1j} = 0 \text{ versus } H_1 : \beta_{1j} \neq 0$$

Epigenome-wide association studies

Perform this test on data of each probe

$$H_0 : \beta_{1j} = 0 \text{ versus } H_1 : \beta_{1j} \neq 0$$

$$j = 1, \dots, M \text{ (number of probes)}$$


Wald's score test

Wilks' likelihood ratio test


Correct the p-values of each individual test by a procedure controlling the false discovery rate (e.g., Benjamini-Hochberg procedure)

Construct a manhattan plot as learned for the genome-wide association studies


Report the significant probes and their location




Manhattan plots adjusting the significance level via Bonferroni correction for multiple testing



Manhattan plots adjusting p-values via Benjamini-Hochberg procedure




Decreased methylation in patients with ME/CFS



More expression for genes associated with these probes

Discussion



Is it reasonable to use linear regression in this kind of data?

What are the eventual problems?

Exercise: Data_lecture_12_ACE_ACE2.csv

Repeat previous analysis using linear regression adjusting the effect of study and gender. Perform a residual analysis to validate the model for each probe.

Analysing M values instead

Use of linear regression again under an appropriate transformation of the outcome

Y_{ij} = methylation levels of probe j in individual i

$$\underbrace{Y_{ij}}_{\text{Beta values}} \rightarrow \underbrace{Y_{ij}^*}_{\text{M values}} = \log \frac{Y_{ij}}{1 - Y_{ij}} \text{ or } \log_2 \frac{Y_{ij}}{1 - Y_{ij}}$$

$$Y_{ij}^* = \beta_{0j} + \beta_{1j}x_{group,i} + \sum_{k=2}^p \beta_{kj}x_{k,i} + \epsilon_{ij}$$

$$\epsilon_{ij} \rightsquigarrow N(\mu = 0; \sigma = \sigma_0)$$

What are the theoretical advantages of this approach?

Alternative statistical analysis of a single probe adjusting for covariates

Du et al. *BMC Bioinformatics* 2010, **11**:587
<http://www.biomedcentral.com/1471-2105/11/587>



RESEARCH ARTICLE

Open Access

Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis

Pan Du^{1,3*}, Xiao Zhang², Chiang-Ching Huang², Nadereh Jafari⁴, Warren A Kibbe^{1,3}, Lifang Hou^{2,3}, Simon M Lin^{1,3*}

Abstract

Background: High-throughput profiling of DNA methylation status of CpG islands is crucial to understand the epigenetic regulation of genes. The microarray-based Infinium methylation assay by Illumina is one platform for low-cost high-throughput methylation profiling. Both Beta-value and M-value statistics have been used as metrics to measure methylation levels. However, there are no detailed studies of their relations and their strengths and limitations.

Results: We demonstrate that the relationship between the Beta-value and M-value methods is a Logit transformation, and show that the Beta-value method has severe heteroscedasticity for highly methylated or unmethylated CpG sites. In order to evaluate the performance of the Beta-value and M-value methods for identifying differentially methylated CpG sites, we designed a methylation titration experiment. The evaluation results show that the M-value method provides much better performance in terms of Detection Rate (DR) and True Positive Rate (TPR) for both highly methylated and unmethylated CpG sites. Imposing a minimum threshold of difference can improve the performance of the M-value method but not the Beta-value method. We also provide guidance for how to select the threshold of methylation differences.

Conclusions: The Beta-value has a more intuitive biological interpretation, but the M-value is more statistically valid for the differential analysis of methylation levels. Therefore, we recommend using the M-value method for conducting differential methylation analysis and including the Beta-value statistics when reporting the results to investigators.




Figure 1 The relationship curve between M-value and Beta-value.

M values can be positive and negative

The range of M values is wider especially at the extremes of the Beta value scale

Alternative statistical analysis of a single probe adjusting for covariates

Du et al. *BMC Bioinformatics* 2010, **11**:587
<http://www.biomedcentral.com/1471-2105/11/587>



RESEARCH ARTICLE

Open Access

Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis

Pan Du^{1,3*}, Xiao Zhang², Chiang-Ching Huang², Nadereh Jafari⁴, Warren A Kibbe^{1,3}, Lifang Hou^{2,3}, Simon M Lin^{1,3*}

Abstract

Background: High-throughput profiling of DNA methylation status of CpG islands is crucial to understand the epigenetic regulation of genes. The microarray-based Infinium methylation assay by Illumina is one platform for low-cost high-throughput methylation profiling. Both Beta-value and M-value statistics have been used as metrics to measure methylation levels. However, there are no detailed studies of their relations and their strengths and limitations.

Results: We demonstrate that the relationship between the Beta-value and M-value methods is a Logit transformation, and show that the Beta-value method has severe heteroscedasticity for highly methylated or unmethylated CpG sites. In order to evaluate the performance of the Beta-value and M-value methods for identifying differentially methylated CpG sites, we designed a methylation titration experiment. The evaluation results show that the M-value method provides much better performance in terms of Detection Rate (DR) and True Positive Rate (TPR) for both highly methylated and unmethylated CpG sites. Imposing a minimum threshold of difference can improve the performance of the M-value method but not the Beta-value method. We also provide guidance for how to select the threshold of methylation differences.

Conclusions: The Beta-value has a more intuitive biological interpretation, but the M-value is more statistically valid for the differential analysis of methylation levels. Therefore, we recommend using the M-value method for conducting differential methylation analysis and including the Beta-value statistics when reporting the results to investigators.




Figure 4 Performance comparisons of Beta- and M-value in the range of low, middle and high methylation levels based on the relationship of 1 - Detection Rate versus True Positive Rate.

Exercise: Data_lecture_12_ACE_ACE2.csv

Repeat previous analysis using linear regression adjusting the effect of study and gender. Perform a residual analysis to validate the model for each probe.

Analysing Beta values with beta regression

Y_{ij} = methylation levels of probe j in individual i

$$Y_{ij} \in (0,1)$$

$$Y_{ij} \rightsquigarrow \text{Beta}(\alpha_{ij}, \beta_{ij})$$

$$\mu_{ij} = \frac{\alpha_{ij}}{\alpha_{ij} + \beta_{ij}}$$

$$\phi_{ij} = \alpha_{ij} + \beta_{ij}$$


Useful
reparametrization

$$Y_{ij} \rightsquigarrow \text{Beta}(\mu_{ij}, \phi_{ij})$$

Beta regression

$$Y_{ij} \rightsquigarrow \text{Beta}(\mu_{ij}, \phi_{ij}) \rightarrow Y_{ij} \rightsquigarrow \text{Beta}(\mu_{ij}, \phi_j)$$

$$\mu_{ij} = \beta_{0j} + \beta_{1j}x_{\text{group},i} + \sum_{k=2}^p \beta_{kj}x_{k,i}$$



Beta distribution is very flexible

$$f_{X|\alpha,\beta}(x) = \frac{x^{\alpha-1}(1-x)^{\beta-1}}{Be(\alpha,\beta)} I_{(0,1)}(x)$$


Exercise: Data_lecture_12_ACE_ACE2.csv

Repeat previous analysis using Beta regression adjusting the effect of study and gender.
Install and use the package “betareg”.

Compare the results from all the analyses done so far.

What are your conclusions?

DNA methylation and age prediction



DNA methylation age of human tissues
and cell types

Horvath

Infinium Illumina 27k - 27,578 CpG probes

Infinium Illumina 450k - 485,512 CpG probes

Training data sets

n=7844 samples from 82 data sets

Samples from the different tissues (PBMC, lung, kidney, etc)

% methylation in more than 20k CpG

Final linear regression model (DNAm)

Outcome = Age

Covariates = % methylation of 353 “clock” CpG probes


Penalized regression using elastic net.

DNA methylation and age prediction



[https://www.ted.com/talks/
steve_horvath_epigenetic_clocks_help_to_find_anti_aging_treatments](https://www.ted.com/talks/steve_horvath_epigenetic_clocks_help_to_find_anti_aging_treatments)


Prediction in the training set



Can we know whether we are aging faster than we should?

Age acceleration =

DNA methylation age - Chronological age



Age acceleration < 0

Age acceleration > 0


Younger than your chronological age

Older than your chronological age



What about the aging of patients with ME/CFS?

Reference (tissue)	ME/CFS patients			Healthy controls, n	Technology (manufacturer)	GEO ID
	n	Sample characteristics	Case definition			
De Vega et al. (2014) (PBMC)	12	Female adults Mean age: 41 yo Mean BMI: 23 kg/m ²	1994 CDC/Fukuda & 2003 CCC	12	Infinium HumanMethylation450K	GSE59489
De Vega et al. (2017) (PBMC)	49	Female adults Mean age: 50 Mean BMI: 23 kg/m ²	1994 CDC/Fukuda & 2003 CCC	25	Infinium HumanMethylation450 Array (Illumina)	GSE93266
Herrera et al. (2018) (T Lymphocytes)	61	Female/male adults Mean age: 32 yo Mean BMI: 27 kg/m²	1994 CDC/Fukuda & 2003 CCC	48	Infinium HumanMethylation450 K Array (Illumina)	GSE156792
Trivedi et al. (2018) (PBMC)	13	Female adults Mean age: 50 yo Mean BMI: 26 kg/m²	1994 CDC/Fukuda & 2003 CCC	12	Methylation EPIC Array (Illumina)	GSE111183

Herrera et al data




Trivedi et al data




De Vega et al studies

De Vega et al (2014) data



De Vega et al (2017) data



How to construct your own model for predicting the biological age based on DNA methylation data?