



Study samples, sampling timepoints and analysis populations:
maternal-cord-infant systems-serology substudy of the Gambian Pertussis Study (GaPs)



Cord blood and *InfMon2 (V2)* analysis populations, by infant group

Tdap-IPV / aP	Tdap-IPV / wP	TT / aP	TT / wP
randomised n = 73	randomised n = 78	randomised n = 83	randomised n = 78
InfMon2 endpoint, n analysed	InfMon2 endpoint, n analysed	InfMon2 endpoint, n analysed	InfMon2 endpoint, n analysed
Whole-cell B. pertussis IgG51	Whole-cell B. pertussis IgG63	Whole-cell B. pertussis IgG58	Whole-cell B. pertussis IgG51
PT-neutralisation53	PT-neutralisation63	PT-neutralisation64	PT-neutralisation57
Serum bactericidal activity51	Serum bactericidal activity54	Serum bactericidal activity62	Serum bactericidal activity55

InfMon2 totals — antigen-specific total-IgG panel: 287 · whole-cell IgG (*WT_IgG*): 223 · PT-neutralisation: 237 · SBA: 222 (238 assayed).
Sample size varies by assay and model; complete measurements across the full panel were not required.

Reconciliation of the analysis population: why n = 287, not n = 239

- The *InfMon2* sample is drawn at ~8 weeks of age, immediately before the first pertussis-containing dose. It therefore predominantly reflects maternally derived passive immunity, so infants assigned to future DTaP (aP) and DTwP (wP) schedules are pooled for the pre-primary analyses.
- Because no infant pertussis vaccine has yet been given, infants need not have completed the per-protocol vaccination series to contribute the *InfMon2* baseline. The previously published per-protocol immunogenicity population (n = 239) was, by contrast, restricted to infants who completed the protocol-defined primary series — a stricter and smaller set.
- The systems-serology *InfMon2* population is therefore larger (n = 287 with the antigen-specific total-IgG panel) than the per-protocol population. The analysis population is defined per analysis as randomised mother-infant pairs with the infant outcome and the antibody measurements required for that analysis; longitudinal maternal-cord-infant analyses are restricted to pairs with the relevant paired samples.

aP = acellular-pertussis (DTaP) primed; wP = whole-cell-pertussis (DTwP) primed. Visit recoding: P02→*PregEarly*, P09→*MatBirth*, M00→*CordBlood*, M02→*InfMon2 (V2)*.