

A WEAKER NOTION OF THE FINITE FACTORIZATION PROPERTY

HENRY JIANG, SHIHAN KANUNGO, AND HWISOO KIM

ABSTRACT. An (additive) commutative monoid is called atomic if every given non-invertible element can be written as a sum of atoms (i.e., irreducible elements), in which case, such a sum is called a factorization of the given element. The number of atoms (counting repetitions) in the corresponding sum is called the length of the factorization. Following Geroldinger and Zhong, we say that an atomic monoid M is a length-finite factorization monoid if each $b \in M$ has only finitely many factorizations of any prescribed length. An additive submonoid of $\mathbb{R}_{\geq 0}$ is called a positive monoid. Factorizations in positive monoids have been actively studied in recent years. The main purpose of this paper is to give a better understanding of the non-unique factorization phenomenon in positive monoids through the lens of the length-finite factorization property. To do so, we identify a large class of positive monoids which satisfy the length-finite factorization property. Then we compare the length-finite factorization property to the bounded and the finite factorization properties, which are two properties that have been systematically investigated for more than thirty years.

1. Introduction

Following Cohn [14], we say that a commutative monoid is atomic provided that every non-invertible element can be written as a product of atoms (also called irreducibles), while an integral domain is called atomic provided that its multiplicative monoid is atomic. Following Anderson, Anderson, and Zafrullah [2], we say that a monoid/domain satisfy the finite factorization property if every nonzero nonunit element has only finitely many factorizations (into atoms). A weaker version of the finite factorization property, also introduced in [2], is the bounded factorization property: for each nonzero nonunit element there exists a bound for the number of atoms (counting repetitions) in any factorization of such an element. The bounded and the finite factorization properties have received a great deal of attention since they were introduced back in 1990 (see the recent paper [3] by Anderson and Gotti for a survey on bounded and finite factorization domains).

2020 *Mathematics Subject Classification.* Primary: 11Y05, 20M13; Secondary: 06F05, 20M14.

Key words and phrases. length-finite factorization, positive monoid, positive semiring, finite factorization, bounded factorization, length-factoriality.

A commutative monoid is called factorial if every nonunit element has a unique factorization. Factorial monoids are, therefore, atomic. An atomic monoid is called half-factorial if any two factorizations of the same element have the same number of atoms (counting repetitions). The half-factorial property was coined by Zaks [34] in the context of integral domains, and the same property has been systematically studied since then: see the recent paper [18] by Gao et al. (and references therein) and see the paper [10] by Chapman and Coykendall for a survey comprising the most relevant studies of half-factoriality until 2000). The following complementary notion of half-factoriality has been considered more recently: a monoid is called length-factorial if any two distinct factorizations of the same element have distinct lengths (it is clear that a monoid is factorial if and only if it is both half-factorial and length-factorial). The notion of length-factoriality was introduced by Coykendall and Smith in [17], and it has been recently investigated by several authors (see, for instance, [7, 11, 20]).

We say that the length-factorial property complements the half-factorial property because the former is precisely what a half-factorial monoid needs to be factorial. Similarly, we can define a property that complements the bounded factorization property with respect to the finite factorization property. An atomic monoid is said to satisfy the length-finite factorization property if each element has only finitely many factorizations with any prescribed number of atoms (counting repetitions). Observe that a monoid satisfies the finite factorization property if and only if it satisfies both the bounded and the length-finite factorization properties. The length-finite factorization property was recently introduced by Geroldinger and Zhong in [20]. The main purpose of this paper is to offer a better understanding of factorizations in positive monoids and positive semirings using the light shed by the length-finite factorization property. While doing so, we provide the first dedicated study of the length-finite factorization property.

A positive monoid is an additive submonoid of \mathbb{R} consisting of nonnegative real numbers. On the other hand, a positive semiring is a positive monoid that contains 1 and is closed under multiplication. The atomic structure and the arithmetic of factorizations of both positive monoids and positive semirings have been actively studied in recent years. For instance, atomicity and factorizations in positive monoids have been studied by Chapman et al. in [13, 15, 22] and more recently by Gotti and Vulakh in [29]. On the other hand, atomicity and factorizations in positive semirings have been studied by Baeth et al. in [4, 5, 35] and more recently by Gotti and Polo in [27, 28]. Factorizations in positive semirings consisting of rationals have also been considered in recent literature by various authors (see, for instance, [12, 32]).

In Section 2, we briefly discuss the notation, terminology, and main known results we will use throughout the rest of the paper.

Section 3, which is the first section of content, is devoted to study the length-finite factorization property in the setting of positive monoids. The main result we establish in Section 3 is Theorem 3.4, where we identify a large class of positive monoids satisfying the length-finite factorization property. In the second part of the same section, we discuss a few examples connecting the length-finite factorization property with some well-studied atomic properties. It is known that every commutative monoid satisfying the ascending chain on principal ideals (ACCP) is atomic [19, Proposition 1.1.4], and it is also known that every monoid that satisfies the bounded factorization property also satisfies the ACCP [31, Corollary 1]. We provide in Section 3 an example of a positive monoid that satisfies the length-finite factorization property but not the ACCP and an example of a positive monoid that satisfies the bounded factorization property but not the length-finite factorization property.

In Section 4, we consider the length-finite factorization property in the setting of positive semirings. Although the main purpose of Section 4 is to discuss examples of positive semirings in connection with the length-finite factorization property, we also discuss a method to construct positive semirings out of positive monoids using certain exponentiation construction. In this direction, we prove that the ACCP, the bounded factorization property, and the finite factorization property are all preserved under the mentioned exponentiation construction. Additionally, we show exponentiation construction is in fact isomorphic to the monoid semiring over the natural numbers. It is still unknown to the authors whether the length-finite factorization property is preserved under the same exponentiation construction, and we pose this as Question 4.4.

2. Background

2.1. General Notation

Following standard notation, we let \mathbb{Z} , \mathbb{Q} , and \mathbb{R} denote the set of integers, rational numbers, and real numbers, respectively. Also, we let \mathbb{N} , \mathbb{N}_0 , and \mathbb{P} denote the set of positive integers, nonnegative integers, and primes, respectively. For $b, c \in \mathbb{Z}$ with $b \leq c$, we let $\llbracket b, c \rrbracket$ denote the set of integers between b and c , i.e.,

$$\llbracket b, c \rrbracket = \{n \in \mathbb{Z} \mid b \leq n \leq c\}.$$

For $S \subseteq \mathbb{R}$ and $r \in \mathbb{R}$, we set $S_{\geq r} = \{s \in S \mid s \geq r\}$ and $S_{>r} = \{s \in S \mid s > r\}$.

2.2. Commutative Monoids

Throughout this paper, we reserve the term *monoid* for a cancellative and commutative semigroup with an identity element. Monoids here are written additively unless we specify otherwise. Let M be a monoid. We let M^\bullet denote the set of nonzero elements of M , while we let $\mathcal{U}(M)$ denote the group of invertible elements of M . The set $M/\mathcal{U}(M) = \{b + \mathcal{U}(M) \mid b \in M\}$ is a monoid under the natural operation induced by the operation of M . We say

that M is *reduced* provided that $\mathcal{U}(M)$ is the trivial group, in which case we can canonically identify $M/\mathcal{U}(M)$ with M . A *submonoid* of M is a subset of M that is closed under addition and contains 0. The monoids we are mostly concerned with in the scope of this paper are positive monoids, which are reduced. A *positive monoid* is an additive submonoid of $\mathbb{R}_{\geq 0}$. If S is a subset of M , then $\langle S \rangle$ stands for the smallest (under inclusion) submonoid of M that contains S . If $M = \langle S \rangle$ for some finite subset S , then we say that M is *finitely generated*. Additive submonoids of \mathbb{N}_0 , called *numerical monoids*, are always finitely generated. A map $\varphi: M \rightarrow M'$, where M' is a monoid, is called a (*monoid*) *homomorphism* if $\varphi(b + c) = \varphi(b) + \varphi(c)$ for all $b, c \in M$.

An element $a \in M \setminus \mathcal{U}(M)$ is called an *atom* if for all $b, c \in M$ the fact that $a = b + c$ guarantees that $b \in \mathcal{U}(M)$ or $c \in \mathcal{U}(M)$. We let $\mathcal{A}(M)$ denote the set of all the atoms of M . Following [16], we say that M is *antimatter* if $\mathcal{A}(M)$ is empty. An element $b \in M$ is called *atomic* if either $b \in \mathcal{U}(M)$ or if b can be written as a sum of atoms (repetitions are allowed). Following [14], we say that M is *atomic* if every element of M is atomic. Atomic monoids play an important role in this paper. A subset I of M is called an *ideal* if $I + M \subseteq I$. An ideal I of M is called *principal* if $I = b + M$ for some $b \in M$. A sequence of ideals $(I_n)_{n \geq 1}$ of M is called an *ascending chain* if $I_n \subseteq I_{n+1}$ for every $n \in \mathbb{N}$. The monoid M is said to satisfy the *ascending chain condition on principal ideals* (ACCP) if every ascending chain of principal ideals $(I_n)_{n \geq 1}$ stabilizes, which means that there exists $N \in \mathbb{N}$ such that $I_n = I_N$ for every $n \geq N$. If a monoid satisfies the ACCP, then it is atomic [19, Proposition 1.1.4]. The converse does not hold (see, for instance, Example 3.6). In addition, finitely generated monoids satisfy the ACCP (see, for instance, [19, Proposition 2.7.8]), and so they are atomic.

2.3. Factorizations and Lengths

Let $Z(M)$ denote the monoid consisting of all formal sums of atoms in $\mathcal{A}(M/\mathcal{U}(M))$; that is, $Z(M)$ is the free commutative monoid on $\mathcal{A}(M/\mathcal{U}(M))$. The elements of $Z(M)$ are called *factorizations* in M . Let the function $\pi: Z(M) \rightarrow M/\mathcal{U}(M)$ be the only monoid homomorphism such that $\pi(a) = a$ for all $a \in \mathcal{A}(M/\mathcal{U}(M))$. Then for each $b \in M$, we set

$$Z(b) := \{z \in Z(M) \mid \pi(z) = b + \mathcal{U}(M)\},$$

and call the element of $Z(b)$ *factorizations of b* . Observe that M is atomic if and only if $Z(b)$ is nonempty for every $b \in M$. The monoid M is called a *unique factorization monoid* (UFM) if $|Z(b)| = 1$ for every $b \in M$, while M is called a *finite factorization monoid* (FFM) if $1 \leq |Z(b)| < \infty$ for every $b \in M$. It follows from the definitions that every UFM is an FFM. Also, it is well known that every finitely generated monoid is an FFM [19, Proposition 2.7.8].

If $z := a_1 \cdots a_\ell \in Z(M)$ for some $a_1, \dots, a_\ell \in \mathcal{A}(M/\mathcal{U}(M))$, then we call ℓ the *length* of z , and we often denote ℓ by $|z|$. For $b \in M$, the set

$$L(b) := \{|z| \mid z \in Z(b)\}$$

is called the *set of lengths* of b . We say that M is a *bounded factorization monoid* (BFM) if $1 \leq |L(b)| < \infty$ for every $b \in M$. It follows from the definitions that every FFM is a BFM. Also, it is well known that every BFM satisfies the ACCP [31, Corollary 1]. For each $b \in M$ and $\ell \in \mathbb{N}$, set

$$Z_\ell(b) := \{z \in Z(b) \mid |z| = \ell\}.$$

Following [20], we say that M is a *length-finite factorization monoid* (LFFM) if M is atomic and $|Z_\ell(b)| < \infty$ for all $b \in M$ and $\ell \in \mathbb{N}$. From the definitions, we obtain that every FFM is an LFFM. As we shall see in the next section, the notions of a BFM and an LFFM are not comparable.

2.4. Commutative Semirings

A nonempty set S endowed with two binary operations ‘+’ and ‘·’ is called a *commutative semiring*, or simply a *semiring*¹ in the scope of this paper, if the following three conditions are satisfied:

- (1) $(S, +)$ is a monoid with its identity element denoted by 0;
- (2) (S^\bullet, \cdot) is a monoid with identity element denoted by 1;
- (3) $b \cdot (c + d) = b \cdot c + b \cdot d$ for all $b, c, d \in S$.

The operations ‘+’ and ‘·’ are called *addition* and *multiplication*, respectively. For any $b, c \in S$, we often write bc rather than $b \cdot c$. Let S be a semiring. We refer to the invertible elements of the multiplicative monoid S^\bullet simply as *units* of S , and we denote the set of units of S by S^\times . A subset S' of a semiring S is called a *subsemiring* of S if $(S', +)$ is a submonoid of $(S, +)$ that contains 1 and is closed under multiplication. In this paper, we restrict our attention to positive semirings. A *positive semiring* is a subsemiring of $\mathbb{R}_{\geq 0}$ under the standard addition and multiplication. Observe that the additive monoid of a positive semiring is a positive monoid.

Let S be a positive semiring. Since the additive monoids of most of the positive semirings we consider in this paper are UFM, when we mention a monoidal divisibility or atomic property of a semiring, we will be referring to its multiplicative structure. Accordingly, we say that S is *antimatter* (resp., *atomic*, an *LFFS*, a *BFS*, an *FFS*) provided that the multiplicative monoid S^\bullet is antimatter (resp., atomic, an LFFM, a BFM, an FFM). Similarly, we say that S satisfies the *ACCP* if its multiplicative monoid S^\bullet satisfies the ACCP.

¹Although in the scope of this paper we are only interested in semirings that are cancellative and commutative with respect to both operations, it is worth emphasizing that a more general and standard definition of a semiring does not assume any of these conditions (see [21]).

3. Length-Finite Factorization Positive Monoids

Following [25], we say that a positive monoid is *decreasing* (resp., *increasing*) if it can be generated by a decreasing (resp., increasing) sequence, and we say that a positive monoid is *monotone* if it is either decreasing or increasing. Monotone positive monoids were also studied in [8, 9, 22, 33]. It is clear that every numerical monoid is both decreasing and increasing. In general, a positive monoid is both decreasing and increasing if and only if it is finitely generated [22, Proposition 5.4].

We say that a sequence of real numbers is *well-ordered* (resp., *co-well-ordered*) if it contains no strictly decreasing (resp., increasing) subsequence. Following Polo [33], we say that a positive monoid M is *well-ordered* (resp., *co-well-ordered*) if it can be generated by a well-ordered (resp., co-well-ordered) sequence. It follows from the definitions that every increasing (resp., decreasing) positive monoid is *well-ordered* (resp., *co-well-ordered*). As the following example illustrates, there are atomic positive monoids that are neither well-ordered nor co-well-ordered.

Example 3.1. Let $(p_n)_{n \geq 1}$ be a strictly increasing sequence of primes, and consider the positive monoid

$$M := \left\langle 1 + \frac{(-1)^n}{p_n} \mid n \in \mathbb{N} \right\rangle.$$

Set $a_n := 1 + \frac{(-1)^n}{p_n}$ for every $n \in \mathbb{N}$. Since for each $n \in \mathbb{N}$ there is only one defining generator, namely a_n , whose p_n -adic valuation is negative, the same defining generator must be an atom. As a consequence,

$$\mathcal{A}(M) = \{a_n \mid n \in \mathbb{N}\},$$

which implies that M is atomic. Since M is reduced, any generating set of M must contain $\mathcal{A}(M)$. Therefore the fact that $(a_{2n})_{n \geq 1}$ is a strictly decreasing sequence implies that M cannot contain any well-ordered generating sequence, while the fact that $(a_{2n-1})_{n \geq 1}$ is strictly increasing implies that M cannot contain any co-well-ordered generating sequence. Hence M is neither a well-ordered nor a co-well-ordered positive monoid.

3.1. A Class of Length-Finite Factorization Monoids

In this subsection, we identify a large class consisting of positive monoids that are LFFM: indeed, as our main result, we prove that every co-well-ordered positive monoid is an LFFM. In order to prove this, we need the following two lemmas.

Lemma 3.2. *The sum of finitely many co-well-ordered sequences is a co-well-ordered sequence.*

Proof. It suffices to argue that the sum of two co-well-ordered sequences is again a co-well-ordered sequence as, after proving this we can induct on the

number of sequences being added to obtain the statement of the lemma. Let $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ be two co-well-ordered sequences. Now suppose, for the sake of a contradiction, that the sequence $(a_n + b_n)_{n \geq 1}$ is not co-well-ordered. Let $(c'_n)_{n \geq 1}$ be a strictly increasing subsequence of $(a_n + b_n)_{n \geq 1}$. Take subsequences $(a'_n)_{n \geq 1}$ and $(b'_n)_{n \geq 1}$ of $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$, respectively, such that $(c'_n)_{n \geq 1} = (a'_n)_{n \geq 1} + (b'_n)_{n \geq 1}$. In light of the Monotone Subsequence Theorem, after passing to suitable subsequences, we can assume that $(a'_n)_{n \geq 1}$ and $(b'_n)_{n \geq 1}$ are both monotone sequences. Since $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ are both co-well-ordered, neither $(a'_n)_{n \geq 1}$ nor $(b'_n)_{n \geq 1}$ can contain a strictly increasing subsequence. Hence there exists $k \in \mathbb{N}$ such that the sequences $(a'_n)_{n \geq k}$ and $(b'_n)_{n \geq k}$ are decreasing. This implies that $(c'_n)_{n \geq k}$ is also a decreasing sequence, which contradicts that $(c'_n)_{n \geq 1}$ is a strictly increasing sequence and, therefore, contains no decreasing subsequences. Hence $(a_n + b_n)_{n \geq 1}$ is a co-well-ordered sequence. \square

A pair (z_1, z_2) of factorizations in $Z(M)$ of the same element of M is called *irredundant* if z_1 and z_2 do not share any atoms. In addition, we say that a subset Z of $Z(M)$ is *irredundant* if any two distinct factorizations in Z are irredundant.

Lemma 3.3. *Let M be an atomic co-well-ordered positive monoid. For all $x \in M$ and $\ell \in \mathbb{N}$, every irredundant subset of $Z_\ell(x)$ is finite.*

Proof. Fix $x \in M$ and $\ell \in \mathbb{N}$. For the sake of a contradiction, suppose that there exists an infinite irredundant subset of $Z_\ell(x)$. Therefore there exists a sequence of factorizations $(z_n)_{n \geq 1}$ whose terms are pairwise distinct and whose underlying set is an irredundant subset of $Z_\ell(x)$. For a factorization $z := a_1 \cdots a_\ell \in Z(M)$ with $a_1, \dots, a_\ell \in \mathcal{A}(M)$ such that $a_1 \leq \cdots \leq a_\ell$ and for each $k \in \llbracket 1, \ell \rrbracket$, we call a_k the k -th atom of z . For each $n \in \mathbb{N}$ and $k \in \llbracket 1, \ell \rrbracket$, let $a_n^{(k)}$ denote the k -th atom of z_n . By virtue of the Monotone Subsequence Theorem, the sequence $(a_n^{(1)})_{n \geq 1}$ has a monotone subsequence. As a result, after replacing $(z_n)_{n \geq 1}$ by a suitable subsequence, we can assume that $(a_n^{(1)})_{n \geq 1}$ is monotone. Since the underlying set of $(z_n)_{n \geq 1}$ is irredundant, the sequence $(a_n^{(1)})_{n \geq 1}$ must be either strictly decreasing or strictly increasing. The fact that M is co-well-ordered, however, ensures that no sequence of atoms of M can be strictly increasing, and so we can assume that $(a_n^{(1)})_{n \geq 1}$ is strictly decreasing. This, along with the fact that $\sum_{i=1}^{\ell} a_n^{(i)} = x$ for every $n \in \mathbb{N}$, ensures that the sequence $(s_n)_{n \geq 1} := (a_n^{(2)})_{n \geq 1} + \cdots + (a_n^{(\ell)})_{n \geq 1}$ is strictly increasing. Therefore $(s_n)_{n \geq 1}$ is not co-well-ordered, and so it follows from Lemma 3.2 that $(a_n^{(j)})_{n \geq 1}$ is not co-well-ordered for some $j \in \llbracket 2, \ell \rrbracket$. Thus, there exists a strictly increasing sequence whose terms are atoms of M , which contradicts the fact that M is co-well-ordered. Hence for all $x \in M$ and $\ell \in \mathbb{N}$, every irredundant subset of $Z_\ell(x)$ must be finite. \square

We are in a position to prove the main result of this section.

Theorem 3.4. *Every atomic co-well-ordered positive monoid is an LFFM.*

Proof. Let M be an atomic co-well-ordered positive monoid. Now suppose, by way of contradiction, that M is not an LFFM. Then for some $\ell \in \mathbb{N}$ there exists $x \in M$ such that $|Z_\ell(x)| = \infty$. Furthermore, assume that we have chosen ℓ as the smallest as it can possibly be. Since $|Z_1(x)| \leq 1$, we see that $\ell \geq 2$. Our next step is to argue the existence of a maximal irredundant subset of $Z_\ell(x)$ under inclusion.

Let \mathcal{Z} be the poset consisting of all irredundant subsets of $Z_\ell(x)$ under inclusion. As $Z_\ell(x)$ is infinite, and so nonempty, \mathcal{Z} is a nonempty poset. In addition, it follows from Lemma 3.3 that every element in \mathcal{Z} is a finite set. Now suppose that $\mathcal{C} := \{Z_\gamma \mid \gamma \in \Gamma\}$ is a nonempty chain in the poset \mathcal{Z} . We claim that $U := \bigcup_{\gamma \in \Gamma} Z_\gamma$ is an upper bound for \mathcal{C} in \mathcal{Z} . If z and z' are two distinct factorizations in U , then as \mathcal{C} is a chain, we can pick $\gamma \in \Gamma$ such that z and z' are both contained in Z_γ . As $Z_\gamma \in \mathcal{Z}$, it must be an irredundant set of factorizations, and so z and z' do not share any atoms. Hence U is also an irredundant set of factorizations, all of them contained in $Z_\ell(x)$. As a result, $U \in \mathcal{Z}$, and so U is an upper bound for the chain \mathcal{C} in \mathcal{Z} . Now we show that \mathcal{Z} has a maximal element. Pick some element $Z_1 \in \mathcal{Z}$. If it is maximal, then we are done. Otherwise we can pick $Z_2 \in \mathcal{Z}$ such that $Z_1 \subsetneq Z_2$. If Z_2 is maximal, we are done. Otherwise we can pick $Z_3 \in \mathcal{Z}$ such that $Z_2 \subsetneq Z_3$. If Z_3 is maximal, we are done. Continuing this process, this process either terminates, in which case we are done, or we have an infinite chain $Z_1 \subsetneq Z_2 \subsetneq \cdots \subset U$ of factorizations in \mathcal{Z} . Since each Z_i adds at least one new element to U , this implies that U must have infinitely many elements, contradicting Lemma 3.3. Thus \mathcal{Z} has a maximal element.

Let Z be a maximal element of \mathcal{Z} , that is, an irredundant subset of $Z_\ell(x)$ that is not properly contained in any other irredundant subset of $Z_\ell(x)$. Let A be the set consisting of all the atoms of M that appear in at least one factorization in Z . Because Z is finite, so is A . The maximality of Z guarantees that every factorization in $Z_\ell(x)$ shares an atom with at least one factorization in Z , and so every factorization in $Z_\ell(x)$ contains an atom from A . As a consequence, after setting

$$Z_a := \{z \in Z_\ell(x) \mid a \text{ appears in } z\}$$

for each $a \in A$, we obtain that $Z_\ell(x) = \bigcup_{a \in A} Z_a$. Since A is finite, the fact that $Z_\ell(x)$ contains infinitely many factorizations ensures the existence of $a \in A$ such that $|Z_a| = \infty$. By definition of Z_a , every factorization of Z_a contains the atom a . Therefore the set of factorizations $Z'_a := \{z - a \mid z \in Z_a\}$ is a subset of $Z_{\ell-1}(x - a)$. Since $|Z'_a| = |Z_a| = \infty$, it follows that $Z_{\ell-1}(x - a)$ is infinite, which contradicts the minimality of ℓ . Hence M is an LFFM. \square

The converse of Theorem 3.4 does not hold in general. The following example, which is a continuation of Example 3.1, illustrates this observation.

Example 3.5. Let $(p_n)_{n \geq 1}$ be a strictly increasing sequence of primes, and consider the positive monoid

$$M := \left\langle 1 + \frac{(-1)^n}{p_n} \mid n \in \mathbb{N} \right\rangle.$$

We have already seen in Example 3.1 that M is an atomic positive monoid with $\mathcal{A}(M) = \{a_n \mid n \in \mathbb{N}\}$, where $a_n := 1 + \frac{(-1)^n}{p_n}$ for every $n \in \mathbb{N}$. We have also observed in the same example that M is not co-well-ordered. We claim that M is an FFM and, therefore, an LFFM. To argue this, fix a nonzero $q \in M$. Take $n_q \in \mathbb{N}$ such that $p_k \nmid \mathbf{d}(q)$ for any $k \geq n_q$, and then set $N := \max\{n_q, q + 1\}$. Now suppose that the atom a_k appears in a factorization of q . We will argue that $k \leq N$. Write $q = \sum_{i=1}^n c_i a_i$ for some $n \in \mathbb{N}_{\geq k}$ and $c_1, \dots, c_n \in \mathbb{N}_0$ such that $c_k \neq 0$. If $p_k \mid \mathbf{d}(q)$, then $k < n_q \leq N$. Assume, on the other hand, that $p_k \nmid \mathbf{d}(q)$. This being the case, after applying p_k -adic valuations to both sides of the equality $q = \sum_{i=1}^n c_i a_i$, we find that $p_k \mid c_k$. Therefore $q \geq c_k a_k \geq p_k a_k = p_k + (-1)^k$, which implies that $k \leq p_k \leq q + 1 \leq N$. Hence $k \leq N$, as desired, and so q is divisible in M by only finitely many atoms. Since every element of M is divisible by only finitely many atoms in M , it follows from [31, Theorem 2] that M is an FFM and, in particular, an LFFM.

3.2. Some Concrete Examples

Although every FFM is an LFFM, the converse of this implication does not hold in general. As the following examples illustrate, there are positive monoids that are LFFM but do not even satisfy the ACCP.

Example 3.6.

- (a) Let $(p_n)_{n \geq 0}$ be the strictly increasing sequence whose underlying set is $\mathbb{P} \setminus \{2\}$. Then consider the positive monoid defined as follows:

$$M := \left\langle \frac{1}{2^n p_n} \mid n \in \mathbb{N}_0 \right\rangle.$$

This monoid is often called the Grams' monoid as it is the essential ingredient used by Grams in her construction of the first atomic domain that does not satisfy the ACCP (see [30] for more details and [26] for a recent generalization). It is well known that M is atomic with

$$\mathcal{A}(M) = \left\{ \frac{1}{2^n p_n} \mid n \in \mathbb{N}_0 \right\}$$

but does not satisfy the ACCP: indeed, $(\frac{1}{2^n} + M)_{n \geq 0}$ is an ascending chain of principal ideals that does not stabilize. Finally, since the sequence of defining generators of M is decreasing, M is a decreasing monoid and, therefore, M is an LFFM by virtue of Theorem 3.4.

- (b) Choose $q \in \mathbb{Q} \cap (0, 1)$ such that $q^{-1} \notin \mathbb{N}$, and consider the positive monoid $\mathbb{N}_0[q] := \langle q^n \mid n \in \mathbb{N}_0 \rangle$. Using the fact that $q^{-1} \notin \mathbb{N}$, we can argue that $\mathbb{N}_0[q]$ is atomic with $\mathcal{A}(\mathbb{N}_0[q]) = \{q^n \mid n \in \mathbb{N}_0\}$ (this is well known: see [24, Theorem 6.2] and also [15, Theorem 4.2]). On the other hand, $\mathbb{N}_0[q]$ does not satisfy the ACCP because $d(q)q^n = (d(q) - n(q))q^n + d(q)q^{n+1}$ for every $n \in \mathbb{N}_0$, where $n(q)$ and $d(q)$ denote the numerator and denominator of q , respectively. Lastly, note that because the sequence $(q^n)_{n \geq 0}$ is decreasing, Theorem 3.4 ensures that $\mathbb{N}_0[q]$ is an LFFM. Hence the positive monoid $\mathbb{N}_0[q]$ is an LFFM that does not satisfy the ACCP.

Thus, being an LFFM does not imply satisfying the ACCP and, therefore, not every LFFM is a BFM. Let us proceed to exhibit a positive monoid that is a BFM but not an LFFM.

Example 3.7. Consider the positive monoid $M := \{0\} \cup \mathbb{Q}_{\geq 1}$. One can readily verify that $\mathcal{A}(M) = \mathbb{Q} \cap [1, 2)$ and, therefore, that M is atomic. Since 0 is not a limit point of M^\bullet , it follows from [22, Proposition 4.5] that M is a BFM. In order to argue that M is not an LFFM, it suffices to observe that the element 3 has infinitely many length-2 factorizations in M : indeed, the equality $3 = (\frac{3}{2} - \frac{1}{n}) + (\frac{3}{2} + \frac{1}{n})$ yields a length-2 factorization of 3 for each $n \in \mathbb{N}$ with $n \geq 3$.

We have seen in Example 3.6 that being an LFFM is not a sufficient condition for satisfying the ACCP. On the other hand, even if a positive monoid is an LFFM satisfying the ACCP it still may not be a BFM. The following example sheds some light upon this observation.

Example 3.8. Consider the positive monoid $M = \langle \frac{1}{p} \mid p \in \mathbb{P} \rangle$. It is well known that $\mathcal{A}(M) = \{\frac{1}{p} \mid p \in \mathbb{P}\}$ and also that M satisfies the ACCP (see [3, Example 3.3] or [23, Proposition 4.2(2)] for more details). In addition, since the set of atoms of M is the underlying set of the decreasing sequence $(\frac{1}{p})_{p \in \mathbb{P}}$, it follows from Theorem 3.4 that M is an LFFM. Finally, observe that M is not a BFM: indeed, $Z(1) = \{p\frac{1}{p} \mid p \in \mathbb{P}\}$ and so $L(1) = \mathbb{P}$. Hence the positive monoid M is an LFFM, satisfies the ACCP, but is not a BFM.

4. Length-Finite Factorization Positive Semirings

4.1. A Class of Length-Finite Factorization Positive Semirings

We can use any positive monoid to construct positive semirings by using certain exponentiation. Let M be a positive monoid. Following [4], we set

$$E(M) := \langle e^m \mid m \in M \rangle;$$

that is, $E(M)$ is the positive monoid generated by the set of positive real numbers $e(M) := \{e^m \mid m \in M\}$. We observe that $1 \in E(M)$ and also that $E(M)$ is closed under multiplication. Hence $E(M)$ is a positive semiring. Also,

the fact that $1 = \min E(M)^\bullet$, implies that the multiplicative monoid $E(M)^\bullet$ is reduced, and so $E(M)$ is a reduced positive semiring. Observe that $e(M)$ is a multiplicative monoid that is naturally isomorphic to the positive monoid M .

Suppose now that M consists of algebraic numbers, which is the case we primarily target here. Then it follows from Lindemann-Weierstrass Theorem that the set $e(M)$ is linearly independent over the algebraic numbers (see [6, Chapter 1]), and so the additive monoid $E(M)$ is a free commutative monoid on $e(M)$. Additionally, it follows from [4, Lemma 2.7]² that $e(M)$ is a divisor-closed submonoid of the multiplicative monoid of $E(M)$. In this case, every nonzero element $r \in E(M)$ can be written uniquely as $r = c_1 e^{m_1} + \dots + c_k e^{m_k}$ for some coefficients $c_1, \dots, c_k \in \mathbb{N}$ and exponents $m_1, \dots, m_k \in M$ such that $m_1 < \dots < m_k$: as for polynomials, we call $\text{LC}(r) := c_k$ and $\deg r := m_k$ the *leading coefficient* and the *degree* of r , respectively. In addition, we call $S_E(r) := \{m_1, \dots, m_k\}$ the *exponent set* of r .

It turns out that the properties of satisfying the ACCP, being a BFM, and being an FFM ascend from a positive monoid M to $E(M)$ provided that every element of M is algebraic. We proceed to prove this.

Proposition 4.1. *Let M be a positive monoid consisting of algebraic numbers. Then the following statements hold.*

- (1) *If M satisfies the ACCP, then $E(M)$ satisfies the ACCP.*
- (2) *If M is a BFM, then $E(M)$ is a BFS.*
- (3) *If M is an FFM, then $E(M)$ is an FFS.*

Proof. (1) Assume that M satisfies the ACCP. Now suppose, by way of contradiction, that $E(M)$ does not satisfy the ACCP. Then there exists an element $s \in E(M)$ and two sequences $(s_n)_{n \geq 1}$ and $(t_n)_{n \geq 1}$ whose terms are nonunits of $E(M)$ such that $s = t_n \prod_{i=1}^n s_i$ for every $n \in \mathbb{N}$. Therefore, for each $n \in \mathbb{N}$,

$$\deg s = \deg t_n + \sum_{i=1}^n \deg s_i \quad \text{and} \quad \text{LC}(s) = \text{LC}(t_n) \prod_{i=1}^n \text{LC}(s_i).$$

Let ℓ be the length of the only factorization of $\text{LC}(s)$ in \mathbb{N} . Then at most ℓ of the real numbers $\text{LC}(s_1), \dots, \text{LC}(s_n), \text{LC}(t_n)$ can be greater than one. Since the terms s_k and t_k are nonunits for every $k \in \mathbb{N}$, at least $n + 1 - \ell$ of the real numbers $\deg s_1, \dots, \deg s_n, \deg t_n$ must be nonzero. Thus, at least $n - \ell$ of the real numbers $\deg s_1, \dots, \deg s_n$ must be nonzero. Now consider the ascending chain of principal ideals $\deg t_1 + M, \deg t_2 + M, \dots$ in M . Since at least $n - \ell$ of the real numbers $\deg s_1, \dots, \deg s_n$ must be nonzero for arbitrarily large n , and $\deg s_n = \deg t_{n-1} - \deg t_n$, this means that this ascending chain of principal ideals must not stabilize. Thus, M does not satisfy the ACCP, a contradiction. Thus $E(M)$ satisfies the ACCP.

²Although [4, Lemma 2.7] is stated for positive monoids, it only applies to positive monoids consisting of algebraic numbers.

(2) Since M is a BFM, it must satisfy the ACCP, and so it follows from part (1) that $E(M)$ also satisfies the ACCP. Thus, $E(M)$ is atomic. Suppose, by way of contradiction, that $E(M)$ is not a BFS. Then there exists $q \in E(M)$ such that there is no bound to the lengths of factorizations of q . Suppose $q = q_1 \cdots q_n$ for nonunits q_1, \dots, q_n . Then

$$\deg q = \sum_{i=1}^n \deg q_i \quad \text{and} \quad \text{LC}(q) = \prod_{i=1}^n \text{LC}(q_i).$$

Let ℓ be the length of the factorization $\text{LC}(q)$ in \mathbb{N} . Then we see that at most ℓ of the real numbers $\text{LC}(q_1), \dots, \text{LC}(q_n)$ can be greater than one. Since the q_1, \dots, q_n are nonunits, at least $n - \ell$ of them must have degree different from zero. Therefore, after replacing each $\deg q_i$ by one of its factorizations in M , we obtain a factorization for $\deg q$ with length at least $n - \ell$. But as n approaches infinity, $n - \ell$ approaches infinity, which means that there is no upper bound to the length of factorizations of $\deg q$ in M , contradicting the fact that M is a BFM. Hence $E(M)$ is a BFS.

(3) Assume that M is an FFM. Since M is a BFM, it follows from (2) that $E(M)$ is a BFS. We claim that, for any positive integers ℓ and $q \in E(M)$, there are finitely many ways to choose $a_1, \dots, a_\ell \in \mathcal{A}(E(M))$ such that $q = a_1 \cdots a_\ell$. Since $E(M)$ is a BFS, this would imply that $E(M)$ is also an FFS.

First, we prove that there are finitely many choices for a_1 . Set $m := m_1 + \cdots + m_\ell$, where $m_1 \in S_E(a_1), \dots, m_\ell \in S_E(a_\ell)$. Observe that $m \in S_E(q)$. Therefore m_1 is the sum of a subset of atoms in a factorization of m with respect to monoid M . Since M is an FFM, there are finitely many such factorizations. Given any such factorization of length ℓ_m , there are at most 2^{ℓ_m} ways to choose m_1 . Since there are finitely many factorizations, there are finitely many ways to choose m_1 . Thus, there are finitely many ways to choose the exponent set of a_1 .

Now observe that $a_1 \leq q$. Therefore $c \leq q$ for every coefficient c of a_1 . If a_1 has ℓ_1 terms, then there are at most q^{ℓ_1} ways to choose all the coefficients of a_1 , so there are finitely many ways to choose a_1 . By symmetry, there are also finitely many ways to choose a_2, \dots, a_ℓ . Let N be a positive integer such that there are less than N ways to choose each of a_1, \dots, a_ℓ . As a result, q has at most N^ℓ factorizations of length ℓ . Hence we conclude that $E(M)$ is an FFS. \square

We now define another method to construct positive semirings, the monoid semiring of M over the naturals. As seen from the definition, this construction is quite similar to the $E(M)$ construction.

Definition. Let M be a positive monoid consisting of algebraic numbers. Then, the *monoid semiring of M over the naturals*, denoted by $\mathbb{N}_0[M]$, is the

semiring of polynomials with coefficients in \mathbb{N}_0 and exponents in M , i.e.

$$\mathbb{N}_0[M] = \left\{ \sum_{i=1}^k c_i x^{m_i} \mid c_i, k \in \mathbb{N}, m_i \in M \right\}.$$

It turns out that for any positive monoid consisting of algebraic numbers, the semirings $E(M)$ and $\mathbb{N}_0[M]$ are isomorphic.

Proposition 4.2. *Let M be a positive monoid consisting of algebraic numbers. Then, the semirings $E(M)$ and $\mathbb{N}_0[M]$ are isomorphic.*

Proof. Since M consists of algebraic numbers, the set $\{x^m \mid m \in M\}$ is linearly independent over the natural numbers. Suppose

$$c_1 x^{m_1} + \cdots + c_k x^{m_k} = 0,$$

where c_1, \dots, c_k are nonzero and $k \geq 1$, and m_1, \dots, m_k are pairwise distinct. Then

$$c_1 e^{m_1} + \cdots + c_k e^{m_k} = 0,$$

so e^{m_1}, \dots, e^{m_k} are linearly dependent, but this contradicts the Lindemann-Weierstrass Theorem.

Thus, every element of $\mathbb{N}_0[M]$ can be represented uniquely as $s = c_1 x^{m_1} + \cdots + c_k x^{m_k}$ for some coefficients $c_1, \dots, c_k \in \mathbb{N}$ and exponents $m_1, \dots, m_k \in M$ such that $m_1 < \cdots < m_k$. Then, by defining $\varphi(s) = c_1 e^{m_1} + \cdots + c_k e^{m_k}$, we get an isomorphism from $\mathbb{N}_0[M]$ to $E(M)$. \square

Proposition 4.2 shows that the results of Proposition 4.1 also hold true for $\mathbb{N}_0[M]$. We can use Proposition 4.1 to construct positive semirings satisfying some desired prescribed properties that are easier to achieve with positive monoids. The following example illustrates this observation.

Example 4.3. Consider the positive monoid $M := \{0\} \cup \mathbb{Q}_{\geq 1}$ (which is also a positive semiring). We have seen in Example 3.7 that M is a BFM with $\mathcal{A}(M) = \mathbb{Q} \cap [1, 2)$ and also that M is not an LFFM. Since M is a positive monoid consisting of algebraic numbers, it follows from Proposition 4.1 that the exponentiation semiring $E(M)$ is also a BFS. However, also from the fact that M consists of algebraic numbers, we obtain via [4, Lemma 2.7] that e^M is a divisor-closed submonoid of $E(M)$. Since e^M is isomorphic to M , the fact that M is not an LFFM guarantees that $E(M)$ is not an LFFS. Hence $E(M)$ is a positive semiring that is a BFS but not an LFFS.

We have seen in Proposition 4.1 that the properties of satisfying the ACCP, having bounded factorizations, and having finite factorizations all ascend from a positive monoid M to its exponentiation semiring $E(M)$ provided that M consists of algebraic elements. We still do not know, however, whether this is also the case for the length-finite factorization property.

Question 4.4. *For a positive monoid M consisting of algebraic numbers, is $E(M)$ an LFFS provided that M is an LFFM?*

4.2. Further Concrete Examples

In this final subsection, we provide two further classes of positive semirings. The first class consists of positive semirings that generalize the positive semiring M used in Example 4.3: this class contains positive semirings that, as the positive semiring $E(M)$ in Example 4.3, are BFS but not LFFS. The second class consists of positive semirings that are FFS and, therefore, both BFS and LFFS.

Example 4.5. For $r \in \mathbb{R}_{\geq 1}$, consider the positive monoid $S_r := \mathbb{N}_0 \cup \mathbb{R}_{\geq r}$. Because S_r contains 1 and is closed under multiplication, it is a positive semiring. If $r = 1$, then for each $s \in \mathbb{R}_{>1}$ the equality $s = (s \frac{n}{n+1})(\frac{n+1}{n})$ for $n \in \mathbb{N}$ large enough implies that s is not an atom of the multiplicative monoid S_r^\bullet and, therefore, we conclude that S_r is antimatter. For $r > 1$, it follows from [4, Theorem 5.1] that

$$\mathcal{A}_+(S_r) = (\{1\} \cup [r, r+1)) \setminus \{r\} \quad \text{and} \quad \mathcal{A}_\times(S_r) = (\mathbb{P}_{<r^2} \cup [r, r^2)) \setminus \mathbb{P} \cdot (S_r)_{>1},$$

and also that both the additive and the multiplicative monoids of S_r are BFMs. Even when $r > 1$, neither the additive nor the multiplicative monoid of S_r may be FFM (the case $r = 2$ was illustrated in [4, Example 6.4]). We proceed to argue that S_r is not even an LFFS when $r > 1$. To verify that the additive monoid of S_r is not an LFFM, it suffices to observe that the formal sum $(r + \frac{1}{n}) + (r + 1 - \frac{1}{n})$ is a length-2 factorization of $2r + 1$ in $(S_r, +)$ for every $n \in \mathbb{N}$ sufficiently large. In a similar way, we can argue that the multiplicative monoid (S_r^\bullet, \cdot) is not an LFFM since the formal product $(s \frac{n+1}{n})(s \frac{n}{n+1})$ for some $s \in (r, r^2)$ is a length-2 factorization of s^2 for every sufficiently large $n \in \mathbb{N}$.

The positive monoid in Example 3.6(b) is also a positive semiring, and the more general case corresponding to algebraic parameters was studied in [15] and, more recently, in [1]. However, only the additive structure of these semirings has been systematically investigated.

Example 4.6. Let $\alpha \in \mathbb{R}_{>0}$, and consider the positive monoid $\mathbb{N}_0[\alpha] := \langle \alpha^n \mid n \in \mathbb{N}_0 \rangle$ (the special case when α is rational was briefly considered in Example 3.6(b)). Observe that $1 \in \mathbb{N}_0[\alpha]$ and also that $\mathbb{N}_0[\alpha]$ is closed under multiplication. Hence $\mathbb{N}_0[\alpha]$ is a positive semiring. There are choices of α such that the additive monoid $\mathbb{N}_0[\alpha]$ is antimatter (for instance, $\alpha = \frac{1}{2}$) and there are choices of α such that the additive monoid $\mathbb{N}_0[\alpha]$ is atomic but does not satisfy the ACCP (for instance, $\alpha = \frac{2}{3}$). In addition, it follows from [15, Theorem 4.11] that the positive monoid $\mathbb{N}_0[\alpha]$ satisfies the ACCP if and only if it is a BFM if and only if it is an FFM. However, not too much is known about the atomic structure of the multiplicative monoid of $\mathbb{N}_0[\alpha]$.

In connection with Example 4.6, we were not able to settle the following question.

Question 4.7. For which $q \in \mathbb{Q}_{>0}$ is the positive semiring $\mathbb{N}_0[q]$ an LFFS?

Acknowledgments

We are grateful to our mentors, Prof. Jim Coykendall and Dr. Felix Gotti, for proposing this project and for their guidance during the preparation of this project. While working on this paper, we were part of PRIMES-USA, a year-long math research program hosted by MIT. We would like to express our collective gratitude to the MIT PRIMES program for arranging such an engaging research experience.

References

- [1] K. Ajran, J. Bringas, B. Li, E. Singer, and M. Tirador: *Factorization in additive monoids of evaluation polynomial semirings*, Comm. Algebra **51** (2023) 4347–4362.
- [2] D. D. Anderson, D. F. Anderson, and M. Zafrullah: *Factorization in integral domains*, J. Pure Appl. Algebra **69** (1990) 1–19.
- [3] D. F. Anderson and F. Gotti: *Bounded and finite factorization domains*. In: Rings, Monoids, and Module Theory (Eds. A. Badawi and J. Coykendall) pp. 7–57 Springer Proceedings in Mathematics & Statistics, Vol. **382**, Singapore, 2022.
- [4] N. R. Baeth, S. T. Chapman, and F. Gotti: *Bi-atomic classes of positive semirings*, Semigroup Forum **103** (2021) 1–23.
- [5] N. R. Baeth and F. Gotti: *Factorizations in upper triangular matrices over information semialgebras*, J. Algebra **562** (2020) 466–496.
- [6] A. Baker: *Transcendental Number Theory* (2nd ed.), Cambridge Mathematical Library, Cambridge University Press, 1990.
- [7] A. Bu, J. Vulakh, and A. Zhao: *Length-factoriality and pure irreducibility*, Comm. Algebra **51** (2023) 3745–3755.
- [8] M. Bras-Amorós: *Increasingly enumerable submonoids of R : Music Theory as a Unifying Theme*, Amer. Math. Monthly **127** (2020) 33–44.
- [9] M. Bras-Amorós and M. Gotti: *Atomicity and density of Puiseux monoids*, Comm. Algebra **49** (2021) 1560–1570.
- [10] S. T. Chapman and J. Coykendall: *Half-factorial domains, a survey*. In: *Non-Noetherian Commutative Ring Theory* (Eds. S. T. Chapman and S. Glaz) pp. 97–115, Mathematics and Its Applications, vol. 520, Kluwer Academic Publishers, Springer, Boston 2000.
- [11] S. T. Chapman, J. Coykendall, F. Gotti, and W. W. Smith: *Length-factoriality in commutative monoids and integral domains*, J. Algebra **578** (2021) 186–212.
- [12] S. T. Chapman, F. Gotti, and M. Gotti: *Factorization invariants of Puiseux monoids generated by geometric sequences*, Comm. Algebra **48** (2020) 380–396.
- [13] S. T. Chapman and M. Gotti: *Atomicity of positive monoids*. Submitted. Preprint on arXiv: <https://arxiv.org/abs/2108.05561>.
- [14] P. M. Cohn: *Bezout rings and their subrings*, Proc. Cambridge Philos. Soc. **64** (1968) 251–264.
- [15] J. Correa-Morris and F. Gotti: *On the additive structure of algebraic valuations of polynomial semirings*, J. Pure Appl. Algebra **226** (2022) 107104.
- [16] J. Coykendall, D. E. Dobbs, and B. Mullins: *On integral domains with no atoms*, Comm. Algebra **27** (1999) 5813–5831.
- [17] J. Coykendall and W. W. Smith: *On unique factorization domains*, J. Algebra **332** (2011) 62–70.
- [18] W. Gao, C. Liu, S. Tringali, and Q. Zhong: *On half-factoriality of transfer Krull monoids*, Comm. Algebra **49** (2021) 409–420.
- [19] A. Geroldinger and F. Halter-Koch: *Non-unique Factorizations: Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics Vol. **278**, Chapman & Hall/CRC, Boca Raton, 2006.

- [20] A. Geroldinger and Q. Zhong: *A characterization of length-factorial Krull monoids*, New York J. Math. **27** (2021) 1347–1374.
- [21] J. S. Golan: *Semirings and their Applications*, Kluwer Academic Publishers, 1999.
- [22] F. Gotti: *Increasing positive monoids of ordered fields are FF-monoids*, J. Algebra **518** (2019) 40–56.
- [23] F. Gotti: *On semigroup algebras with rational exponents*, Comm. Algebra **50** (2022) 3–18.
- [24] F. Gotti and M. Gotti: *Atomicity and boundedness of monotone Puiseux monoids*, Semigroup Forum **96** (2018) 536–552.
- [25] F. Gotti and M. Gotti: *On the molecules of numerical semigroups, Puiseux monoids, and Puiseux algebras*. In: Numerical Semigroups (Eds. V. Barucci, S. T. Chapman, M. D’Anna, and R. Fröberg) pp. 141–161. Springer INdAM Series, Vol. **40**, Switzerland, 2020.
- [26] F. Gotti and B. Li: *Atomic semigroup rings and the ascending chain condition on principal ideals*, Proc. Amer. Math. Soc. **151** (2023) 2291–2302.
- [27] F. Gotti and H. Polo: *On the arithmetic of polynomial semidomains*, Forum Math. **35** (2023) 1179–1197.
- [28] F. Gotti and H. Polo: *On the subatomicity of polynomial semidomains*. In: Algebraic, Number Theoretic, and Topological Aspects of Ring Theory (Eds. J. L. Chabert, M. Fontana, S. Frisch, S. Glaz, and K. Johnson) pp. 197–212. Springer International Publishing, Switzerland, 2023.
- [29] F. Gotti and J. Vulakh: *On the atomic structure of torsion-free monoids*, Semigroup Forum (to appear). Preprint on arXiv: <https://arxiv.org/abs/2212.08347>
- [30] A. Grams: *Atomic rings and the ascending chain condition for principal ideals*, Proc. Cambridge Philos. Soc. **75** (1974) 321–329.
- [31] F. Halter-Koch: *Finiteness theorems for factorizations*, Semigroup Forum **44** (1992) 112–117.
- [32] N. Jiang, B. Li, and S. Zhu: *On the primality and elasticity of algebraic valuations of cyclic free semirings*, Internat. J. Algebra Comput. **33** (2023) 197–210.
- [33] H. Polo: *A characterization of finite factorization positive monoids*, Commun. Korean Math. Soc. **37** (2022) 669–679.
- [34] A. Zaks: *Half-factorial domains*, Bull. Amer. Math. Soc. **82** (1976) 721–723.
- [35] S. Zhu: *Factorizations in evaluation monoids of Laurent semirings*, Comm. Algebra **50** (2022) 2719–2730.

HENRY JIANG
DETROIT COUNTRY DAY SCHOOL
BEVERLY HILLS, MI 48025
Email address: jiangstem@gmail.com

SHIHAN KANUNGO
HENRY M. GUNN SCHOOL
PALO ALTO, CA 94306
Email address: shihankanungo@gmail.com

HWISOO KIM
PHILLIPS ACADEMY
ANDOVER, MA 01810
Email address: hkim25@andover.edu