

❖ MACHINE LEARNING - WORKSHEET 1 SOLUTION

(* Answers are represented in red font)

In Q1 to Q11, only one option is correct, choose the correct option:

1. Which of the following methods do we use to find the best fit line for data in Linear Regression?

- A) Least Square Error**
- B) Maximum Likelihood
- C) Logarithmic Loss
- D) Both A and B

2. Which of the following statement is true about outliers in linear regression?

- A) Linear regression is sensitive to outliers**
- B) linear regression is not sensitive to outliers
- C) Can't say
- D) none of these

3. A line falls from left to right if a slope is _____?

- A) Positive
- B) Negative**
- C) Zero
- D) Undefined

4. Which of the following will have symmetric relation between dependent variable and independent variable?

- A) Regression
- B) Correlation**
- C) Both of them
- D) None of these

5. Which of the following is the reason for over fitting condition?

- A) High bias and high variance
- B) Low bias and low variance
- C) Low bias and high variance**
- D) none of these

6. If output involves label then that model is called as:

- A) Descriptive model
- B) Predictive model**
- C) Reinforcement learning
- D) All of the above

7. Lasso and Ridge regression techniques belong to _____?

- A) Cross validation
- B) Removing outliers
- C) SMOTE
- D) Regularization**

8. To overcome with imbalance dataset which technique can be used?

- A) Cross validation
- B) Regularization
- C) Kernel
- D) SMOTE**

9. The AUC Receiver Operator Characteristic (AUCROC) curve is an evaluation metric for binary

classification problems. It uses _____ to make graph?

- A) TPR and FPR
- B) Sensitivity and precision
- C) Sensitivity and Specificity**
- D) Recall and precision

10. In AUC Receiver Operator Characteristic (AUCROC) curve for the better model area under the

curve should be less.

- A) True
- B) False**

11. Pick the feature extraction from below:

- A) Construction bag of words from a email
- B) Apply PCA to project high dimensional data**
- C) Removing stop words
- D) Forward selection

In Q12, more than one options are correct, choose all the correct options:

12. Which of the following is true about Normal Equation used to compute the coefficient of the Linear

Regression?

- A) We don't have to choose the learning rate.**

B) It becomes slow when number of features is very large.

C) We need to iterate.

D) It does not make use of dependent variable.

Q13 and Q15 are subjective answer type questions, Answer them briefly.

13. Explain the term regularization?

Solution:

Regularization is one of the most important concepts of machine learning. It is a technique to prevent the model from overfitting by adding extra information to it.

May it be in statistics or mathematics or finance – particularly in machine learning and inverse problems – regularization is any modification that one makes in a learning algorithm that is intended to not reduce its training error but its generalization error. In layman's language, we can say the word "regularize" simply means to make things acceptable or regular.

So, in what sort of situations do we use regularization? Sometimes, we encounter a situation where our machine learning algorithm fails to perform well on testing data but works well on training data exceptionally. This is a situation where regularization comes into play and solves the issues. It discourages learning a flexible or more complex model in order to prevent overfitting.

This technique can be used in such a way that it will allow us to maintain all variables or features in the model by reducing the magnitude of the variables. Hence, regularization maintains accuracy as well as generalization of the model.

In the same way cancer treatment destroys the cancer cells rather than affecting the healthy cells in the body, the regularization approach attacks the noise alone without affecting the signal.

Working :

Regularization works by adding a penalty or complexity term to the complex model.

Let's consider the simple linear regression equation:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \dots + \beta_n x_n + b$$

where, Y represents the value to be predicted

X1, X2, ...Xn are the features for Y.

$\beta_0, \beta_1, \dots, \beta_n$ are the weights or magnitude attached to the features, respectively. Here, b represents the intercept.

Linear regression models try to optimize the β_0 and b to minimize the cost function.

So, the equation for the cost function for the linear model is as stated below:

$$\sum_{i=1}^M (y_i - y'_i)^2 = \sum_{i=1}^M (y_i - \sum_{j=0}^n \beta_j * X_{ij})^2$$

Now, we will add a loss function and optimize parameter to make the model that can predict the accurate value of Y. The loss function for the linear regression is called as RSS or Residual sum of squares.

14. Which particular algorithms are used for regularization?

Solution:

- **Ridge Regression**
- **Lasso Regression**

Ridge Regression:

Ridge regression is one of the types of linear regression in which a small amount of bias is introduced so that we can get better long-term predictions.

Ridge regression is a regularization technique, which is used to reduce the complexity of the model. It is also called as L2 regularization.

In this technique, the cost function is altered by adding the penalty term to it. The amount of bias added to the model is called Ridge Regression penalty. We can calculate it by multiplying with the lambda to the squared weight of each individual feature.

The equation for the cost function in ridge regression will be:

$$\sum_{i=1}^M (y_i - y'_i)^2 = \sum_{i=1}^M \left(y_i - \sum_{j=0}^n \beta_j * x_{ij} \right)^2 + \lambda \sum_{j=0}^n \beta_j^2$$

In the above equation, the penalty term regularizes the coefficients of the model, and hence ridge regression reduces the amplitudes of the coefficients that decreases the complexity of the model.

As we can see from the above equation, if the values of λ tend to zero, the equation becomes the cost function of the linear regression model. Hence, for the minimum value of λ , the model will resemble the linear regression model.

A general linear or polynomial regression will fail if there is high collinearity between the independent variables, so to solve such problems, Ridge regression can be used.

It helps to solve the problems if we have more parameters than samples.

Lasso Regression:

Lasso regression is another regularization technique to reduce the complexity of the model. It stands for Least Absolute and Selection Operator.

It is similar to the Ridge Regression except that the penalty term contains only the absolute weights instead of a square of weights.

Since it takes absolute values, hence, it can shrink the slope to 0, whereas Ridge Regression can only shrink it near to 0.

It is also called as L1 regularization. The equation for the cost function of Lasso regression will be:

$$\sum_{i=1}^M (y_i - y'_i)^2 = \sum_{i=1}^M \left(y_i - \sum_{j=0}^n \beta_j * x_{ij} \right)^2 + \lambda \sum_{j=0}^n |\beta_j|$$

Some of the features in this technique are completely neglected for model evaluation.

Hence, the Lasso regression can help us to reduce the overfitting in the model as well as the feature selection.

15. Explain the term error present in linear regression equation?

An error term is a residual variable produced by a statistical or mathematical model, which is created when the model does not fully represent the actual relationship between the independent variables and the dependent variables. As a result of this incomplete relationship, the error term is the amount at which the equation may differ during empirical analysis.

The error term is also known as the residual, disturbance, or remainder term, and is variously represented in models by the letters e, ε, or u.

An error term essentially means that the model is not completely accurate and results in differing results during real-world applications. For example, assume there is a multiple linear regression function that takes the following form:

$$Y = \alpha X + \beta \rho + \epsilon$$

where:

α, β = Constant parameters

X, ρ = Independent variables

ϵ = Error term

When the actual Y differs from the expected or predicted Y in the model during an empirical test, then the error term does not equal 0, which means there are other factors that influence Y.