

Topology and Groups - MATH0074

Based on lectures by Dr. Lars Louder

Notes taken by Imran Radzi

(Revision) notes based on the Autumn 2021 Topology and Groups lectures by Dr. Lars Louder. Some parts marked with (*) are taken from Hatcher's Algebraic Topology.

Contents

1	Point-set Topology	1
1.1	Preliminaries	1
1.2	Connectedness	2
1.3	Compactness	3
1.4	Quotient spaces	4
2	Homotopy	5
2.1	Homotopy	5
2.2	Paths and path-homotopy	7
3	Covering spaces	8

1 Point-set Topology

1.1 Preliminaries

Definition (Topological space). A topological space is a pair (X, \mathcal{T}) such that

1. X is a set
2. $\mathcal{T} \subset \mathcal{P}(X)$ is a collection of subsets of X
3. $\emptyset \in \mathcal{T}, X \in \mathcal{T}$
4. \mathcal{T} is closed under finite intersections and arbitrary unions

Definition (Open neighbourhood). If $x \in X$, U open in X , and $x \in U$, then U is an *open neighbourhood* of x .

Definition (Hausdorff spaces). A topological space (X, \mathcal{T}) is *Hausdorff* if $\forall x, y \in X$, there exists U, V open neighbourhoods of x, y respectively such that $U \cap V = \emptyset$.

Definition (Homeomorphisms). A map $f : X \rightarrow Y$ is a *homeomorphism* if

1. f is bijective
2. f is continuous
3. f^{-1} is continuous

Definition (Continuous maps). A map $f : X \rightarrow Y$ is continuous if $\forall U$ (open) $\subset Y$, $f^{-1}(U)$ is open in X .

Definition. If \mathcal{T} and \mathcal{T}' are topologies on X such that $\mathcal{T} \subsetneq \mathcal{T}'$ then \mathcal{T}' is *finer* than \mathcal{T} , and \mathcal{T} is *coarser* than \mathcal{T}' .

Proposition. $\text{id} : (X, \mathcal{T}) \rightarrow (X, \mathcal{T}')$ is continuous if and only if \mathcal{T} is finer than \mathcal{T}' .

Definition (Subspace topology). If X is a topological space, $Y \subset X$, the subspace topology on Y is defined by

$$U \text{ open in } Y \iff \exists V \text{ open in } X \text{ such that } U = Y \cap V$$

Definition. If a map $f : X \rightarrow Y$ is continuous, the *image* of f is the set

$$f(X) = \{f(x) \mid x \in X\} \subset Y$$

with the subspace topology.

Definition (Product topology). Let X, Y be spaces. The *product topology* on $X \times Y$ is the smallest (coarsest) topology making the projections

$$p_X : X \times Y \rightarrow X, \quad p_Y : X \times Y \rightarrow Y$$

continuous.

Proposition. Product of Hausdorff spaces is Hausdorff.

1.2 Connectedness

Definition (Connectedness). A space X is *disconnected* if there exists a surjective continuous map $f : X \rightarrow \{p_1, p_2\}$. A space is *connected* if every continuous function $f : X \rightarrow \{p_1, p_2\}$ is constant.

Definition. A pair of sets $U, V \subset X$ is said to disconnect X if they are non-empty, disjoint, $U \cup V = X$ and both are open.

Definition. X is disconnected if there exists U, V which disconnect X .

Definition (Path). A *path* in X is a continuous map $\gamma : [0, 1] \rightarrow X$. γ is a path from $\gamma(0)$ to $\gamma(1)$. $a, b \in X$ are said to be connected by a path if there is a path from a to b .

Definition (Path-connectedness). A space X is *path-connected* if for all x, y , there exists

$$\gamma : [0, 1] \rightarrow X \text{ such that } \gamma(0) = x, \gamma(1) = y$$

or equivalently,

Definition. We say X is path-connected if there exists a unique equivalence class, where the equivalence relation \sim is defined $a \sim b$ if and only if there exists a path from a to b .

Proposition. Suppose X is connected. Then, if $f : X \rightarrow Y$, then $f(X) \subset Y$ is connected.

Proposition. $[0, 1]$ is connected.

Corollary. If X is path-connected, then X is connected.

Definition. $X \subset \mathbb{R}$ is an *interval* if $a \leq b \leq c$, $a, c \in X \implies b \in X$.

Proposition. A subset of \mathbb{R} is connected if and only if it is an interval.

Definition (Locally (path) connected). A space X is locally (path) connected at a point p if for every open neighbourhood U of p , there exists a (path) connected open neighbourhood V of p such that $p \in V \subset U$.

Proposition. If X is locally path-connected then the path components of X are open.

Proposition. If X is connected and locally path-connected, then X is path connected.

1.3 Compactness

Definition (Open cover). An *open cover* of a space X is a collection of open sets \mathcal{U} such that

$$X = \bigcup_{U \in \mathcal{U}} U$$

Definition. A space X is *compact* if every open cover has a finite subcover.

Lemma. Closed subset of compact spaces are compact.

Theorem. If X, Y are compact, then $X \times Y$ is compact.

Theorem (Heine-Borel theorem). $X \subset \mathbb{R}^n$ is compact if and only if X is closed and bounded.

Theorem. $[0, 1]$ is compact.

Theorem. If $f : X \rightarrow Y$ is continuous, X compact, then $f(X) \subset Y$ is compact with respect to the subspace topology.

Proposition. If $C \subset Y$ is compact, Y Hausdorff, then C is closed.

Proposition. If $f : X \rightarrow Y$ is a continuous bijection, X compact, Y Hausdorff, then f is a homeomorphism

1.4 Quotient spaces

Definition (Quotient map). Let $q : X \rightarrow Y$ be a continuous surjection. Then q is a *quotient map* if $q^{-1}(U)$ is open if and only if U is open. (A bijective quotient map is a homeomorphism)

Definition (Quotient space). Let X be a space, and \sim an equivalence relation on X , and $q : X \rightarrow X/\sim = Y$ the quotient map. The quotient topology on Y is defined by U open in Y if and only if $q^{-1}(U)$ is open in X .

Lemma.

$$\begin{array}{ccc} X & \xrightarrow{f} & Z \\ & \searrow q & \uparrow h \\ & & Y \end{array}$$

Let f be continuous, and suppose f factors through $q : X \rightarrow Y$, a quotient map, i.e., $\exists h : Y \rightarrow Z$ such that $h \circ q = f$. Then h is continuous.

Proposition. Let $f : X \rightarrow Y$ be a continuous surjection with X compact, Y Hausdorff. Then f is a quotient map.

Definition (Disjoint union). Let X_1, X_2 be topological spaces. The *disjoint union* of X_1 and X_2 , $X_1 \sqcup X_2$ is the space with the underlying set $X_1 \sqcup X_2$, with U open in $X_1 \sqcup X_2$ if and only if $U \cap X_1$ is open in X_1 , and $U \cap X_2$ is open in X_2 .

Definition (Cell complex). A *cell complex* is a space built up inductively, as follows

1. ($n = 0$) We start with a discrete set $X^{(0)}$ consisting of points, which we call 0-cells $\{e_i^0 \mid i \in I_0\}$, $e_i^0 \cong pt$. $X^{(0)} = \bigsqcup_i e_i^0$ is called the 0-skeleton.
2. ($n > 0$) We add a (possibly empty) subset of n -cells $\{e_i^n \mid i \in I_n\}$ $e_i^n \cong D^n$, the n -dimensional disk, and a continuous map

$$\phi_i^n : \partial e_i^n \cong S^{n-1} \rightarrow X^{(n-1)}$$

and here the n -skeleton is

$$X^{(n)} = X^{(n-1)} \sqcup \bigsqcup e_i^n / \sim$$

A space X is a cell complex if there exists $X^{(0)} \subset X^{(1)} \subset \dots$ as above, with the condition that U is open in X if and only if $X^{(n)} \cap U$ is open for all n .

$X^{(0)} \subseteq X^{(1)} \subseteq \dots$ is called the *cell decomposition* of X .

Definition (Presentation complex). text

Definition (Cayley graph). text

2 Homotopy

2.1 Homotopy

Definition. Let (X, A) be a pair of spaces, where $A \subseteq X$, $f_0, f_1 : X \rightarrow Y$. We say f_0 and f_1 are *homotopic relative to A* if there exists a

function $F : X \times I \rightarrow Y$ such that $F(-, 0) = f_0$, $F(-, 1) = f_1$ and $F(a, t) = f_0(a) = f_1(a)$ for all t . In this case we write $f_0 \simeq_A f_1$.

If $A = \emptyset$ then we say f_0 and f_1 are *homotopic* and write $f_0 \simeq f_1$.

Lemma (*). A function defined on the union of two closed sets is continuous if it is continuous when restricted to each of the closed sets separately.

Proposition. Any two continuous maps $f_0, f_1 : X \rightarrow \mathbb{R}^n$ are homotopic via the homotopy

$$F(x, t) = tf_1(x) + (1 - t)f_0(x)$$

Definition (Homotopy equivalence). Two spaces X and Y are *homotopy equivalent* if there exists $f : X \rightarrow Y$, $g : Y \rightarrow X$ such that $f \circ g \simeq \text{id}_Y$, $g \circ f \simeq \text{id}_X$. In this case, we write $X \simeq Y$.

Proposition. Homotopy equivalence is an equivalence relation on (topological) spaces.

Proposition. $\mathbb{R}^n \simeq pt$

Definition. A space X is *contractible* if $X \simeq pt$, or in other words, $\text{id} : X \rightarrow X$ is homotopic to a constant map. In this case the map id_X is said to be *null-homotopic*.

Proposition. $\mathbb{R}^n \setminus pt \simeq S^{n-1}$

Proposition. If X is contractible then X is path-connected.

Definition (Retract). Let $A \subseteq X$ be a subspace. A is a *retract* of X if there exists a continuous map $f : X \rightarrow A$ (retraction) such that $f|_A = \text{id}_A$. A is a *deformation retract* of X if there exists such a function r such that r is homotopic to id_X relative to A .

Proposition. If A is a deformation retract of X then $X \simeq A$.

2.2 Paths and path-homotopy

Definition (Path-homotopy). Two paths γ_0 and γ_1 are *path-homotopic* if they are homotopic relative to $\{0, 1\} \subseteq I$. In particular $\gamma_0(0) = \gamma_1(0)$, $\gamma_0(1) = \gamma_1(1)$. If F is a homotopy from γ_0 to γ_1 ,

$$F(-, 0) = \gamma_0(0), F(-, 1) = \gamma_1(1)$$

F is a family of paths connecting $\gamma_0(0)$ and $\gamma_0(1)$

Proposition. Path-homotopy is an equivalence relation on the set of paths in (a topological space) X .

Definition (Based loop). A *based loop* at $x_0 \in X$ is a path $\gamma : I \rightarrow X$ such that $\gamma(0) = \gamma(1) = x_0$.

Definition (Fundamental group of a space). The *fundamental group* of X at x_0 is the set (group)

$$\{[\gamma] \mid \gamma \text{ is a loop based at } x_0\}$$

which is denoted by $\Pi_1(X, x_0)$.

Definition (n^{th} homotopy group). The n^{th} homotopy group of a space X at x_0 is the set (group)

$$\pi_n(X, x_0) = \{[f : I^n \rightarrow X \mid f(\partial I^n) \rightarrow x_0]\}$$

Definition. A loop based at x_0 is null-homotopic if it is path-homotopic to a constant path.

Definition (Free homotopy). If γ_0 and γ_1 are based loops (not necessarily at the same point), then γ_0 and γ_1 are *freely homotopic* if they are homotopic through based loops, so if F is a free homotopy between γ_0 and γ_1 , then,

$$\begin{aligned} F(x_0) &= \gamma_0, F(x, 1) = \gamma_1 \\ F(0, t) &= F(1, t) \text{ for all } t \end{aligned}$$

Proposition. Free homotopy is an equivalence relation on the set of based loops in (a topological space) X .

Definition. A based loop *bounds a disk* if the induced map

$$\bar{\gamma} : [0, 1]/0=1 \cong S^1 \subseteq D^2$$

extends to a continuous function $D^2 \rightarrow X$.

Lemma. The following are equivalent

1. γ bounds a disk.
2. γ is null-homotopic.
3. γ is freely homotopic to a constant path.

3 Covering spaces

Definition (Covering map). A map $p : X' \rightarrow X$ is a *covering map* if $\forall x \in X$, there exists U and open neighbourhood of x , and a discrete set Δ and a homeomorphism $h_U : U \times \Delta \rightarrow p^{-1}(U)$ such that

$$p \circ h_u = \pi_U : U \times \Delta \rightarrow U$$

and such a neighbourhood U is called a *covering neighbourhood*.

Definition (Lift). Let $f : Y \rightarrow X$ and $g : Z \rightarrow X$ be two maps,

$$\begin{array}{ccc} & & Z \\ & \nearrow \tilde{f} & \downarrow g \\ Y & \xrightarrow{f} & X \end{array}$$

a lift of f is a map $\tilde{f} : Y \rightarrow Z$ such that

$$g \circ \tilde{f} = f$$

Lemma (Path/Homotopy lifting lemma). Let $p : X' \rightarrow X$ be a covering map and $f : I^n \rightarrow X$ a continuous map. Then for any $x' \in p^{-1}(f(U))$, there exists a unique lift \tilde{f} of f to X' , where $\tilde{f}(0) = x'$. Similarly, a homotopy can be lifted to a homotopy.