



ANALYSIS I, TUTORIAL 3 / Real-valued Function with a real variable Differentiability

Exercise 1. Provide the derivative of the following functions and exhibit their domain.

$$f(x) = \frac{1}{\sqrt[3]{x^2}} - \frac{1}{\sqrt{x^3}}, \quad g(x) = \sin(\cos(3x)), \quad h(x) = 2^{\ln(x)} + 3^x.$$

Exercise 2. Provide the n -th derivative of the following functions

$$f(x) = e^{2x}, \quad g(x) = \frac{1}{1+x}, \quad h(x) = \sin(x).$$

Exercise 3. Let f be a real-valued function with a real variable defined by

$$\forall x \in \mathbb{R} : \quad f(x) = \frac{x}{1+|x|}.$$

- Provide the domain, the parity and the derivative of f .
- Provide the limit of f at its domain's boundary.
- Find the largest sets $\mathcal{A}, \mathcal{B} \subset \mathbb{R}$ such as the maps g defined as

$$\begin{aligned} g : \mathcal{A} &\longrightarrow \mathcal{B} \\ x &\longmapsto g(x) = f(x), \end{aligned}$$

be a bijective one; in this case, exhibit the inverse of g .

Exercise 4.

- Show that for any $x \in [-1, +1]$ we have $\arcsin(x) + \arccos(x) = \pi/2$.
- Show that for any $x \in [0, +1]$ we have $\arcsin(x) + \arcsin(\sqrt{1-x^2}) = \pi/2$.

Exercise 5. [The inverse of the hyperbolic sinus] Let f be the function defined over \mathbb{R} with value in \mathbb{R} and satisfies

$$f(x) = \operatorname{sh}(x) = \frac{e^x - e^{-x}}{2}.$$

Show that f has an inverse, denoted it arcsh , and provide the derivative of f^{-1} .

Exercise 6. [The inverse of the hyperbolic consinus] Let f be the function defined over \mathbb{R} with value in \mathbb{R}_+^* and satisfies

$$f(x) = \operatorname{ch}(x) = \frac{e^x + e^{-x}}{2}.$$

Show that f has an inverse, denoted it arcch , and provide the derivative of f^{-1} .

Exercise 7. [The inverse of the hyperbolic tangent] Let f be the function defined over \mathbb{R} with value in $] -1, 1[$ and satisfies

$$f(x) = \operatorname{tanh}(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

Show that f has an inverse, denoted it $\operatorname{arctanh}$, and provide the derivative of f^{-1} .

Exercise 8. Study the differentiability of the following functions

$$f(x) = x|x|, \quad g(x) = \frac{1}{1+|x|}, \quad h(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Exercise 9. Let $a, b, c \in \mathbb{R}$, we define the function f as

$$f(x) = \begin{cases} \sqrt{x} & \text{if } x \in [0; 1], \\ ax^2 + bx + c & \text{if } x > 1. \end{cases}$$

Find a, b, c so that the function f be differentiable over \mathbb{R}_+^* .

Exercise 10.

- Show that the equation $\cos(x) - x = 0$ has a solution in $[0, 1]$.
- Show that the equation $x + e^x = 0$ has a unique solution in \mathbb{R} .
- Provide an example of function g from $]0, 1[$ to $]0, 1[$ without fixed point.

Exercise 11. Let $n \in \mathbb{N}^*$, we defined the function \mathcal{L}_n over $I = [0, 1]$ by

$$\forall x \in I : \quad \mathcal{L}_n(x) = x^n \sin(\pi x).$$

Show the existence of $x_n \in]0, 1[$ such that $\mathcal{L}_n'(x_n) = 0$, write $\mathcal{L}_n(x)$ in terms of $\mathcal{L}_n'(x)$, and calculate the limit of $\mathcal{L}_n(x_n)$ when n tends to $+\infty$.

Exercise 12. Let the function $f(x) = \text{Ln}(x)$ defined over \mathbb{R}_+^* with value in \mathbb{R} .

- Show that

$$\forall x \in \mathbb{R}_+^* : \quad \frac{1}{1+x} < f(x+1) - f(x) < \frac{1}{x}.$$

- Calculate the value of the following limits

$$\lim_{x \rightarrow +\infty} \sqrt{x} (\text{Ln}(x+1) - \text{Ln}(x)), \quad \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x}\right)^x.$$

Exercise 13. Let the function f defined over \mathbb{R} with value in \mathbb{R} by

$$f(x) = \begin{cases} \frac{3-x^2}{2} & \text{si } x < 1, \\ \frac{1}{x} & \text{si } x \geq 1. \end{cases}$$

Show that f is continuous and differentiable on \mathbb{R} and there exists $x_0 \in]0, 2[$ such that

$$2f^{(1)}(x_0) = f(2) - f(0).$$

Exercise 14. Show that for ever $x \in \mathbb{R}_+$ we have $x \leq e^x$ and prove that

$$\forall \alpha \in \mathbb{R}_+, \quad \exists C_\alpha \in \mathbb{R}_+^*, \quad \exists K_\alpha \in \mathbb{R}_+^*, \quad \forall x \in \mathbb{R}_+ : \quad C_\alpha x^\alpha \leq e^x, \quad K_\alpha x^\alpha \geq \text{Ln}(x).$$

Show that for any $s \in \mathbb{R}_+^*$ we have

$$\lim_{x \rightarrow +\infty} \frac{e^x}{x^s} = +\infty, \quad \lim_{x \rightarrow +\infty} \frac{x^s}{\text{Ln}(x)} = +\infty.$$

Exercise 15. Let n be an integer greater than or equal to two; we define the fuction f_n from \mathbb{R} to \mathbb{R} by

$$\forall x \in \mathbb{R} : \quad f_n(x) = x^n + x^{n-1} + x^2 + x - 1.$$

- Show that f_n has a unique root noted u_n in \mathbb{R}_+^* .
- Show that $u_n \in]0, 2/3[$ for every $n \in \mathbb{N}^*$.
- Show that the sequence $(u_n)_n$ is strictly increasing.
- Show that the sequence $(u_n)_n$ converges and calculate its limit.