

Knot Traces and Sliceness


Isaac M. Craig

Bryn Mawr College

MAA EPaDel Section Meeting
25 March 2019

A **framed knot** $K \subset S^3$ is a knot decorated with an integer n , called the **framing**.


A **framed knot** $K \subset S^3$ is a knot decorated with an integer n , called the **framing**.




A 0-framed knot (left) with the 0-framed push-off (right).


The framing induces a **push-off** K' with linking number $\text{lk}(K, K') = n$.

We want to use framed knots to build a 4-manifold by (essentially) attaching a disk to a knot in $S^3 = \partial B^4$. To visualize this, pretend we're down a dimension:






* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.




To a framed knot $K \subset S^3 = \partial B^4$, we associate a 4-manifold* $X(K)$ by attaching a framed 2-handle to B^4 along K according to its framing:


* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.




To a framed knot $K \subset S^3 = \partial B^4$, we associate a 4-manifold* $X(K)$ by attaching a framed 2-handle to B^4 along K according to its framing:




* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.




To a framed knot $K \subset S^3 = \partial B^4$, we associate a 4-manifold* $X(K)$ by attaching a framed 2-handle to B^4 along K according to its framing:



* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.




To a framed knot $K \subset S^3 = \partial B^4$, we associate a 4-manifold* $X(K)$ by attaching a framed 2-handle to B^4 along K according to its framing:




The associated 4-manifold to an n -framed knot K is called the **n -trace** of K .

* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.



To a framed knot $K \subset S^3 = \partial B^4$, we associate a 4-manifold* $X(K)$ by attaching a framed 2-handle to B^4 along K according to its framing:




The associated 4-manifold to an n -framed knot K is called the **n -trace** of K .

Question. Do properties of K correspond to properties of $X(K)$, and conversely?


* Recall, an n -manifold is a sufficiently nice topological space that locally looks n -dimensional.

A knot $K \subset S^3 = \partial B^4$ is **slice** if it bounds a smoothly embedded disk $D^2 \hookrightarrow B^4$.

A knot $K \subset S^3 = \partial B^4$ is **slice** if it bounds a smoothly embedded disk $D^2 \hookrightarrow B^4$.




A knot $K \subset S^3 = \partial B^4$ is **slice** if it bounds a smoothly embedded disk $D^2 \hookrightarrow B^4$.




Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

A knot $K \subset S^3 = \partial B^4$ is **slice** if it bounds a smoothly embedded disk $D^2 \hookrightarrow B^4$.



Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

Theorem. (Piccirillo 2018) The Conway knot is not slice.



The Conway Knot

Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

Theorem. (Piccirillo 2018) The Conway knot C is not slice.



Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

Theorem. (Piccirillo 2018) The Conway knot C is not slice.



Proof Technique. Construct knot C' with $X(C) \cong X(C')$. Show C' is not slice.


The Conway Knot

Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

Theorem. (Piccirillo 2018) The Conway knot C is not slice.



Proof Technique. Construct knot C' with $X(C) \cong X(C')$. Show C' is not slice.




The Conway Knot

Theorem. Let K and K' be knots with diffeomorphic traces $X(K) \cong X(K')$. Then K is slice if and only if K' is slice.

Theorem. (Piccirillo 2018) The Conway knot C is not slice.




Proof Technique. Construct knot C' with $X(C) \cong X(C')$. Show C' is not slice.



Remark. The construction of C' is known to work for unknotting $\# 1$ knots.


Knot Traces of Knots with Unknotting Number 1

A knot K with (positive) unknotting number 1 has a diagram:





Knot Traces of Knots with Unknotting Number 1



A knot K with (positive) unknotting number 1 has a diagram:



The associated knot K' then has the form:



New C and K'



Bibliography I



Lisa Piccirillo,

Shake genus and slice genus,

arXiv:1803.09834,, to appear in G&T (2018).