

# Knot Traces and Sliceness

Isaac M. Craig

Joint with Paul Melvin

Bryn Mawr College


GSTGC

30 March 2019

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:


# Knot Traces

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:




# Knot Traces

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:




# Knot Traces

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:




# Knot Traces

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:



The associated 4-manifold  $X(K)$  to an  $n$ -framed knot  $K$  is called its  **$n$ -trace**.

To a framed knot  $K \subset S^3 = \partial B^4$ , we associate a 4-manifold  $X(K)$  by attaching a framed 2-handle to  $B^4$  along  $K$  according to its framing:




The associated 4-manifold  $X(K)$  to an  $n$ -framed knot  $K$  is called its  **$n$ -trace**.

*Question.* Do properties of  $K$  correspond to properties of  $X(K)$ , and conversely?

# Slice Knots


A knot  $K \subset S^3 = \partial B^4$  is **slice** if it bounds a smoothly embedded disk  $D^2 \hookrightarrow B^4$ .






# Slice Knots

A knot  $K \subset S^3 = \partial B^4$  is **slice** if it bounds a smoothly embedded disk  $D^2 \hookrightarrow B^4$ .




A knot  $K \subset S^3 = \partial B^4$  is **slice** if it bounds a smoothly embedded disk  $D^2 \hookrightarrow B^4$ .



*Remark.* The core of the 2-handle and slice-disk form a smoothly embedded  $S^2$ .


A knot  $K \subset S^3 = \partial B^4$  is **slice** if it bounds a smoothly embedded disk  $D^2 \hookrightarrow B^4$ .



*Remark.* The core of the 2-handle and slice-disk form a smoothly embedded  $S^2$ .

**Theorem.** Let  $K$  and  $K'$  be knots with diffeomorphic traces  $X(K) \cong X(K')$ . Then  $K$  is slice if and only if  $K'$  is slice.


A knot  $K \subset S^3 = \partial B^4$  is **slice** if it bounds a smoothly embedded disk  $D^2 \hookrightarrow B^4$ .



*Remark.* The core of the 2-handle and slice-disk form a smoothly embedded  $S^2$ .

**Theorem.** Let  $K$  and  $K'$  be knots with diffeomorphic traces  $X(K) \cong X(K')$ . Then  $K$  is slice if and only if  $K'$  is slice.

**Theorem.** (Piccirillo [1]) The Conway knot is not slice.



**Theorem.** (Piccirillo [1]) The Conway knot  $C$  is not slice.


**Theorem.** (Piccirillo [1]) The Conway knot  $C$  is not slice.

**Proof Technique.** Construct knot  $C'$  with  $X(C) \cong X(C')$ . Show  $C'$  is not slice.

# The Conway Knot


**Theorem.** (Piccirillo [1]) The Conway knot  $C$  is not slice.

**Proof Technique.** Construct knot  $C'$  with  $X(C) \cong X(C')$ . Show  $C'$  is not slice.



**Theorem.** (Piccirillo [1]) The Conway knot  $C$  is not slice.

**Proof Technique.** Construct knot  $C'$  with  $X(C) \cong X(C')$ . Show  $C'$  is not slice.




*Remark.* This construction is known to work for unknotting number 1 knots.




# The Conway Knot

**Theorem.** (Piccirillo [1]) The Conway knot  $C$  is not slice.



**Proof Technique.** Construct knot  $C'$  with  $X(C) \cong X(C')$ . Show  $C'$  is not slice.



*Remark.* This construction is known to work for unknotting number 1 knots.



## New $C$ and $K'$



# Bibliography I



Lisa Piccirillo,  
*The Conway knot is not slice,*  
*arXiv:1808.02923*, (2018).