


Khovanov homology and exotic surfaces in the 4-ball

Isaac Sundberg

Max-Planck-Institut für Mathematik

PATCH - Penn



11 November 2022



Non-isotopic Seifert surfaces in B^4

Theorem (Hayden-Miller-Kim-Park-S. 22)



There are knots bounding pairs of non-isotopic Seifert surfaces, which remain non-isotopic when pushed into B^4 .



Non-isotopic Seifert surfaces in B^4

Theorem (Hayden-Miller-Kim-Park-S. 22)


There are knots bounding pairs of non-isotopic Seifert surfaces, which remain non-isotopic when pushed into B^4 .



Non-isotopic Seifert surfaces in B^4

Theorem (Hayden-Miller-Kim-Park-S. 22)

There are knots bounding pairs of non-isotopic Seifert surfaces, which remain non-isotopic when pushed into B^4 .




(All x labels)

Non-isotopic Seifert surfaces in B^4

Theorem (Hayden-Miller-Kim-Park-S. 22)

There are knots bounding pairs of non-isotopic Seifert surfaces, which remain non-isotopic when pushed into B^4 .




(All x labels)

Non-isotopic Seifert surfaces in B^4

Theorem (Hayden-Miller-Kim-Park-S. 22)

There are knots bounding pairs of non-isotopic Seifert surfaces, which remain non-isotopic when pushed into B^4 .



Thank You!

Thank you!