

Relative Khovanov-Jacobsson classes

Isaac Sundberg

Joint with Jonah Swann

Bryn Mawr College

AIM Research Community in 4-dimensional Topology:
Current Events Seminar

14 April 2021

Table of Contents

- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes
- 4 Obstructing Sliceness
- 5 Future work

Table of Contents

- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes
- 4 Obstructing Sliceness
- 5 Future work

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

- Objects: oriented links

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

- Objects: oriented links
- Morphisms: link cobordisms

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

- Objects: oriented links
- Morphisms: link cobordisms

The algebraic category we will get information from:

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

- Objects: oriented links
- Morphisms: link cobordisms

The algebraic category we will get information from:

- Objects: chain complexes (more generally, \mathbb{Z} -modules)

Categories

We will be using Khovanov homology to get algebraic information about the topological things we are interested in.

The topological category we are interested in:

- Objects: oriented links
- Morphisms: link cobordisms

The algebraic category we will get information from:


- Objects: chain complexes (more generally, \mathbb{Z} -modules)
- Morphisms: chain maps (more generally, \mathbb{Z} -linear maps)

Link Cobordisms

Definition. A **link cobordism** $\Sigma: L_0 \rightarrow L_1$ is a smooth, compact, oriented, properly embedded surface $\Sigma \subset \mathbb{R}^3 \times [0, 1]$ with boundary a pair $(i \in \{0, 1\})$ of oriented links $L_i = \Sigma \cap (\mathbb{R}^3 \times \{i\})$.


Link Cobordisms

Definition. A **link cobordism** $\Sigma: L_0 \rightarrow L_1$ is a smooth, compact, oriented, properly embedded surface $\Sigma \subset \mathbb{R}^3 \times [0, 1]$ with boundary a pair $(i \in \{0, 1\})$ of oriented links $L_i = \Sigma \cap (\mathbb{R}^3 \times \{i\})$.



Link Cobordisms


Definition. A **link cobordism** $\Sigma: L_0 \rightarrow L_1$ is a smooth, compact, oriented, properly embedded surface $\Sigma \subset \mathbb{R}^3 \times [0, 1]$ with boundary a pair $(i \in \{0, 1\})$ of oriented links $L_i = \Sigma \cap (\mathbb{R}^3 \times \{i\})$.



Examples: concordance, closed surfaces in B^4 ($\emptyset \rightarrow \emptyset$), slices ($\emptyset \rightarrow K$), Seifert surfaces ($\emptyset \rightarrow K$)

Link Cobordisms

Definition. A **link cobordism** $\Sigma: L_0 \rightarrow L_1$ is a smooth, compact, oriented, properly embedded surface $\Sigma \subset \mathbb{R}^3 \times [0, 1]$ with boundary a pair $(i \in \{0, 1\})$ of oriented links $L_i = \Sigma \cap (\mathbb{R}^3 \times \{i\})$.




Examples: concordance, closed surfaces in B^4 ($\emptyset \rightarrow \emptyset$), slices ($\emptyset \rightarrow K$), Seifert surfaces ($\emptyset \rightarrow K$)

Definition. A link cobordism $\Sigma: L_0 \rightarrow L_1$ can be represented as a **movie**: a finite sequence of diagrams $\{D_{t_i}\}_{i=0}^n$, called *frames*, with each successive pair related by an isotopy, Morse move, or Reidemeister move.

Link Cobordisms

Definition. A **link cobordism** $\Sigma: L_0 \rightarrow L_1$ is a smooth, compact, oriented, properly embedded surface $\Sigma \subset \mathbb{R}^3 \times [0, 1]$ with boundary a pair $(i \in \{0, 1\})$ of oriented links $L_i = \Sigma \cap (\mathbb{R}^3 \times \{i\})$.



Examples: concordance, closed surfaces in B^4 ($\emptyset \rightarrow \emptyset$), slices ($\emptyset \rightarrow K$), Seifert surfaces ($\emptyset \rightarrow K$)

Definition. A link cobordism $\Sigma: L_0 \rightarrow L_1$ can be represented as a **movie**: a finite sequence of diagrams $\{D_{t_i}\}_{i=0}^n$, called *frames*, with each successive pair related by an isotopy, Morse move, or Reidemeister move.



Background
○○●○○○

Definition
○○○

Isotopy Classes
○○○○○○○○○

Sliceness
○○○○

Future work
○○○○○

Khovanov homology

Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Properties:

Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Properties:

- elements of $\mathcal{CKh}(D)$ are *labeled smoothings* of D




Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Properties:

- elements of $\mathcal{CKh}(D)$ are *labeled smoothings* of D




Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Properties:

- elements of $\mathcal{CKh}(D)$ are *labeled smoothings* of D



- the chain complex can be bigraded $\mathcal{CKh}^{h,q}(D)$


Khovanov homology

Theorem (Khovanov '00)

A diagram D of an oriented link L induces a chain complex $\mathcal{CKh}(D)$ with homology $\text{Kh}(D)$, called the Khovanov homology.

Properties:

- elements of $\mathcal{CKh}(D)$ are *labeled smoothings* of D



- the chain complex can be bigraded $\mathcal{CKh}^{h,q}(D)$
- different diagrams have isomorphic Khovanov homology, so $\text{Kh}(L)$ means choose a diagram D for L and consider $\text{Kh}(D)$

Background
○○○○●○○

Definition
○○○

Isotopy Classes
○○○○○○○○○○

Sliceness
○○○○

Future work
○○○○○

Maps on Khovanov homology

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$



Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$



Maps on Khovanov homology


Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$



Maps on Khovanov homology


Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$



Maps on Khovanov homology


Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

- Define chain maps for isotopies, Morse moves, and Reidemeister moves
- Adjacent frames $D_{t_i} \rightarrow D_{t_{i+1}}$ induce one of these maps
- Compose these chain maps to produce $\mathcal{CKh}(\Sigma)$



Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

- This map has a bigrading

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}^{h,q}(D_0) \rightarrow \mathcal{CKh}^{h,q+\chi(\Sigma)}(D_1)$$

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

- This map has a bigrading

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}^{h,q}(D_0) \rightarrow \mathcal{CKh}^{h,q+\chi(\Sigma)}(D_1)$$

- This map is invariant, up to sign, under boundary-preserving isotopy of Σ .

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

- This map has a bigrading

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}^{h,q}(D_0) \rightarrow \mathcal{CKh}^{h,q+\chi(\Sigma)}(D_1)$$

- This map is invariant, up to sign, under boundary-preserving isotopy of Σ .
In other words, if $\Sigma': L_0 \rightarrow L_1$ satisfies $\Sigma \cong \Sigma'$ (rel $L_{0,1}$)

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

- This map has a bigrading

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}^{h,q}(D_0) \rightarrow \mathcal{CKh}^{h,q+\chi(\Sigma)}(D_1)$$

- This map is invariant, up to sign, under boundary-preserving isotopy of Σ .
In other words, if $\Sigma': L_0 \rightarrow L_1$ satisfies $\Sigma \cong \Sigma'$ (rel $L_{0,1}$), then

$$\mathcal{CKh}(\Sigma'): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

satisfies $\mathcal{CKh}(\Sigma) \simeq \pm \mathcal{CKh}(\Sigma')$.

Maps on Khovanov homology

Theorem (Khovanov '00)

A movie $\{D_{t_i}\}_{i=0}^n$ of a link cobordism $\Sigma: L_0 \rightarrow L_1$ induces a chain map

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

with induced homomorphism $\text{Kh}(\Sigma)$ on homology.

Properties:

- This map has a bigrading

$$\mathcal{CKh}(\Sigma): \mathcal{CKh}^{h,q}(D_0) \rightarrow \mathcal{CKh}^{h,q+\chi(\Sigma)}(D_1)$$

- This map is invariant, up to sign, under boundary-preserving isotopy of Σ .
In other words, if $\Sigma': L_0 \rightarrow L_1$ satisfies $\Sigma \cong \Sigma'$ (rel $L_{0,1}$), then

$$\mathcal{CKh}(\Sigma'): \mathcal{CKh}(D_0) \rightarrow \mathcal{CKh}(D_1)$$

satisfies $\mathcal{CKh}(\Sigma) \simeq \pm \mathcal{CKh}(\Sigma')$. Or equivalently $\text{Kh}(\Sigma) = \pm \text{Kh}(\Sigma')$.

Example

Let's look at the simplest example:

Example

Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$.

Example


Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$.

Example

Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:


Example

Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



Example


Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



- a. Choose your favorite movie for $\Sigma: \emptyset \rightarrow \emptyset$

Example


Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow \emptyset$
- Construct induced map $\text{Kh}(\Sigma)$ on Khovanov homology.

Example


Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow \emptyset$
- Construct induced map $\text{Kh}(\Sigma)$ on Khovanov homology. Note that $\text{Kh}(\Sigma)$ is determined by $\text{Kh}(\Sigma)(1)$.

Example


Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow \emptyset$
- Construct induced map $\text{Kh}(\Sigma)$ on Khovanov homology. Note that $\text{Kh}(\Sigma)$ is determined by $\text{Kh}(\Sigma)(1)$.
- $\text{Kh}(\Sigma)$ is invariant, up to sign, under boundary-preserving isotopy of Σ

Example

Let's look at the simplest example: consider a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$. Note that $\text{Kh}(\emptyset) = \mathbb{Z}$. We then have the following construction:



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow \emptyset$
- Construct induced map $\text{Kh}(\Sigma)$ on Khovanov homology. Note that $\text{Kh}(\Sigma)$ is determined by $\text{Kh}(\Sigma)(1)$.
- $\text{Kh}(\Sigma)$ is invariant, up to sign, under boundary-preserving isotopy of Σ , so the element

$$\text{KJ}_\Sigma := |\text{Kh}(\Sigma)(1)|$$

is also an invariant of the (ambient) isotopy class of Σ .

Khovanov-Jacobsson classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$, the element

$$KJ_{\Sigma} := |\mathrm{Kh}(\Sigma)(1)| \in \mathbb{Z}$$

is called the *Khovanov-Jacobsson class* of Σ .

Khovanov-Jacobsson classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$, the element

$$KJ_{\Sigma} := |\mathrm{Kh}(\Sigma)(1)| \in \mathbb{Z}$$

is called the *Khovanov-Jacobsson class* of Σ .

The Khovanov-Jacobsson class of a link cobordism Σ is an invariant of the (ambient) isotopy class of Σ .

Khovanov-Jacobsson classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$, the element

$$KJ_{\Sigma} := |\text{Kh}(\Sigma)(1)| \in \mathbb{Z}$$

is called the *Khovanov-Jacobsson class* of Σ .

The Khovanov-Jacobsson class of a link cobordism Σ is an invariant of the (ambient) isotopy class of Σ .

Question. Does KJ_{Σ} distinguish any surfaces up to isotopy?

Khovanov-Jacobsson classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow \emptyset$, the element

$$KJ_{\Sigma} := |\text{Kh}(\Sigma)(1)| \in \mathbb{Z}$$

is called the *Khovanov-Jacobsson class* of Σ .

The Khovanov-Jacobsson class of a link cobordism Σ is an invariant of the (ambient) isotopy class of Σ .

Question. Does KJ_{Σ} distinguish any surfaces up to isotopy?

Theorem (Rasmussen '05, Tanaka '05)

Khovanov-Jacobsson classes of connected Σ are determined by genus:

- if $g(\Sigma) = 1$, then $KJ_{\Sigma} = 2$
- if $g(\Sigma) \neq 1$, then $KJ_{\Sigma} = 0$

Table of Contents


- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes
- 4 Obstructing Sliceness
- 5 Future work

Construction

Idea: follow the same procedure for a link cobordism $\Sigma: \emptyset \rightarrow L$, that is, a surface with boundary.

Construction


Idea: follow the same procedure for a link cobordism $\Sigma: \emptyset \rightarrow L$, that is, a surface with boundary.



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow L$.

Construction


Idea: follow the same procedure for a link cobordism $\Sigma: \emptyset \rightarrow L$, that is, a surface with boundary.



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow L$.
- Construct induced map $\text{Kh}(\Sigma): \mathbb{Z} \rightarrow \text{Kh}(L)$, which is determined by $\text{Kh}(\Sigma)(1)$.

Construction

Idea: follow the same procedure for a link cobordism $\Sigma: \emptyset \rightarrow L$, that is, a surface with boundary.



- Choose your favorite movie for $\Sigma: \emptyset \rightarrow L$.
- Construct induced map $\text{Kh}(\Sigma): \mathbb{Z} \rightarrow \text{Kh}(L)$, which is determined by $\text{Kh}(\Sigma)(1)$.
- Produce invariant KJ_Σ of the boundary-preserving isotopy class of Σ .

Relative Khovanov-Jacobsson Classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow L$, the element

$$KJ_{\Sigma} := |\mathrm{Kh}(\Sigma)(1)| \in \mathrm{Kh}(L)$$

is called the *relative Khovanov-Jacobsson class* of Σ .

Relative Khovanov-Jacobsson Classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow L$, the element

$$KJ_{\Sigma} := |\mathrm{Kh}(\Sigma)(1)| \in \mathrm{Kh}(L)$$

is called the *relative Khovanov-Jacobsson class* of Σ .

The relative Khovanov-Jacobsson class of a link cobordism Σ is an invariant of the **boundary-preserving** isotopy class of Σ .

Relative Khovanov-Jacobsson Classes

Definition

For a link cobordism $\Sigma: \emptyset \rightarrow L$, the element

$$KJ_{\Sigma} := |\mathrm{Kh}(\Sigma)(1)| \in \mathrm{Kh}(L)$$

is called the *relative Khovanov-Jacobsson class* of Σ .

The relative Khovanov-Jacobsson class of a link cobordism Σ is an invariant of the **boundary-preserving** isotopy class of Σ .

Will this relative invariant have the same fate as the absolute ($L = \emptyset$) case?

Table of Contents

- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes**
- 4 Obstructing Sliceness
- 5 Future work

Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

If so, then the relative classes have more information than the absolute classes.

Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

If so, then the relative classes have more information than the absolute classes.

Plan:


Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

If so, then the relative classes have more information than the absolute classes.

Plan:




Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

If so, then the relative classes have more information than the absolute classes.

Plan:




Obstructing Boundary-Preserving Isotopy Classes

Obstruction: if a pair of link cobordisms $\Sigma, \Sigma' : \emptyset \rightarrow L$ have relative classes $KJ_{\Sigma'} \neq KJ_{\Sigma}$, then $\Sigma \not\approx \Sigma' \text{ rel } L$.

Question. Can we find $\Sigma, \Sigma' : \emptyset \rightarrow L$ having $KJ_{\Sigma} \neq KJ_{\Sigma'}$?

If so, then the relative classes have more information than the absolute classes.


Plan:



Example: two slices for 9_{46}

Theorem (Swann '10)


The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.



Example: two slices for 9_{46}

Theorem (Swann '10)


The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.



Example: two slices for 9_{46}

Theorem (Swann '10)


The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.



Example: two slices for 9_{46}

Theorem (Swann '10)


The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.



Example: two slices for 9_{46}

Theorem (Swann '10)


The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.




Example: two slices for 9_{46}

Theorem (Swann '10)

The two slices below for 9_{46} have distinct relative Khovanov-Jacobsson classes.



These surfaces Σ and Σ' have distinct relative classes.

Calculation for 9_{46} 

Example: two slices for \mathfrak{g}_1

Theorem (S. '21)


The two slices below for \mathfrak{g}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{b}_1

Theorem (S. '21)


The two slices below for \mathfrak{b}_1 have distinct relative Khovanov-Jacobsson classes.

 Σ  Σ'

Example: two slices for \mathfrak{g}_1

Theorem (S. '21)


The two slices below for \mathfrak{g}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{g}_1

Theorem (S. '21)


The two slices below for \mathfrak{g}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{g}_1

Theorem (S. '21)


The two slices below for \mathfrak{g}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{G}_1

Theorem (S. '21)


The two slices below for \mathfrak{G}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{G}_1

Theorem (S. '21)


The two slices below for \mathfrak{G}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{G}_1

Theorem (S. '21)


The two slices below for \mathfrak{G}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{G}_1

Theorem (S. '21)


The two slices below for \mathfrak{G}_1 have distinct relative Khovanov-Jacobsson classes.



Example: two slices for \mathfrak{b}_1

Theorem (S. '21)

The two slices below for \mathfrak{b}_1 have distinct relative Khovanov-Jacobsson classes.



These surfaces Σ and Σ' also have distinct relative classes.

Good news!

This is good news!

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

On the other hand, this isn't anything new (we have known these slices are distinct, using different techniques).

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

On the other hand, this isn't anything new (we have known these slices are distinct, using different techniques).

There are a few follow-up questions we could ask:

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

On the other hand, this isn't anything new (we have known these slices are distinct, using different techniques).

There are a few follow-up questions we could ask:

- Can we find knots with more than 2 slices?

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

On the other hand, this isn't anything new (we have known these slices are distinct, using different techniques).

There are a few follow-up questions we could ask:

- Can we find knots with more than 2 slices?
- If so, can we find *prime* knots with more than 2 slices?

Good news!

This is good news! Relative Khovanov-Jacobsson classes can be used to distinguish link cobordisms $\emptyset \rightarrow L$.

On the other hand, this isn't anything new (we have known these slices are distinct, using different techniques).

There are a few follow-up questions we could ask:

- Can we find knots with more than 2 slices?
- If so, can we find *prime* knots with more than 2 slices?
- Can we distinguish other slices? Are there any other known/unknown examples to check?

Knots with many slices

Question: Can we find knots with more than 2 slices?

Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices.


Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices. Slices are obtained by choosing one of the band moves for each sail of the windmill (or boundary connect summing the slices).




Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices. Slices are obtained by choosing one of the band moves for each sail of the windmill (or boundary connect summing the slices).




Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices. Slices are obtained by choosing one of the band moves for each sail of the windmill (or boundary connect summing the slices).




Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices. Slices are obtained by choosing one of the band moves for each sail of the windmill (or boundary connect summing the slices).




Knots with many slices

Question: Can we find knots with more than 2 slices?

Theorem (S.-Swann '21)

There are knots with an arbitrarily large number of slices having distinct relative Khovanov-Jacobsson classes.

The knot $\#_k(9_{46})$ has 2^k slices. Slices are obtained by choosing one of the band moves for each sail of the windmill (or boundary connect summing the slices).



This was also shown in [4] through different techniques.

Prime knots with many slices

Question: can we find *prime* knots with more than 2 slices?

Prime knots with many slices

Question: can we find *prime* knots with more than 2 slices?

Theorem (S. '21)


There are prime knots with an arbitrarily large number of slices.

Prime knots with many slices

Question: can we find *prime* knots with more than 2 slices?

Theorem (S. '21)

There are prime knots with an arbitrarily large number of slices.




Prime knots with many slices

Question: can we find *prime* knots with more than 2 slices?

Theorem (S. '21)

There are prime knots with an arbitrarily large number of slices.




Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$



Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S



Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S



Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S
- Then Σ and $\Sigma \# S$ are (generally) not isotopic rel K .



Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S
- Then Σ and $\Sigma \# S$ are (generally) not isotopic rel K .



Definition. A surface Σ is *locally knotted* if $\Sigma = \Sigma' \# S$ for some surface Σ' .

Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S
- Then Σ and $\Sigma \# S$ are (generally) not isotopic rel K .



Definition. A surface Σ is *locally knotted* if $\Sigma = \Sigma' \# S$ for some surface Σ' .

Theorem (Swann '10)

Relative Khovanov-Jacobsson classes do not detect local knottedness.

Local knottedness

In general, it is (perhaps too) easy to change isotopy classes:

- Given $\Sigma: \emptyset \rightarrow L$
- Choose your favorite knotted 2-sphere S
- Then Σ and $\Sigma \# S$ are (generally) not isotopic rel K .



Definition. A surface Σ is *locally knotted* if $\Sigma = \Sigma' \# S$ for some surface Σ' .

Theorem (Swann '10)

Relative Khovanov-Jacobsson classes do not detect local knottedness.


That is, for a surface Σ and knotted 2-sphere S we have $KJ_{\Sigma} = KJ_{\Sigma \# S}$

Table of Contents

- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes
- 4 Obstructing Sliceness**
- 5 Future work


A note on ribbon disks

Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .



A note on ribbon disks


Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .



\therefore ribbon concordances induces injections on Khovanov homology

A note on ribbon disks


Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .



\therefore ribbon concordances induces injections on Khovanov homology

A note on ribbon disks


Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .



\therefore ribbon concordances induces injections on Khovanov homology

A note on ribbon disks

Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .




\therefore ribbon concordances induces injections on Khovanov homology

$\therefore KJ_\Sigma$ must be nontrivial.

A note on ribbon disks

Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .




\therefore ribbon concordances induces injections on Khovanov homology

$\therefore KJ_{\Sigma}$ must be nontrivial.

We might (optimistically) hope that relative Khovanov-Jacobsson classes can disprove the Slice-Ribbon Conjecture:

A note on ribbon disks

Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .




- ∴ ribbon concordances induces injections on Khovanov homology
- ∴ KJ_{Σ} must be nontrivial.

We might (optimistically) hope that relative Khovanov-Jacobsson classes can disprove the Slice-Ribbon Conjecture:

- find a slice Σ for a knot with $KJ_{\Sigma} = 0$
- then Σ cannot be ribbon

A note on ribbon disks

Suppose we have a ribbon disk $\Sigma: \emptyset \rightarrow K$ for some knot K .



- ∴ ribbon concordances induces injections on Khovanov homology
- ∴ KJ_{Σ} must be nontrivial.

We might (optimistically) hope that relative Khovanov-Jacobsson classes can disprove the Slice-Ribbon Conjecture:

- find a slice Σ for a knot with $KJ_{\Sigma} = 0$
- then Σ cannot be ribbon

Unfortunately, none can exist...

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

In particular, slices have nontrivial relative Khovanov-Jacobson classes.

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

In particular, slices have nontrivial relative Khovanov-Jacobsson classes.

Although the relative classes do not determine slice vs ribbon questions, they can be used to obstruct sliceness:

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

In particular, slices have nontrivial relative Khovanov-Jacobson classes.

Although the relative classes do not determine slice vs ribbon questions, they can be used to obstruct sliceness:

- Given a knot K with unknown sliceness

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

In particular, slices have nontrivial relative Khovanov-Jacobsson classes.

Although the relative classes do not determine slice vs ribbon questions, they can be used to obstruct sliceness:

- Given a knot K with unknown sliceness
- Find a link cobordism $\Sigma: \emptyset \rightarrow K$ with $g(\Sigma) = 1$

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.

In particular, slices have nontrivial relative Khovanov-Jacobsson classes.

Although the relative classes do not determine slice vs ribbon questions, they can be used to obstruct sliceness:

- Given a knot K with unknown sliceness
- Find a link cobordism $\Sigma: \emptyset \rightarrow K$ with $g(\Sigma) = 1$
- Show $KJ_\Sigma \neq 0$

Obstructing Sliceness

Theorem (Swann '10)

If K is slice, and $\Sigma: \emptyset \rightarrow K$ is a surface with $g(\Sigma) \leq 1$, then KJ_Σ is nontrivial.


In particular, slices have nontrivial relative Khovanov-Jacobsson classes.

Although the relative classes do not determine slice vs ribbon questions, they can be used to obstruct sliceness:

- Given a knot K with unknown sliceness
- Find a link cobordism $\Sigma: \emptyset \rightarrow K$ with $g(\Sigma) = 1$
- Show $KJ_\Sigma \neq 0$

Note: there are classes of knots with 4-ball genus at most 1 (e.g. Whitehead doubles, unknotting number 1 knots)

Example: Three-stranded Pretzel Knots




$$q_{146} = P(3, -3, 3)$$



$$P(3, 3, 3)$$


Example: Three-stranded Pretzel Knots

 $9_{46} = P(3, -3, 3)$  $P(3, 3, 3)$

Corollary (Swann '10)

For $p, q, r \geq 3$ and odd, the pretzel knot $P(p, q, r)$ is not slice.

Example: Three-stranded Pretzel Knots


 $q_{146} = P(3, -3, 3)$  $P(3, 3, 3)$

Corollary (Swann '10)

For $p, q, r \geq 3$ and odd, the pretzel knot $P(p, q, r)$ is not slice.

Proof idea: relative class of Seifert surface is 0.

Example: Three-stranded Pretzel Knots

 $9_{46} = P(3, -3, 3)$  $P(3, 3, 3)$

Corollary (Swann '10)


For $p, q, r \geq 3$ and odd, the pretzel knot $P(p, q, r)$ is not slice.

Proof idea: relative class of Seifert surface is 0.

Corollary (Swann '10)

For $p, q \leq -3$ and odd, the pretzel knot $P(p, q, 1)$ is not slice.

Example: Three-stranded Pretzel Knots

 $q_{46} = P(3, -3, 3)$  $P(3, 3, 3)$

Corollary (Swann '10)

For $p, q, r \geq 3$ and odd, the pretzel knot $P(p, q, r)$ is not slice.

Proof idea: relative class of Seifert surface is 0.

Corollary (Swann '10)

For $p, q \leq -3$ and odd, the pretzel knot $P(p, q, 1)$ is not slice.

Proof idea: relative class of Seifert surface is trivial (the cycle representing KJ_Σ is a boundary)

Table of Contents

- 1 Background
- 2 Defining relative Khovanov-Jacobsson classes
- 3 Obstructing Boundary-Preserving Isotopy Classes
- 4 Obstructing Sliceness
- 5 Future work

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero?

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

- General work done by A. Elliott ('09) and Swann ('10)

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

- General work done by A. Elliott ('09) and Swann ('10)
- Can get lucky (if homology in bigrading containing KJ_{Σ} is zero)

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

- General work done by A. Elliott ('09) and Swann ('10)
- Can get lucky (if homology in bigrading containing KJ_{Σ} is zero)
- Can map to a "nicer" knot by a cobordism and hope nontriviality is easier to check there

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

- General work done by A. Elliott ('09) and Swann ('10)
- Can get lucky (if homology in bigrading containing KJ_{Σ} is zero)
- Can map to a "nicer" knot by a cobordism and hope nontriviality is easier to check there

Advertisement: I wrote code that determines nontriviality of explicit classes from Khovanov homology (works for knots up to 12 crossings, and occasionally others).

Determining nontriviality of Khovanov classes

When is a class in Khovanov homology zero? This is hard to answer...

- General work done by A. Elliott ('09) and Swann ('10)
- Can get lucky (if homology in bigrading containing KJ_{Σ} is zero)
- Can map to a "nicer" knot by a cobordism and hope nontriviality is easier to check there

Advertisement: I wrote code that determines nontriviality of explicit classes from Khovanov homology (works for knots up to 12 crossings, and occasionally others).

Email me at icraig@brynmawr.edu

Relations to other invariants

Is there a relation to the s -invariant?

Relations to other invariants

Is there a relation to the s -invariant?

Conjecture

Relative Khovanov-Jacobsson classes do not detect any sliceness not already detected by the s -invariant.

Relations to other invariants

Is there a relation to the s -invariant?

Conjecture

Relative Khovanov-Jacobsson classes do not detect any sliceness not already detected by the s -invariant.

i.e. if $s(K) = 0$ for a knot K , then $KJ_{\Sigma} \neq 0$ for any link cobordism $\Sigma: \emptyset \rightarrow K$.

Ideally, this is false.

Relations to other invariants

Is there a relation to the s -invariant?

Conjecture

Relative Khovanov-Jacobsson classes do not detect any sliceness not already detected by the s -invariant.

i.e. if $s(K) = 0$ for a knot K , then $KJ_{\Sigma} \neq 0$ for any link cobordism $\Sigma: \emptyset \rightarrow K$.

Ideally, this is false.

Is there a relation to the invariant $t_{\Sigma,P}$ [2] from knot Floer homology?

Relations to other invariants

Is there a relation to the s -invariant?

Conjecture

Relative Khovanov-Jacobsson classes do not detect any sliceness not already detected by the s -invariant.

i.e. if $s(K) = 0$ for a knot K , then $\text{KJ}_\Sigma \neq 0$ for any link cobordism $\Sigma: \emptyset \rightarrow K$.

Ideally, this is false.

Is there a relation to the invariant $t_{\Sigma,P}$ [2] from knot Floer homology?

Conjecture

The Dowker spectral sequence relates KJ_Σ to the invariant $t_{\Sigma,P}$.

Relations to other invariants

Is there a relation to the s -invariant?

Conjecture

Relative Khovanov-Jacobsson classes do not detect any sliceness not already detected by the s -invariant.

i.e. if $s(K) = 0$ for a knot K , then $KJ_\Sigma \neq 0$ for any link cobordism $\Sigma: \emptyset \rightarrow K$.

Ideally, this is false.

Is there a relation to the invariant $t_{\Sigma,P}$ [2] from knot Floer homology?

Conjecture






The Dowker spectral sequence relates KJ_Σ to the invariant $t_{\Sigma,P}$.

If true, relative Khovanov-Jacobsson classes should be able to prove results similar to those from [3] about slice disks obtained from deform-spinning knots.

Thank You!

Thank you!

Bibliography I

-  Andrew Elliott, Graphical methods establishing nontriviality of state cycle Khovanov homology classes, *arXiv:0907.0396*, 2009
-  András Juhász and Marco Marengon, Concordance maps in knot Floer homology, *Geom. Topol.*,20(6):3623–3673, 2016. MR359035
-  András Juhász and Ian Zemke, Distinguishing slice disks using knot floer homology, *Seceta Math. (N.S.)*,20(1), 2020. Paper No. 5, 18 pp. MR404515
-  Allison N. Miller and Mark Powell, Stabilization distance between surfaces, *Enseign. Math.*,**65**:397-440, 2020. MR4113047
-  Jonah Swann, Relative Khovanov-Jacobsson classes of spanning surfaces, *Ph.D. Thesis, Bryn Mawr College*, 2010