

## Homework Assignment #2

Due: Feb 26

Student: Shuo Yang

1. Algorithm**Function** *FindClosePair*( $A[1 : n]$ )
$$\text{distance} := \text{ComputeCloseDistance}(A[1 : n])$$

$$(x, y) := \text{ComputeClosePair}(A[1 : n], \text{distance})$$
**Function** *ComputeCloseDistance*( $A[1 : n]$ )

Linearly scan the input array  $A[1 : n]$  from left to right, find the minimum and maximum numbers, put them into variables  $\text{min}$  and  $\text{max}$  respectively, and let

$$\text{distance} := \lfloor (\text{max} - \text{min}) / (n - 1) \rfloor.$$
**Function** *ComputeClosePair*( $A[1 : n], \text{distance}$ )

(1) Linearly scan the input array  $A$  from left to right, find the minimum and maximum numbers, put them into variables  $\text{min}$  and  $\text{max}$  respectively.

(2) Divide the  $n$  elements into  $\lfloor n/5 \rfloor$  groups of 5 elements, and  $\leq 1$  group of  $< 5$  elements.

(3) Find the median of each group using selection sort.

(4) Recursively find the median  $\text{med}$  of the  $\lfloor n/5 \rfloor$  medians found in step (3).

(5) Partition the input array  $A$  around  $\text{med}$  from step (4) into two subarrays  $A_{\text{low}}$  and  $A_{\text{high}}$  such that  $A_{\text{low}}$  contains all elements  $\leq \text{med}$  and  $A_{\text{high}}$  contains all elements  $\geq \text{med}$ .

(6) **if**  $\text{med} - \text{min} \leq \text{distance}$

**return** ( $\text{med}, \text{min}$ )

**if**  $\text{max} - \text{med} \leq \text{distance}$

**return** ( $\text{max}, \text{med}$ )

**if**  $n$  is odd

**if**  $\text{max} - \text{med} \geq \text{med} - \text{min}$

$(x, y) := \text{ComputeClosePair}(A_{\text{low}}, \text{distance})$

**else**

$(x, y) := \text{ComputeClosePair}(A_{\text{high}}, \text{distance})$

**else** //  $n$  is even

**if**  $\text{max} - \text{med} - \text{distance} \geq \text{med} - \text{min}$  //  $\text{med}$  is lower median

$(x, y) := \text{ComputeClosePair}(A_{\text{low}}, \text{distance})$

**else**

$(x, y) := \text{ComputeClosePair}(A_{\text{high}}, \text{distance})$

Correctness

**Lemma1:** Given an unsorted array  $A$  of  $n$  distinct numbers, a close pair always exists.

*Proof.* Prove by contradiction, that is, no such a close pair exists, this means for any pair  $(x, y)$  where  $x > y$ , we have:

$$x - y > \frac{1}{n-1}(\text{max} - \text{min}) \quad (1)$$

Suppose we order numbers in the array  $A$  in ascending order as a sequence:  $a_1, a_2, \dots, a_n$  where  $a_i > a_j$  if  $i > j$ . Let  $\text{distance} = \frac{1}{n-1}(\text{max} - \text{min})$ , according to the assumption, we have:

$$\begin{aligned}
a_n - a_{n-1} &> distance \\
a_{n-1} - a_{n-2} &> distance \\
&\dots \\
a_3 - a_2 &> distance \\
a_2 - a_1 &> distance
\end{aligned} \tag{2}$$

Summing the above equations together produces:

$$\begin{aligned}
a_n - a_1 &> distance \times (n - 1) \\
&= \frac{1}{n - 1} (max - min) \times (n - 1) \\
&= max - min
\end{aligned} \tag{3}$$

Since we have sorted the array  $A$  in ascending order, this means that  $a_n = max$  and  $a_1 = min$ , therefore the above equation says that  $max - min > max - min$ , this is clearly a contradiction, thus the lemma1 must be true.  $\square$

**Lemma2:** Given an unsorted array  $A$  of  $n$  distinct numbers partitioned around its median into two subarrays  $A_{low}$  and  $A_{high}$ .  $A_{low}$  contains all elements  $\leq$  median and  $A_{high}$  contains all elements  $\geq$  median. Let the maximum element be  $max$ , minimum element be  $min$  and median be  $med$ . A close pair must exist in  $A_{low}$  if  $max - med \geq med - min$  (when  $n$  is odd) or  $max - med - distance \geq med - min$  (when  $n$  is even), and in  $A_{high}$  otherwise.

*Proof.* Prove by contradiction. If  $max - med \geq med - min$  (when  $n$  is odd) or  $max - med - distance \geq med - min$  (when  $n$  is even), assume that  $A_{low}$  does not contain any close pairs. There are two cases to consider:

(a)  $n$  is odd.

In this case,  $A_{low}$  contains  $\frac{n-1}{2} + 1 = \frac{n+1}{2}$  elements, including the  $med$  itself. Applying the same method used in proving Lemma1, we sort  $A_{low}$  as:  $a_0, a_1, \dots, a_{(n+1)/2}$  in ascending order. According to the assumption, we must have,

$$\begin{aligned}
a_{\frac{n+1}{2}} - a_{\frac{n+1}{2}-1} &> distance \\
&\dots \\
a_3 - a_2 &> distance \\
a_2 - a_1 &> distance
\end{aligned} \tag{4}$$

Summing the above equations together produces:

$$\begin{aligned}
a_{\frac{n+1}{2}} - a_1 &> distance \times \left(\frac{n+1}{2} - 1\right) \\
&= \frac{1}{n-1} (max - min) \times \left(\frac{n+1}{2} - 1\right) \\
&= \frac{1}{n-1} (max - min) \times \frac{n-1}{2} \\
&= \frac{max - min}{2}
\end{aligned} \tag{5}$$

Since  $A_{low}$  is sorted in ascending order,  $a_{\frac{n+1}{2}} = med$  and  $a_1 = min$ . Thus,

$$\begin{aligned} med - min &> \frac{max - min}{2} \\ med &> \frac{max + min}{2} \end{aligned} \tag{6}$$

And because  $max - med \geq med - min$ , we have  $med \leq \frac{max+min}{2}$ . But we have just proved that  $med > \frac{max+min}{2}$ , clearly it is a contradiction. Thus, there must exist a close pair in  $A_{low}$ .

(b)  $n$  is even.

In this case, there are two medians, left median and right median. Assume that we pick the low median.  $A_{low}$  contains  $\frac{n}{2}$  elements, including the  $med$  itself. Applying the same method used in proving Lemma1, we sort  $A_{low}$  as:  $a_1, a_2, \dots, a_{n/2}$  in ascending order. According to the assumption, we must have,

$$\begin{aligned} a_{\frac{n}{2}} - a_{\frac{n}{2}-1} &> distance \\ &\dots \\ a_3 - a_2 &> distance \\ a_2 - a_1 &> distance \end{aligned} \tag{7}$$

Summing the above equations together produces:

$$\begin{aligned} a_{\frac{n}{2}} - a_1 &> distance \times \left(\frac{n}{2} - 1\right) \\ &= \frac{1}{n-1}(max - min) \times \left(\frac{n}{2} - 1\right) \\ &= \frac{max - min}{2} \times \frac{n-2}{n-1} \end{aligned} \tag{8}$$

Since  $A_{low}$  is sorted in ascending order,  $a_{\frac{n}{2}} = med$  and  $a_1 = min$ , we have:

$$\begin{aligned} med - min &> \frac{max - min}{2} \times \frac{n-2}{n-1} \\ med &> \frac{max + min}{2} - \frac{max - min}{2(n-1)} \end{aligned} \tag{9}$$

Substituting  $\frac{max-min}{n-1}$  with  $distance$ , we have:

$$med > \frac{max + min}{2} - \frac{distance}{2} \tag{10}$$

And because  $max - med - distance \geq med - min$ , we have  $med \leq \frac{max+min}{2} - \frac{distance}{2}$ . But we have just proved that  $med > \frac{max+min}{2} - \frac{distance}{2}$ , clearly it is a contradiction. Thus, there must exist a close pair in  $A_{low}$ .

The proof for the case when  $max - med < med - min$  (when  $n$  is odd) or  $max - med - distance < med - min$  (when  $n$  is even) is symmetrical.  $\square$

So combining Lemma1 and Lemma2, we can conclude that our algorithm can always find a pair of close elements.

### Run time analysis

The sub-routine *ComputeCloseDistance* takes  $\Theta(n)$  time. Let the total run time for *ComputeClosePair* be  $T(n)$  where  $n$  is the number of elements in the input array  $A$ . Step (1) takes  $\Theta(n)$  time, step (2) takes  $\Theta(n)$  time, step (3) takes  $\Theta(n)$  time, step (4) takes  $T(\lfloor n/5 \rfloor)$  time, step (5) takes  $\Theta(n)$  time, step (6) takes  $T(n/2)$  time because for each recursive call, we reduce the size of the input for the sub-problem into half, that is, either recurse on lower partition or higher partition. Thus, we have:

$$\begin{aligned} T(n) &= T(n/5) + T(n/2) + \Theta(n) \\ &= \Theta(n) \end{aligned} \tag{11}$$

By the master theorem, the total run time for *FindClosePair* is  $T(n) + \Theta(n) = \Theta(n)$ .

## 2. Algorithm

The algorithm works as the following: (1) Compute the index of the median:  $m = \lfloor n/2 \rfloor$ .  
(2) Find the  $m_{th}$  smallest element *median*, which is also the median of the input array.  
(3) Partition the input array around *median* into lower subarray and higher array.  
(4) recursively call on the lower sub-arrays, find it  $(k/2)_{th}$  quantiles, then output *median*, then recursively call on the lower sub-arrays, find it  $(k/2)_{th}$  quantiles. Return when  $k == 1$ .

### Correctness

The algorithm first finds the median of the input array, which is also the median of the  $k_{th}$  quantiles. Then partition the array around the median. So now we can solve two sub-problems recursively, each contains at most  $(k-1)/2$  order statistics of the original input. Thus the recursion will eventually hit the base case where  $k == 1$  and return back the  $k_{th}$  quantiles.

### Run time analysis

Step (1) takes constant time, step (2) takes  $\Theta(n)$  time, step (3) takes  $\Theta(n)$  time. The recursion has the depth of  $\log k$ .

Thus the recurrence equation is:

$$T(n, k) = 2T\left(\frac{n}{2}, \frac{k}{2}\right) + \Theta(n) \tag{12}$$

By the master theorem, the run time is  $\Theta(n \log k)$ .

## 3. Algorithm

**Function** *FindSmallestInMerge*( $A[1 : m], B[1 : n], k$ )

$i := \lfloor k/2 \rfloor$

$j := \lceil k/2 \rceil$

**if**  $k == 1$  // base case

**return**  $\min(A[1], B[1])$

**if**  $A[i] > B[j]$

$s_k := \text{FindSmallestInMerge}(A[1 : i], B[j + 1 : n], i)$

**else if**  $A[i] < B[j]$

$s_k := \text{FindSmallestInMerge}(A[i + 1 : m], B[1 : j], j)$

### Correctness

If  $A[i] > B[j]$ , where  $i := \lfloor k/2 \rfloor$  and  $j := \lceil k/2 \rceil$ , then in the merged array, there can be at most  $k - 2$  elements that are  $< B[j]$ , that is,  $A[1 : i - 1]$  and  $B[1 : j - 1]$ . So we must have the  $k_{th}$  smallest element  $s_k > B[j]$ . On the other hand, there are at least  $k - 1$  elements that are  $< A[i]$  in the merged array, that is,  $A[1 : i - 1]$  and  $B[1 : j]$ , thus we have  $s_k \leq A[i]$ .

The above analysis shows that  $s_k$  can only appear in the subarray  $B[j + 1 : n]$  or  $A[1 : i]$ . Moreover, we have thrown out  $j$  elements that  $< s_k$ , thus, the problem is reduced to finding the  $i_{th}$  smallest element in the merged array of  $B[j + 1 : n]$  and  $A[1 : i]$ .

In the case when  $A[i] < B[j]$ , the argument is symmetric.

With the above argument, we can conclude that our algorithm can find the  $k_{th}$  smallest element in the merge of two sorted arrays.

### **Run time analysis**

In each recursive call, we reduce the problem into half of its original size, and other operations executes in constant time, thus the recurrence equation is:

$$T(k) = T\left(\frac{k}{2}\right) + \Theta(1) \tag{13}$$

By the master theorem, the run time is  $\Theta(\log k)$ .