

Math4CS
Date: Dec 28
Student: Shuo Yang

1 Induction

1. Courtyard Tiling

The problem is to tile a courtyard with dimensions $2^n \times 2^n$. We are required to install a statue of a wealthy donor in one of the central square, and only special L-shaped tiles can be used. We need to prove this is feasible.

Theorem For all $n \geq 0$ there exists a tiling of a $2^n \times 2^n$ courtyard with the donor in a central square.

Proof. Prove by induction. Let $P(n)$ be the proposition that there exists a tiling of a $2^n \times 2^n$ courtyard with the donor placed in any location.

Base case: $P(0)$ is true because the donor fills the whole courtyard.

Inductive step: Suppose $P(n)$ is true, we need to prove that $P(n) \rightarrow P(n+1)$. A $2^{n+1} \times 2^{n+1}$ courtyard consists of four $2^n \times 2^n$ quadrants, each of them can be tiled with the donor placed in any location. Let the donor be in one of the four central squares, and the remaining three central squares can fit a L-shaped tile. Now we can tile each of the four quadrants by the induction hypothesis. This proves that $P(n) \rightarrow P(n+1)$. The theorem follows as a special case. \square

2 Graph

1. **Theorem.** Every graph $G = (V, E)$ has at least $|V| - |E|$ connected components.

Proof. We use induction on the number of edges. Let $P(n)$ be the proposition that every graph $G = (V, E)$ has at least $|V| - n$ connected components where $|E| = n$.

Base case: In a graph where $|E| = 0$, every vertex is a connected component itself, thus the graph has exactly $|V| - 0 = |V|$ connected components.

Inductive step: Assume that $P(n)$ holds for $n \geq 0$, that is, a graph with $|E| = n$ has at least $|V| - n$ connected components. Consider a graph with $n+1$ edges. Remove an edge (u, v) to create a n -edge graph G' , which has at least $|V| - n$ connected components. Now add (u, v) to obtain the original graph G . If u and v were in the same connected component of G' , then G has the same number of connected components as G' . If u and v were in the different connected components of G' , then adding (u, v) would merge these two components of G' into one in G , but all other components remain. In both cases, the number of connected components in G is at least $|V| - n - 1 = |V| - (n+1)$.

The theorem follows by induction. \square

2. **Theorem.** Let G be a digraph(possible with self-loops) with vertices v_1, \dots, v_n . Let M be the adjacency matrix of G . Then M_{ij}^k is equal to the number of length- k walk from v_i to v_j .

Proof. We use induction on k . Let $P(k)$ be the proposition that the number of length- k walk from v_i to v_j is M_{ij}^k , for all i, j .

Base case: for $k = 0$, a vertex only has length-0 walk to itself. Since M^0 is the identity matrix, $P(0)$ holds.

Inductive step: Assume $P(k)$ holds, we will prove that M_{ij}^{k+1} is equal to the number of length- $(k+1)$ walk from v_i to v_j . A length- $(k+1)$ walk from v_i to v_j consists of a walk of length- k from v_i to some intermediate vertex v_m followed by an edge (v_m, v_j) . Therefore, the number of M_{ij}^{k+1} is equal to:

$$M_{iv_1}^k M_{v_1j} + M_{iv_2}^k M_{v_2j} + \cdots + M_{iv_n}^k M_{v_nj} \quad (1)$$

This is exactly M_{ij}^{k+1} , thus $P(k+1)$ also holds. The theorem follows by induction. \square