

Problem Set 1 - Solution

January 8, 2015

1. Prove by contradiction. Suppose that $\sqrt[3]{2}$ is sensible. Then by definition, there exist positive integers a and b such that $\sqrt{a/b} = \sqrt[3]{2}$. Squaring both sides of the equation gives us $\sqrt{a/b} = \sqrt[3]{4}$, which implies that $\sqrt[3]{4}$ is rational.

This means there exists positive integers x and y such that $x/y = \sqrt[3]{4}$ and x/y is in lowest term. Therefore, we have:

$$x/y = \sqrt[3]{4} \quad (1)$$

$$x^3/y^3 = 4 \quad (2)$$

$$x^3 = 4y^3 \quad (3)$$

In the last equation, the right side is even, so is the left side. Since x^3 is even, x must be even. Therefore x^3 is a multiple of 8, this implies that y^3 is also a even number, thus y is even, too. x and y are both even, this contradicts the assumption that x/y is in lowest term.

Thus $\sqrt[3]{4}$ is irrational, the original assumption must not be true.

2. **A Wrong Attempt:**

$$\exists x. (E(x, y) \wedge E(x, z) \wedge x \neq y \wedge x \neq z \wedge y \neq z) \quad (4)$$

This doesn't say " x emailed exactly two other people in the class.". It also doesn't existentially quantify y and z .

Right Attempt:

First, we modify the above predicate to express that there exist students x , y and z such that x have emailed y and z .

$$\exists x \exists y \exists z. (E(x, y) \wedge E(x, z) \wedge x \neq y \wedge x \neq z \wedge y \neq z) \quad (5)$$

The following predicate restricts that x emailed exactly y and z , besides herself.

$$\forall s, E(x, s) \implies s = x \vee s = y \vee s = z \quad (6)$$

Combining these two predicates, we can say that there exists some student x who has emailed to exactly two other students y and z , besides possibly herself.

$$\exists x \exists y \exists z. (E(x, y) \wedge E(x, z) \wedge \quad (7)$$

$$x \neq y \wedge x \neq z \wedge y \neq z \wedge \quad (8)$$

$$\forall s, E(x, s) \implies s = x \vee s = y \vee s = z) \quad (9)$$

3. (a) $\exists a, b, c. (n = a \cdot a + b \cdot b + c \cdot c)$

- (b) We can express $x = 1$ as: $\forall y. (xy = y)$. Further we can express $x > 1$ as: $\exists y. (y = 1 \wedge x > y)$. Replacing $y = 1$ with the previous predicate gives us: $\exists y. (\forall z. (yz = z) \wedge x > y)$.

(c)

$$\neg(\exists x. (x > 1 \wedge x < n \wedge \exists y. (y > 1 \wedge y < n \wedge xy = n))) \quad (10)$$

A better version:

$$IS - PRIME(n) \equiv (n > 1) \wedge \neg(\exists x \exists y. (x > 1 \wedge y > 1 \wedge x \cdot y = n)) \quad (11)$$

(d)

$$\exists n \exists p \exists q. IS-PRIME(p) \wedge IS-PRIME(q) \wedge (n = p \cdot q) \wedge (p \neq q) \quad (12)$$

(e) **My Attempt:**

$$\neg(\exists n. IS-PRIME(n) \wedge IS-PRIME(p) \wedge \forall p, n > p) \quad (13)$$

Right Solution:

$$\neg(\exists n. IS-PRIME(n) \wedge (\forall p, IS-PRIME(p) \implies n \geq p)) \quad (14)$$

- (f) We can express $n > 2$ as:

$$\exists k. (k = 1) \wedge (n > k + k). \quad (15)$$

So the predicate is:

$$\forall(n), (n > 2 \wedge \exists k. n = k + k) \implies \exists p \exists q. IS-PRIME(p) \wedge IS-PRIME(q) \wedge (n = p + q) \quad (16)$$

(g)

$$\forall(n), (n > 1 \implies \exists p. IS-PRIME(p) \wedge (n < p) \wedge (p < n + n)) \quad (17)$$