

Linear Regression Models

Segment 1 – Simple Linear Regression Model

Topic 5 – Accuracy of the Model

Sudarsan N.S. Acharya (sudarsan.acharya@manipal.edu)

Topics

1. Residual Standard Error

2. R^2 Statistic

3. Summary

Residual Standard Error



MANIPAL
ACADEMY of HIGHER EDUCATION
(Institution of Eminence Deemed to be University)



Residual Standard Error

- The residual standard error (RSE) is a measure of lack of fit of the model to the data.



Residual Standard Error

- The residual standard error (RSE) is a measure of lack of fit of the model to the data.
- In an SLRM $Y = \beta_0 + \beta_1 X_1 + \varepsilon$, the random error term prevents a perfect prediction even if the population coefficients are exactly known.



Residual Standard Error

- The residual standard error (RSE) is a measure of lack of fit of the model to the data.
- In an SLRM $Y = \beta_0 + \beta_1 X_1 + \varepsilon$, the random error term prevents a perfect prediction even if the population coefficients are exactly known.
- The RSE is an in-sample estimate of the standard deviation σ of the random error term ε :



Residual Standard Error

- The residual standard error (RSE) is a measure of lack of fit of the model to the data.
- In an SLRM $Y = \beta_0 + \beta_1 X_1 + \varepsilon$, the random error term prevents a perfect prediction even if the population coefficients are exactly known.
- The RSE is an in-sample estimate of the standard deviation σ of the

random error term ε :
$$\text{RSE} = \sqrt{\frac{1}{n-2} \underbrace{\sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2}_{RSS}}$$



Residual Standard Error

- The residual standard error (RSE) is a measure of lack of fit of the model to the data.
- In an SLRM $Y = \beta_0 + \beta_1 X_1 + \varepsilon$, the random error term prevents a perfect prediction even if the population coefficients are exactly known.
- The RSE is an in-sample estimate of the standard deviation σ of the

random error term ε :
$$\text{RSE} = \sqrt{\frac{1}{n-2} \underbrace{\sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2}_{RSS}}$$

- The RSE is the amount by which the response will deviate from the true regression line on an average.

R^2 Statistic



MANIPAL
ACADEMY of HIGHER EDUCATION
(Institution of Eminence Deemed to be University)

R^2 Statistic



MANIPAL
ACADEMY of HIGHER EDUCATION
(Institution of Eminence Deemed to be University)

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 :



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model?



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**;



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**; the associated error is

R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**; the associated error is **residual sum of squares (RSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n \left(y^{(i)} - \left(\hat{\beta}_0 + \hat{\beta}_1 x_1^{(i)} \right) \right)^2$.

R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**; the associated error is **residual sum of squares (RSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n \left(y^{(i)} - \left(\hat{\beta}_0 + \hat{\beta}_1 x_1^{(i)} \right) \right)^2$.
- The R² statistic is defined as



R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**; the associated error is **residual sum of squares (RSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n \left(y^{(i)} - (\hat{\beta}_0 + \hat{\beta}_1 x_1^{(i)}) \right)^2$.
- The R² statistic is defined as $\frac{\text{TSS} - \text{RSS}}{\text{TSS}} = 1 - \frac{\text{RSS}}{\text{TSS}}$.

R² Statistic

- Suppose we have a dataset with n samples and want a simple model to predict the response Y from a single predictor X_1 : **the average model** $\hat{y}^{(i)} = \bar{y}_n$.
- What is the error associated with this model? **Total sum of squares (TSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n (y^{(i)} - \bar{y}_n)^2$.
- Suppose now we build an **SLRM**; the associated error is **residual sum of squares (RSS)** $= \sum_{i=1}^n (y^{(i)} - \hat{y}^{(i)})^2 = \sum_{i=1}^n \left(y^{(i)} - \left(\hat{\beta}_0 + \hat{\beta}_1 x_1^{(i)} \right) \right)^2$.
- The R^2 statistic is defined as $\frac{\text{TSS} - \text{RSS}}{\text{TSS}} = 1 - \frac{\text{RSS}}{\text{TSS}}$.
- The R^2 statistic varies between 0 & 1 and is a measure of the variability in the response Y that the SLRM (built using the predictor X_1) is able to explain.

Summary



MANIPAL
ACADEMY of HIGHER EDUCATION
(Institution of Eminence Deemed to be University)

Summary

Different ways to assess the accuracy of an SLRM using residual standard error (RSE) and R^2 statistic.