

# Linear Regression Models

## Segment 6 – Advanced Topics in Linear Regression

### Topic 3 – F-test in Linear Regression

Sudarsan N.S. Acharya (sudarsan.acharya@manipal.edu)

# Topics

1. Recalling Hypothesis Tests

2. The F Statistic

3. The F-test Recipe

# Recalling Hypothesis Testing



**MANIPAL**  
ACADEMY of HIGHER EDUCATION  
*(Institution of Eminence Deemed to be University)*



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ;



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.
- *Assuming that the null hypothesis is true,*



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.

- Assuming that the null hypothesis is true,  $T = \frac{\hat{\beta}_j - E[\hat{\beta}_j]}{SE(\hat{\beta}_j)} = \frac{\hat{\beta}_j - \underbrace{\beta_j}_{=0}}{SE(\hat{\beta}_j)}$





## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.

- Assuming that the null hypothesis is true,  $T = \frac{\hat{\beta}_j - E[\hat{\beta}_j]}{SE(\hat{\beta}_j)} = \frac{\hat{\beta}_j - \underbrace{\beta_j}_{=0}}{SE(\hat{\beta}_j)}$  follows a  $t$ -distribution with  $n - 2$  degrees of freedom.



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.

- Assuming that the null hypothesis is true,  $T = \frac{\hat{\beta}_j - E[\hat{\beta}_j]}{SE(\hat{\beta}_j)} = \frac{\hat{\beta}_j - \underbrace{\beta_j}_{=0}}{SE(\hat{\beta}_j)}$  follows a  $t$ -distribution with  $n - 2$  degrees of freedom.
- From a particular dataset, we have the estimate  $\hat{\beta}_j$  and the realized value of  $T$  denoted as  $t$ .



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.

- Assuming that the null hypothesis is true,  $T = \frac{\hat{\beta}_j - E[\hat{\beta}_j]}{SE(\hat{\beta}_j)} = \frac{\hat{\beta}_j - \underbrace{\beta_j}_{=0}}{SE(\hat{\beta}_j)}$  follows a  $t$ -distribution with  $n - 2$  degrees of freedom.
- From a particular dataset, we have the estimate  $\hat{\beta}_j$  and the realized value of  $T$  denoted as  $t$ .
- We calculate  $P(T \geq |t|)$ , called the p-value,



## Recalling Hypothesis Testing

- Suppose we want to know if there is a linear relationship between the  $j$ th predictor predictor and the response variable.
- We consider a *null hypothesis*  $\beta_j = 0$ ; that is, we hypothesize that there is no relationship between the  $j$ th predictor and the response variable.

- Assuming that the null hypothesis is true,  $T = \frac{\hat{\beta}_j - E[\hat{\beta}_j]}{SE(\hat{\beta}_j)} = \frac{\hat{\beta}_j - \underbrace{\beta_j}_{=0}}{SE(\hat{\beta}_j)}$  follows a  $t$ -distribution with  $n - 2$  degrees of freedom.
- From a particular dataset, we have the estimate  $\hat{\beta}_j$  and the realized value of  $T$  denoted as  $t$ .
- We calculate  $P(T \geq |t|)$ , called the p-value, and reject the null hypothesis if the p-value is smaller than a threshold, typically 0.05.

# The F Statistic



**MANIPAL**  
ACADEMY of HIGHER EDUCATION  
*(Institution of Eminence Deemed to be University)*



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero.



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?
- Suppose we want to test the null hypothesis that the regression coefficients associated with all the predictors in a particular set  $S$  are equal to zeros.





## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?
- Suppose we want to test the null hypothesis that the regression coefficients associated with all the predictors in a particular set  $S$  are equal to zeros.
- For this purpose, we calculate



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?
- Suppose we want to test the null hypothesis that the regression coefficients associated with all the predictors in a particular set  $S$  are equal to zeros.
- For this purpose, we calculate (1) the residual vector  $r$  using all the  $p$  predictors



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?
- Suppose we want to test the null hypothesis that the regression coefficients associated with all the predictors in a particular set  $S$  are equal to zeros.
- For this purpose, we calculate (1) the residual vector  $r$  using all the  $p$  predictors (2) the residual vector  $r_S$  by excluding the predictors in the set  $S$ ,



## The F Statistic

- The approach so far (called the t-test or the Wald test) lets us test whether the regression coefficient associated with one predictor is zero. What if we want to test for several regression parameters *simultaneously*?
- Suppose we want to test the null hypothesis that the regression coefficients associated with all the predictors in a particular set  $S$  are equal to zeros.
- For this purpose, we calculate (1) the residual vector  $\mathbf{r}$  using all the  $p$  predictors (2) the residual vector  $\mathbf{r}_S$  by excluding the predictors in the set  $S$ , (3) calculate the F statistic

$$\frac{(\|\mathbf{r}_S\|^2 - \|\mathbf{r}\|^2) / (p - |S|)}{\|\mathbf{r}\| / (n - p)}$$

# The F-test Recipe



**MANIPAL**  
ACADEMY of HIGHER EDUCATION  
*(Institution of Eminence Deemed to be University)*



# The F-test Recipe

- The null hypothesis is that  $\beta_j = 0$  for regression coefficients associated with the predictors in the set  $S$ .



## The F-test Recipe

- The null hypothesis is that  $\beta_j = 0$  for regression coefficients associated with the predictors in the set  $S$ .
- If this null hypothesis is true, and if the random error terms are normally distributed, then the  $F$  distribution can be used to determine how unlikely our observed value of the  $F$  statistic is given that the null hypothesis is true.



## The F-test Recipe

- The null hypothesis is that  $\beta_j = 0$  for regression coefficients associated with the predictors in the set  $S$ .
- If this null hypothesis is true, and if the random error terms are normally distributed, then the  $F$  distribution can be used to determine how unlikely our observed value of the  $F$  statistic is given that the null hypothesis is true.
- If the  $F$  statistic much larger than 1, then we reject the null hypothesis;





## The F-test Recipe

- The null hypothesis is that  $\beta_j = 0$  for regression coefficients associated with the predictors in the set  $S$ .
- If this null hypothesis is true, and if the random error terms are normally distributed, then the  $F$  distribution can be used to determine how unlikely our observed value of the  $F$  statistic is given that the null hypothesis is true.
- If the  $F$  statistic much larger than 1, then we reject the null hypothesis; otherwise, we fail to reject the null hypothesis.



## The F-test Recipe

- The null hypothesis is that  $\beta_j = 0$  for regression coefficients associated with the predictors in the set  $S$ .
- If this null hypothesis is true, and if the random error terms are normally distributed, then the  $F$  distribution can be used to determine how unlikely our observed value of the  $F$  statistic is given that the null hypothesis is true.
- If the  $F$  statistic much larger than 1, then we reject the null hypothesis;
- The  $F$ -test is a simple quantification our level of uncertainty about the linear relationship of multiple predictors at once with the response variable.

# Summary



**MANIPAL**  
ACADEMY of HIGHER EDUCATION  
*(Institution of Eminence Deemed to be University)*

# Summary

- F statistic and the F-test in the context of linear regression.

# Summary

- F statistic and the F-test in the context of linear regression.
- t-test and F-test for analyzing predictors in linear regression.