Doubt Clearing Session

Foundation Course on Discrete Mathematics for GATE

Sanchit Jain • Lesson 8 • Feb 12, 2022

Cartesian Product

1. Cartesian Product of two sets A and B in the set of all ordered pairs, whose first member belongs to the first set and second member belongs to the second set, denoted by $A \times B$.

2. For E.g. if $A = \{a, b\}$, $B = \{1, 2, 3\}$

3. $A \times B = \{$ $\}$

a

b

1

2

3

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- It is a kind of maximum relation possible, where every member of the first set belong to every member of the second set.

- $A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}$



Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. In general, commutative law does not hold good $A\times B \mathrel{!=} B\times A$

2. If $|A| = m$ and $|B| = n$ then $|A\times B| =$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The number of binary relations on a set with n elements is: **(GATE-1999) (1 Marks)**

(A) n^2

(B) 2^n

(C) 2^{n^2}

(D) None of the above

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let A be a finite set of size n. The number of elements in the power set of A × A is: **(GATE-1993) (1 Marks)**

a) $2^{(2^n)}$

b) $2^{(n^2)}$

c) 2^n

d) None of the above

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Relation

- **Relation**: - Let A and B are sets then every possible subset of 'A×B' is called a relation from A to B.

- If |A| =m and |B| = n then total no of element(pair) will be m*n, every element will have two choice weather to present or not present in the subset(relation), therefore the total number of relation possible is _____

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, B = {1, 2}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, B = {1, 2}, A×B = {(a, 1), (a, 2), (b, 1), (b, 2)}

(a , 1)	(a , 2)	(b , 1)	(b , 2)	
0	0	0	0	
0	0	0	1	
0	0	1	0	
0	0	1	1	
0	1	0	0	
0	1	0	1	
0	1	1	0	
0	1	1	1	
1	0	0	0	
1	0	0	1	
1	0	1	0	
1	0	1	1	
1	1	0	0	
1	1	0	1	
1	1	1	0	
1	1	1	1	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, B = {1, 2}

(a , 1)	(a , 2)	(b , 1)	(b , 2)	
0	0	0	0	{ }
0	0	0	1	{(b, 2)}
0	0	1	0	{(b, 1)}
0	0	1	1	{(b, 1), (b, 2)}
0	1	0	0	{(a, 2)}
0	1	0	1	{(a, 2), (b, 2)}
0	1	1	0	{(a, 2), (b, 1)}
0	1	1	1	{(a, 2), (b, 1), (b, 2)}
1	0	0	0	{(a, 1)}
1	0	0	1	{(a, 1), (b, 2)}
1	0	1	0	{(a, 1), (b, 1)}
1	0	1	1	{(a, 1), (b, 1), (b, 2)}
1	1	0	0	{(a, 1), (a, 2)}
1	1	0	1	{(a, 1), (a, 2), (b, 2)}
1	1	1	0	{(a, 1), (a, 2), (b, 1)}
1	1	1	1	{(a, 1), (a, 2), (b, 1), (b, 2)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Matrix Representation

	1	2	3
x	$(x,1)$	$(x,2)$	$(x,3)$
y	$(y,1)$	$(y,2)$	$(y,3)$
z	$(z,1)$	$(z,2)$	$(z,3)$

B

A

$A \times B$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- Largest relation possible will be _____

- Smallest possible relation will be _____

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Complement of a relation</u>**: - Let R be a relation from A to B, then the complement of relation will be denoted by R', R^C or ~~R.~~

- $R' = \{(a, b) \mid (a, b) \in A \times B, (a, b) \in! R\}$

- $R' = (A \times B) - R$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- For E.g. if $A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$
- $R = \{(a, 1), (a, 3), (b, 2)\}$
- $R' = \{ \qquad \}$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

$R \cup R' =$

$R \cap R' =$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Inverse of a relation</u>**: - Let R be a relation from A to B, then the inverse of relation will be a relation from B to A, denoted by R^{-1}.

- $R^{-1} = \{(b, a) \mid (a, b) \in R\}$

- $A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$

- $R = \{(a, 1), (a, 3), (b, 2)\}$

- $R^{-1} = \{ \qquad \}$

- $|R| \square |R^{-1}|$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Diagonal relation</u>**: - A relation R on a set A is said to be diagonal relation if, R is a set of all ordered pair (x, x), for every $\forall x \in A$, sometimes it is also denoted by $\blacktriangle_A$

- $R = \{(x, x) \mid \forall x \in A\}$

	1	2	3
1	11		
2		22	
3			33

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Types of a Relation

- To further study types of relations, we consider a set A with n elements, then a cartesian product A×A will have n^2 elements(pairs). Therefore, total number of relation possible is $2^{n * n}$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Reflexive relation</u>**: - A relation R on a set A is said to be reflexive,
- If $\forall x \in A$
  - $(x, x) \in R$

	1	2	3
1	11		
2		22	
3			33

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q consider a set A = {1,2,3}, find which of the following relations are reflexive and Irreflexive?

	Relation	Reflexive	Irreflexive
1	A×A		
2	ϕ		
3	{(1,1), (2,2), (3,3)}		
4	{(1,2), (2,3), (1,3)}		
5	{(1,1), (1,2), (2,1), (2,2)}		
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}		
7	{(1,3), (2,1), (2,3), (3,2)}		

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest reflexive relation is _____

2. Largest reflexive relation is _____

3. Total number of reflexive relations will be _____

4. If two relations R_1 and R_2 are reflexive then their union and intersection will also be reflexive (T / F).

5. Any super set of reflexive relation will also be reflexive(T / F).

6. If a relation is reflexive then its inverse R^{-1} will also be reflexive (T / F).

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest reflexive relation is $\blacktriangle_A$

2. Largest reflexive relation is $A \times A$

3. Total number of reflexive relations will be $2^{n(n-1)}$

4. If two relations R_1 and R_2 are reflexive then their union and intersection will also be reflexive.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	
0	0	0	1	
0	0	1	0	
0	0	1	1	
0	1	0	0	
0	1	0	1	
0	1	1	0	
0	1	1	1	
1	0	0	0	
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q What is the possible number of reflexive relations on a set of 5 elements? **(GATE-2010) (1 Marks)**

(A) 2^{10} **(B)** 2^{15} **(C)** 2^{20} **(D)** 2^{25}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Irreflexive relation</u>**: - A relation R on a set A is said to be Irreflexive,

1. If $\forall x \in A$
2. $(x, x) \notin R$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q consider a set A = {1,2,3}, find which of the following relations are reflexive and Irreflexive?

	Relation	Reflexive	Irreflexive
1	A×A	Y	
2	ϕ	N	
3	{(1,1), (2,2), (3,3)}	Y	
4	{(1,2), (2,3), (1,3)}	N	
5	{(1,1), (1,2), (2,1), (2,2)}	N	
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}	Y	
7	{(1,3), (2,1), (2,3), (3,2)}	N	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest irreflexive relation is _____

2. Largest irreflexive relation is _____

3. Total number of irreflexive relation will be _____

4. If two relations R_1 and R_2 are Irreflexive then their union and intersection will also be Irreflexive (T / F).

5. If a relation R on a set A is reflexive, then R^C is Irreflexive, and vice versa (T / F).

6. Any sub set of irreflexive relation will also be irreflexive(T / F).

7. If a relation is irreflexive then its inverse R^{-1} will also be irreflexive (T / F).

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	
0	0	1	0	{(b, a)}
0	0	1	1	
0	1	0	0	{(a, b)}
0	1	0	1	
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	
1	0	0	0	
1	0	0	1	
1	0	1	0	
1	0	1	1	
1	1	0	0	
1	1	0	1	
1	1	1	0	
1	1	1	1	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- Smallest irreflexive relation is ϕ
- Largest irreflexive relation is $(A \times A) - \blacktriangle_A$
- Total number of irreflexive relation will be $2^{n(n-1)}$
- If two relations R_1 and R_2 are Irreflexive then their union and intersection will also be Irreflexive.
- If a relation R on a set A is reflexive, then R^C is Irreflexive

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Suppose that R_1 and R_2 are reflexive relations on a set A. Which of the following statements is correct? **(NET-July-2016)**

a) $R_1 \cap R_2$ is reflexive and $R_1 \cup R_2$ is irreflexive.

b) $R_1 \cap R_2$ is irreflexive and $R_1 \cup R_2$ is reflexive.

c) Both $R_1 \cap R_2$ and $R_1 \cup R_2$ are reflexive.

d) Both $R_1 \cap R_2$ and $R_1 \cup R_2$ are irreflexive

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Symmetric relation</u>**: - A relation R on a set A is said to be Symmetric,

If $\forall$ a, b $\in$ A

(a, b) $\in$ R

.....

then (b, a) $\in$ R

.....

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q consider a set A = {1,2,3}, find which of the following relations are Symmetric, Anti-Symmetric and Asymmetric?

	Relation	Symmetric	Anti-Symmetric	Asymmetric
1	$A \times A$			
2	ϕ			
3	{(1,1), (2,2), (3,3)}			
4	{(1,2), (2,3), (1,3)}			
5	{(1,1), (1,2), (2,1), (2,2)}			
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}			
7	{(1,3), (2,1), (2,3), (3,2)}			

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest symmetric relation is _____

2. Largest symmetric relation is_____

3. Total number of symmetric relation will be _____

4. If a relation on a set A is symmetric then R __ R^{-1}

5. If two relations R_1 and R_2 are symmetric then their Union and Intersection will also be symmetric. (T / F)

6. If a relation is symmetric then its superset and subset will always be symmetric. (T / F)

7. If a relation is symmetric then its complement R^c will always be symmetric. (T / F)

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	
0	0	1	1	
0	1	0	0	
0	1	0	1	
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	
1	0	1	1	
1	1	0	0	
1	1	0	1	
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest symmetric relation is ϕ

2. Largest symmetric relation is $A \times A$

3. Total number of symmetric relation will be $2^{[n(n+1)]/2}$

4. If a relation on a set A is symmetric then $R = R^{-1}$

5. If two relations R1 and R2 are symmetric then their union and intersection will also be symmetric.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R be a relation on the set of ordered pairs of positive integers such that ((p, q), (r, s)) ∈ R if and only if p–s = q–r. Which one of the following is true about R? **(GATE-2015) (2 Marks)**

(A) Both reflexive and symmetric

(B) Reflexive but not symmetric

(C) Not reflexive but symmetric

(D) Neither reflexive nor symmetric

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q How many relations are there on a set with n elements that are symmetric and a set with n elements that are reflexive and symmetric? **(NET-Dec-2012)**

A) $2^{n(n+1)/2}$ and $2^{n}.3^{n(n-1)/2}$

B) $3^{n(n-1)/2}$ and $2^{n(n-1)}$

C) $2^{n(n+1)/2}$ and $3^{n(n-1)/2}$

D) $2^{n(n+1)/2}$ and $2^{n(n-1)/2}$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Anti-Symmetric relation</u>**: - A relation R on a set A with cartesian product A×A is said to be Anti-Symmetric,

If $\forall\, a, b \in A$

$(a, b) \in R$

$(b, a) \in R$

.....

$a = b$

.....

Conclusion: Symmetry is not allowed but diagonal pairs are allowed

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

	Relation	Symmetric	Anti-Symmetric	Asymmetric
1	A×A	Y		
2	ϕ	Y		
3	{(1,1), (2,2), (3,3)}	Y		
4	{(1,2), (2,3), (1,3)}	N		
5	{(1,1), (1,2), (2,1), (2,2)}	Y		
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}	N		
7	{(1,3), (2,1), (2,3), (3,2)}	N		

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest Anti-symmetric relation is _____

2. Largest Anti-symmetric relation will contain _____elements

3. Total number of Anti-symmetric relation will be _____

4. A relation R on a set A is Anti-Symmetric if $(R \cap R^{-1}) \sqsubseteq \blacktriangle_A$ (T / F)

5. Sub set of a Anti-Symmetric will also be (T / F)

6. Super set of a Anti-Symmetric will also be (T / F)

7. If two relations R_1 and R_2 are Anti-symmetric then their _____need not to be Anti-symmetric but _____ will also be Anti-symmetric.

8. If a relation is Anti-symmetric then its complement R^c will always be Anti-symmetric. (T / F)

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	
0	1	1	1	
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	
1	1	1	1	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest Anti-symmetric relation is ϕ

2. Largest Anti-symmetric relation will contain $n(n+1)/2$ elements

3. Total number of Anti-symmetric relation will be $2^n * 3^{[n(n-1)]/2}$

4. A relation R on a set A is Anti-Symmetric if $(R \cap R^{-1}) \sqsubseteq \blacktriangle_A$

5. Sub set of a Anti-Symmetric will also be Anti-Symmetric

6. If two relations R1 and R2 are Anti - symmetric then their union need not to be Anti-symmetric but intersection will also be Anti-symmetric.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Consider a set A = {a, b, c} and R_1, R_2, R_3 and R_4 are relations on A which of the following is not true?

		Symmetric	Anti-Symmetric	True
1	$R_1 = \{(a, a), (c, c)\}$	Y	Y	
2	$R_2 = \{(a, b), (b, a), (a, c)\}$	N	N	
3	$R_3 = \{(a, b), (b, c), (a, c)\}$	N	Y	
4	$R_4 = \{(a, b), (b, a), (c, c)\}$	Y	N	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Consider the binary relation R = {(x, y), (x, z), (z, x), (z, y)} on the set {x, y, z}. Which one of the following is TRUE? **(GATE-2009) (1 Marks)**

(A) R is symmetric but NOT antisymmetric

(B) R is NOT symmetric but antisymmetric

(C) R is both symmetric and antisymmetric

(D) R is neither symmetric nor antisymmetric

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Asymmetric relation</u>**: - A relation R on a set A is said to be Asymmetric,

If $\forall$ a, b $\in$ A

(a, b) $\in$ R

.....

(b, a) $\notin$ R

.....

Conclusion: Symmetry is not allowed; even diagonal pairs are not allowed

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

	Relation	Symmetric	Anti-Symmetric	Asymmetric
1	$A \times A$	Y	N	
2	ϕ	Y	Y	
3	{(1,1), (2,2), (3,3)}	Y	Y	
4	{(1,2), (2,3), (1,3)}	N	Y	
5	{(1,1), (1,2), (2,1), (2,2)}	Y	N	
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}	N	Y	
7	{(1,3), (2,1), (2,3), (3,2)}	N	N	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest Asymmetric relation is _____

2. Largest Asymmetric relation will contain _____elements

3. Total number of Asymmetric relation will be _____

4. Every asymmetric relation is also anti-symmetric (T / F)

5. Sub set of a Asymmetric will also be Asymmetric (T / F)

6. Super set of a Asymmetric will also be Asymmetric(T / F)

7. If two relations R_1 and R_2 are Asymmetric then their Union will also be Asymmetric(T / F).

8. If two relations R_1 and R_2 are Asymmetric then their Intersection will also be Asymmetric(T / F).

9. If a relation is Asymmetric then its complement R^c will always be Asymmetric. (T / F)

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	
0	0	1	0	{(b, a)}
0	0	1	1	
0	1	0	0	{(a, b)}
0	1	0	1	
0	1	1	0	
0	1	1	1	
1	0	0	0	
1	0	0	1	
1	0	1	0	
1	0	1	1	
1	1	0	0	
1	1	0	1	
1	1	1	0	
1	1	1	1	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Every asymmetric relation is also anti-symmetric

2. Smallest Asymmetric relation is ϕ

3. Largest Asymmetric relation will contain $n(n-1)/2$ elements

4. Total number of Asymmetric relation will be $3^{[n(n-1)]/2}$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Transitive relation</u>**: - A relation R on a set A is said to be Transitive,

If $\forall$ a, b, c $\in$ A

(a, b) $\in$ R

(b, c) $\in$ R

.....

(a, c) $\in$ R

.....

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

	Relation	Transitive
1	$A \times A$	
2	ϕ	
3	{(1,1), (2,2), (3,3)}	
4	{(1,2), (2,3), (1,3)}	
5	{(1,1), (1,2), (2,1), (2,2)}	
6	{(1,1), (2,2), (3,3), (1,3), (2,1)}	
7	{(1,3), (2,1), (2,3), (3,2)}	
8	{(1,2)}	
9	{(1,3), (2,3)}	
10	{(1,2), (1,3)}	
11	{(2,3), (1,2)}	

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest Asymmetric relation is _____

2. Largest Asymmetric relation will contain _____ elements

3. If two relations R_1 and R_2 are Transitive then their _____ need not to be transitive but _____ will also be transitive.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	{(a, b), (b, a)}
0	1	1	1	{(a, b), (b, a), (b, b)}
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	{(a, a), (a, b), (b, a)}
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

For E.g. if A = {a, b}, A×A = {(a, a), (a, b), (b, a), (b, b)}

(a , a)	(a , b)	(b , a)	(b , b)	
0	0	0	0	{ }
0	0	0	1	{(b, b)}
0	0	1	0	{(b, a)}
0	0	1	1	{(b, a), (b, b)}
0	1	0	0	{(a, b)}
0	1	0	1	{(a, b), (b, b)}
0	1	1	0	
0	1	1	1	
1	0	0	0	{(a, a)}
1	0	0	1	{(a, a), (b, b)}
1	0	1	0	{(a, a), (b, a)}
1	0	1	1	{(a, a), (b, a), (b, b)}
1	1	0	0	{(a, a), (a, b)}
1	1	0	1	{(a, a), (a, b), (b, b)}
1	1	1	0	
1	1	1	1	{(a, a), (a, b), (b, a), (b, b)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. Smallest Asymmetric relation is ϕ

2. Largest Asymmetric relation will contain $A \times A$ elements

3. If two relations R_1 and R_2 are Transitive then their union need not to be transitive but intersection will also be transitive.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

$\|A\| = n$	No of transitive relation
0	1
1	2
2	13
3	171
4	3994

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Warshall's Algorithm: -

Q Consider a set A = {1,2,3} and a relation **R** = {(1,1), (1,3), (2,2), (3,1), (3,2)}?

	1	2	3
Column			
Row			

	1	2	3
1			
2			
3			

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Consider a set A = {1,2,3,4} and a relation

R = {(1, 1), (2, 1), (2, 2), (2, 3), (2, 4), (3, 1), (3, 2), (3, 3), (3, 4)}?

	1	2	3	4
Column				
Row				

	1	2	3	4
1				
2				
3				
4				

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Warshall's Algorithm: -

Q Consider a set A = {1,2,3} and a relation **R** = {(1,1), (1,3), (2,2), (3,1), (3,2)}?

	1	2	3
Column			
Row			

	1	2	3
1			
2			
3			

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q A binary relation R on N x N is defined as follows:

(a, b) R (c, d) if a <= c or b <= d

Consider the following propositions:

P: R is reflexive

Q: R is transitive

Which one of the following statements is TRUE? **(GATE- 2016) (2 Marks)**

(A) Both P and Q are true. – 46

(B) P is true and Q is false. – 33

(C) P is false and Q is true. – 15

(D) Both P and Q are false. – 7

(a, b) (c, d)

$((a,b), (c,d))$

$a \le c$ or $b \le d$

$(17, 30)$ $(17, 30)$

$[(1,2), (3,4)] \in R$

$(3,4), (1,2) \notin R$

1 → ~~(1,1)~~ ~~(1,2)~~

2 → ~~(1,1)~~ ~~(1,1)~~

3 → (1,2), (1,2)

$(5,6)$ $(6,3) \in R$

$(6,3)$ $(2,4) \in R$

$(5,6)$ $(2,4) \notin R$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R be the relation on the set of positive integers such that a R b if and only if a and b are distinct and have a common divisor other than 1. Which one of the following statements about R is True? **(GATE-2015) (1 Marks)**

(A) R is symmetric and reflexive but not transitive

(B) R is reflexive but not symmetric and not transitive

(C) R is transitive but not reflexive and not symmetric

(D) R is symmetric but not reflexive and not transitive

(a,b) ∈ R // aRb

(15,b) (6,15)

(15,6) (6,15)

(15,15)

(3,6) (6,3)

(3,3)

(15, 6)

(4,4) (5,5) (6,6) (1,1) (1,1) (1,3)

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The relation "divides" on a set of positive integers is _____. **(NET-June-2013)**

a) Symmetric and transitive

b) Anti symmetric and transitive

c) Symmetric only

d) Transitive only

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q A relation **R** in {1, 2,3,4,5,6} is given by {(1,2), (2,3), (3,4), (4,4), (4,5)}. This relation is: **(NET-Dec-2008)**

a) Reflexive

b) Symmetric

c) Transitive

d) not reflexive, not symmetric and not transitive

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q the transitive closure of a relation R on a set A whose relation matrix $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$ is:

(NET-June-2005)

a) $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$ **b)** $\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$ **c)** $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ **d)** $\begin{bmatrix} 0 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$

20 43 53 4

$$\begin{array}{c|ccc} & a & b & c \\ \hline a & 0 & 1 & 0 \\ b & 0 & 0 & 1 \\ c & 1 & 0 & 0 \end{array} =$$

	1	2	3
c	c	a	b
R	b	c	a
	cb	ac	ba

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The binary relation S = ϕ (empty set) on set A = {1,2,3} is **(GATE-2002) (2 Marks)**

(a) Neither reflexive nor symmetric

(b) Symmetric and reflexive

(c) Transitive and reflexive

(d) Transitive and symmetric

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The binary relation R= {(1,1), (2,1), (2,2), (2,3), (2,4), (3,1), (3,2), (3,3), (3,4)} on the set A = {1,2,3,4} is **(GATE-1998) (2 Marks)**

(a) reflexive, symmetric and transitive

(b) neither reflexive, nor irreflexive but transitive

(c) irreflexive, symmetric and transitive

(d) irreflexive and antisymmetric

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The transitive closure of the relation {(1,2), (2,3), (3,4), (5,4)} on the set {1,2,3,4,5} is _____. **(GATE-1989) (2 Marks)**

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Equivalence Relation</u>**: - A relation R on a set A with cartesian product A×A is said to be Equivalence, if it is

1. **Reflexive**
2. **Symmetric**
3. **Transitive**

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- If two relations R_1 and R_2 are Equivalence then their union need not to be equivalence but intersection will also be Equivalence.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

R_1 : (a, b) iff (a + b) is even over the set of integers

$\in R$

R ✓ : $(z_1 + z_1) \in$ even

S ✓ : $(a,b) \Leftrightarrow (b,a)$ $\quad a+b = b+a$

T ✓ : $a, b \in R$

$b, c \in R$

$a, c \in R$ ✓

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

R_2 : (a, b) iff (a + b) is odd over the set of integers

R $(2, +2)$

X

S

T

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

R_3 : (a, b) iff a. b > 0 over the set of non-zero rational numbers

R: $(\overset{+}{\underset{-}{\ }}) \times (\overset{+}{\underset{-}{\ }}) = +$

E.R

S: $(2, -3) \notin R$

$(-3, 2) \notin R$

$a \times b > 0$

T: $b \times c > 0$

$a \times c > 0$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

R_4 : (a, b) iff $|a - b| \leq 2$ over the set of natural numbers

$\cancel{|a-b|}$

(1, 1)

(2, 1)

R :

(2, 2)

$\cancel{a = -7}$

(a, 1) ∈ R

|a − 2|

$\cancel{7 \leq 2}$

En ✓

$a - b \leq 2$

S :

$|a - b| = |b - a|$

5 − 3 = 3 − 5

2 −2

(5, 7) ← 1

(7, 6) ← R

(5, 6) ∈ R

~~R :~~

~~(5, 3)~~

~~(3, 1)~~

~~(5, 1)~~

2

2

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Which of the relations on {0, 1, 2, 3} is an equivalence relation? **(NET-July-2018)**

a) ~~{(0, 0) (0, 2) (2, 0) (2, 2) (2, 3) (3, 2) (3, 3)}~~

b) {(0, 0) (1, 1) (2, 2) (3, 3)}

Δ_A

R

S

T

c) ~~{(0, 0) (0, 1) (0, 2) (1, 0) (1, 1) (1, 2) (2, 0)}~~

d) ~~{(0, 0) (0, 2) (2, 3) (1, 1) (2, 2)}~~

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let S be a set of n elements. The number of ordered pairs in the largest and the smallest equivalence relations on S are **(GATE-2007) (1 Marks)**

(A) n and n

(B) n^2 and n

(C) n^2 and 0

(D) n and 1

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R and S be any two equivalence relations on a non-empty set A. Which one of the following statements is TRUE? **(GATE-2005) (2 Marks)**

(A) $R \cup S$, $R \cap S$ are both equivalence relations

(B) $R \cup S$ is an equivalence relation

(C) $R \cap S$ is an equivalence relation

(D) Neither $R \cup S$ nor $R \cap S$ is an equivalence relation

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Suppose A is a finite set with n elements. The number of elements in the largest equivalence relation of A is? **(GATE-1998)** **(1 Marks)**

(a) n **(b)** n^2 **(c)** 1 **(d)** n + 1

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R_1 and R_2 be two equivalence relations on a set. Consider the following assertions **(GATE-1998) (1 Marks)**

(i) $R_1 \cup R_2$ is an equivalence relation

(ii) $R_1 \cap R_2$ is an equivalence relation

Which of the following is correct?

a) Both assertions are true

b) Assertions (i) is true but assertions (ii) is not true

c) Assertions (ii) is true but assertions (i) is not true

d) Neither (i) nor (ii) is true

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R be a non-empty relation on a collection of sets defined by A R B if and only if $A \cap B = \phi$. Then, (pick the true statement) **(GATE-1996) (2 Marks)**

(a) R is reflexive and transitive

(b) R is symmetric and not transitive

(c) R is an equivalence relation

(d) R is not reflexive and not symmetric

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. In mathematics, when the elements of some set *S* have a notion of equivalence (formalized as an equivalence relation) defined on them, then one may naturally split the set *S* into **equivalence classes**.

2. These equivalence classes are constructed so that elements *a* and *b* belong to the same **equivalence class** if and only if they are equivalent.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **Equivalence Class:** - of an element is denoted by [x].
  $[x] = \{y \mid y \in A \text{ and } (x, y) \in R\}$ for all $x \in A$

- We can have [x] = [y], even if x != y

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Consider A = {1, 2, 3, 4, 5} an equivalence relation R on A, R = {(1,1),(2,2),(3,3),(4,4),(5,5),(1,4),(4,1),(2,5),(5,2)} find the partition of a set A, defined by R.

[1] = {1,4}

[2] = {2,5}

[3] = {3}

[4] = {1,4}

[5] = {2,5}

|A| = n

n

{1,2,3,4,5}

$P_1 = \{1,4\}$ $P_2 = \{2,5\}$ $P_3 = \{3\}$

|A| = n

1

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Partitions of a Set: - let A be a set, with n elements. Based on our understanding of equivalent classes, a subdivision of A into non-empty and non-overlapping subset is called a partition of A.

$$A_1 \cup A_2 \cup A_3 \cup \ldots\ldots\ldots\ldots \cup A_n = A$$

$$A_1 \cap A_2 \cap A_3 \cap \ldots\ldots\ldots\ldots \cap A_n = \phi$$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Consider A = {1, 2, 3, 4, 5} an equivalence relation R on A, R = {(1,1),(2,2),(3,3),(4,4),(5,5),(1,4),(4,1),(2,5),(5,2)} find the partition of a set A, defined by R.

[1] = {1, 4}

[2] = {2, 5}

[3] = {3}

[4] = {1,4}

[5] = {2, 5}

so we have partitions =

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let A = {1,2,3,4,5} is a set having partitions as {1, 4}, {2, 3, 5}, find the equivalence relation from which these partitions are created?

$\{1,4\} \times \{1,4\}$

$(1,1)\ (1,4)\ (4,1)\ (4,4)$

$\{2,3,5\} \times \{2,3,5\}$

$(1,2)\ (2$

$E - R \longleftrightarrow P$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q A relation R is defined on the set of integers as x Ry iff (x + y) is even. Which of the following statements is true? **(GATE-2000) (2 Marks)**

(a) R is not an equivalence relation — 4

(b) R is an equivalence relation having 1 equivalence class — 19

(c) R is an equivalence relation having 2 equivalence classes — 66

(d) R is an equivalence relation having 3 equivalence classes — 11

$[-1] = \{\pm 1, \pm 3, \pm 5, \pm 7, \ldots\}$

$[0] = \{0, \pm 2, \pm 4, \pm 6, \pm 8, \ldots\}$ ← even

$[\pm 1] = \{\pm 1, \pm 3, \pm 5, \pm 7 \ldots\}$ ← odd

$[+1] = \{0, \pm 2, \pm 4, \ldots\}$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q How many different equivalence relations with exactly three different equivalence classes are there on a set with five elements? **(NET-July-2016)**

(A) 10 **(B)** 15 **(C)** 25 **(D)** 30

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q The number of equivalence relations of the set {1,2,3,4} is **(GATE-1997) (1 Marks)**

a) 15 **b)** 16 **c)** 24 **d)** 4

Will answer this question after functions

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Break

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Partial Order Relation</u>**: - A relation R on a set A with cartesian product A×A is said to be partial order, if it is

1. **Reflexive**
2. **Anti - Symmetric**
3. **Transitive**

$\forall A, B \in U$

$(A, B) \in R$

if $A \subseteq B$

$\forall a, b \in Z$

$(a, b) \in R$

if $a \leq b$

$A \subseteq A$

$A \subseteq B \quad B \subseteq A$

$A \subseteq B$

$B \subseteq C$

$A \subseteq C$

$R = \quad -2 \leq -2$

$+13 \leq +13$

anti $\quad 1 \leq 2 \quad (1 \leq 1)$

$\cancel{(2 \leq 1)}$

$a \leq b$

$b \leq c$

$a \leq c$

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

1. In mathematics, especially order theory, a **partially ordered set** (also **poset**) formalizes and generalizes the intuitive concept of an ordering, sequencing, or arrangement of the elements of a set.

2. A poset consists of a set together with a binary relation indicating that, for certain pairs of elements in the set, one of the elements precedes the other in the ordering.

3. The relation itself is called a "partial order." The word *partial* in the names "partial order" and "partially ordered set" is used as an indication that not every pair of elements needs to be comparable.

4. That is, there may be pairs of elements for which neither element precedes the other in the poset. Partial orders thus generalize total orders, in which every pair is comparable.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Partial ordering set (Poset):</u>** - a set A with partial ordering relation R defined on A is called a POSET and is denoted by [A, R]

- For e.g. [A, /], [A, <=], [P(S), ⊑]

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

- **<u>Total order relation:</u>** - A Poset [A, R] is called a total order set, if every pair of elements are comparable i.e. either (a, b) or (b, a) $\in$ R, for $\forall$ a, b $\in$ A

- For e.g. A = {1, 2, 3, 6}, then Poset [A,/] is not a total order relation but A = {1,2,4,8} will be

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Which of the following statements is true? **(NET-July-2018)**

a) $(Z, \leq)$ is not totally ordered

b) The set inclusion relation $\subseteq$ is a partial ordering on the power set of a set S

c) $(Z, \neq)$ is a Poset

d) The directed graph 

is not a partial order

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q A partial order P is defined on the set of natural numbers as follows. Here x/y denotes integer division. **(GATE-2007) (2 Marks)**

(1) (0,0) ∈ P **(2)** (a, b) ∈ P if and only if a % 10 <= b % 10 and (a/10, b/10) ∈ P.

Consider the following ordered pairs:

(i) (101,22) **(ii)** (22,101) **(iii)** (145,265) **(iv)** (0,153)

a) I & iii **b)** ii & iv **c)** I & iv **d)** iii & iv

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q A relation R is defined on ordered pairs of integers as follows: (x, y) R (u, v) if x < u and y > v. Then R is **(GATR-2006) (1 Marks)**

(A) Neither a Partial Order nor an Equivalence Relation

(B) A Partial Order but not a Total Order

(C) A Total Order

(D) An Equivalence Relation

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q let R_1 be a relation from A = {1,3,5,7} to B = {2,4,6,8} and R_2 be another relation from B to C = {1,2,3,4} as defined below **(GATE-2004) (1 Marks)**

(i) an element x in A is related to an element y in B if x + y is divisible by 3

(ii) an element x in B is related to an element y in C if x + y is even but not divisible by 3.

Which is the composite relation R_1R_2 from A to C?

a) {(1,2), (1,4), (3,3), (5,4), (7,3)}

b) {(1,2), (1,3), (3,2), (5,2), (7,3)}

c) {(1,2), (3,2), (3,4), (5,4), (7,2)}

d) {(3,2), (3,4), (5,1), (5,3), (7,1)}

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q Let R be a symmetric and transitive relation on a set A. Then **(GATE-1995) (1 Marks)**

a) R is reflexive and hence an equivalence relation

b) R is reflexive and hence a partial order

c) R is reflexive and hence not an equivalence relation

d) None of the above

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Q.19 A relation R is said to be circular if aRb and bRc together imply cRa. Which of the following options is/are correct?

(Gate-2021) (1 Marks)

(a) If a relation S is reflexive and circular, then S is an equivalence relation.

(b) If a relation S is circular and symmetric, then S is an equivalence relation.

(c) If a relation S is transitive and circular, then S is an equivalence relation.

(d) If a relation S is reflexive and symmetric, then S is an equivalence relation.

Use Referral Code KGYT for Unacademy Plus(Live) to Get maximum Discount

Conversion of POSET into a Hasse Diagram

- If we want to study Partial order relation further then it will be better to convert it into more convenient notation so that it can be studied easily.

- This graphical representation is called Hasse Diagram

- In order theory, a Hasse diagram is a type of mathematical diagram used to represent a finite partially ordered set, in the form of a drawing of its transitive reduction.

- The diagrams are named after Helmut Hasse (1898–1979)



Steps to convert partial order relation into hasse diagram

1- Draw a vertex for each element in the Set

2- If (a, b) ϵ R then draw an edge from a to b

3- Remove all Reflexive and Transitive edges

4- Remove the direction of edges and arrange them in the increasing order of heights.

Q Consider a Partial order relation and convert it into hasse diagram?

R = {(1,1), (1,2), (1,4), (1,8), (2,2), (2,4), (2,8), (4,4), (4,8), (8,8)}

Q Consider a Partial order relation and convert it into hasse diagram?

R = {(1,1), (1,2), (1,3), (1,6), (2,2), (2,6), (3,3), (3,6), (6,6)}

Break

Q Study the following hasse diagram and find which of the following are valid?



(1)



(2)



(3)



(4)



(5)



(5)



(6)



(7)



(8)



(9)



(10)

Conclusion

- We can not have a horizontal edge in a hasse diagram
- We can not have a reflexive and transitive edge in Hasse Diagram

Q Let X = {2,3,6,12,24}, Let ≤ be the partial order defined by X ≤ Y if x divides y. Number of edges as in the Hasse diagram of (X, ≤) is. **(GATE-1996) (1 Marks)**

(a) 3 **(b)** 4 **(c)** 9 **(d)** None of the above

Break

Elements of a Poset

1. **Maximal Element: -** An element is said to be maximal if it is not related to any other element in the Partial order relation.

2. **Minimal Element: -** An element is said to be minimal if no other element is related to it in the Partial order relation.



	Elements
Minimal	
Least	
Maximal	
Greatest	



	Elements
Minimal	
Least	
Maximal	
Greatest	



	Elements
Minimal	
Least	
Maximal	
Greatest	



	Elements
Minimal	
Least	
Maximal	
Greatest	



	Elements
Minimal	
Least	
Maximal	
Greatest	



	Elements
Minimal	
Least	
Maximal	
Greatest	

- Every hasse diagram will have at least one Maximal and Minimal element(one or more).

Break

- **<u>Greatest Element</u>: -** An element is said to be Maximum/Greatest if it is not related to any other element but every element is related to it in Partial order relation. Or if a hasse diagram has only one Maximal element then it will also be Maximum/Greatest element.

- **<u>Least Element</u>: -** An element is said to be Minimum/Least if no other element is related to it but it is related to every element Partial order relation. Or if a hasse diagram has only one Minimal element then it will also be Minimum/Least element.



	Elements
Minimal	a
Least	
Maximal	i
Greatest	



	Elements
Minimal	a, b
Least	
Maximal	e, f
Greatest	



	Elements
Minimal	a
Least	
Maximal	f
Greatest	



	Elements
Minimal	a, c
Least	
Maximal	b, d
Greatest	



	Elements
Minimal	a
Least	
Maximal	c, d
Greatest	



	Elements
Minimal	c, d, f, i
Least	
Maximal	j, g, h, e
Greatest	

1. Every hasse diagram will have at most one Greatest and Least element(zero or one) (T/F)

2. Every Greatest element is also Maximal (T/F)

3. Every Least element is also Minimal (T/F)

4. If there is only one Maximal element then it is called Greatest (T/F)

5. If there is only one Minimal element then it is called Least (T/F)

Break

- **<u>Upper Bound</u>: -** Upper bound of a subset B with respect to set A, will contain all those element to which all the elements of B is related.

- **<u>Lower Bound</u>: -** lower bound of a subset B with respect to A, will contain all those elements which are related to every element of B.



Elements	B = {g, f, e, c}	B = {d, f}
Upper Bound		
Lower Bound		



Elements	B = {d, e}	B = {c}
Upper Bound		
Lower Bound		



Elements	B = {d, e}	B = {b, c}
Upper Bound		
Lower Bound		



Elements	B = {d, c}	B = {b, e}
Upper Bound		
Lower Bound		

Break

Least Upper Bound / LUB / Join / Supremum / $\vee$

Least value in the upper bound

Greatest Lower Bound / GLB / Meet / Infimum / $\wedge$

Greatest value in the lower bound



Elements	B = {g, f, e, c}	B = {d, f}
Upper Bound	{i}	{i, g, h, f}
Least Upper Bound		
Lower Bound	{a, e}	{a, d}
Greatest Lower Bound		



Elements	B = {d, e}	B = {c}
Upper Bound	{}	{d, e, c}
Least Upper Bound		
Lower Bound	{a, b, c}	{a, b, c}
Greatest Lower Bound		



Elements	B = {d, e}	B = {b, c}
Upper Bound	{f}	{d, e, f}
Least Upper Bound		
Lower Bound	{a, b, c}	{a}
Greatest Lower Bound		



Elements	B = {d, c}	B = {b, e}
Upper Bound	{f, d}	{f}
Least Upper Bound		
Lower Bound	{a, c}	{a}
Greatest Lower Bound		

Q Consider the Poset ({3,5,9,15,24,45}, /). Which of the following is correct for the given Poset? **(NET-JUNE-2019)**

a) There exist a greatest element and a least element

b) There exist a greatest element but not a least element

c) There exist a least element but not a greatest element

d) There does not exist a greatest element and a least element

Break

Join Semi Lattice :- A hasse diagram/Partial order relation is called Join Semi Lattice if for every elements their exists a Join.

Meet Semi Lattice :- A hasse diagram/Partial order relation is called Meet Semi Lattice if for every elements their exists a Meet.

<u>Lattice</u> :- A hasse diagram/Partial order relation is called Lattice if their exist a Join and Meet for every pair of element. Or A hasse diagram/Partial order relation is called Lattice if it is both Join Semi Lattice and Meet Semi Lattice.


e
f
c
d
a
b


e
f
d
b
c
a


i
g
h
b
f
c
d
e
a


d
e
c
a
b


f
e
d
c
b
a


f
d
e
b
c
a


f
d
e
c
a
b


d
e
b
c
a


g
e
f
d
b
c
a


d
b
c
a


f
b
c
d
e
a


d
c
e
b
a


f
d
e
b
c
a


g
e
f
d
b
c
a


e
c
d
a
b


c
d
b
a


i
g
h
d
e
f
b
c
a


d
f
e
b
c
a


d
c
b
a


b
a
c

a


b
d
a
c


f
d
e
b
c
a


g
e
f
d
b
c
a


h
e
f
g
b
c
d
a

Break

Q Consider the set **X**={a, b, c, d, e} under partial ordering **R**={(a, a), (a, b), (a, c), (a, d), (a, e), (b, b), (b, c), (b, e), (c, c), (c, e), (d, d), (d, e), (e, e)}. The Hasse diagram of the partial order (X, R) is shown below. The minimum number of ordered pairs that need to be added to R to make (X, R) a lattice is _____ **(GATE-2017) (1 Marks)**



Q A partially ordered set is said to be a lattice if every two elements in the set have **(NET-Dec-2010)**

a) a unique least upper bound

b) a unique greatest lower bound

c) both (A) and (B)

d) none of the above

Q Consider the following Hasse diagrams

Which all of the above represent a lattice? **(GATE-2008) (2 Marks)**

(A) (i) and (iv) only

(B) (ii) and (iii) only

(C) (iii) only

(D) (i), (ii) and (iv) only



Q the inclusion of which of the following set into S = {{1,2}, {1,2,3}, {1,3,5}, {1,2,4}, {1,2,3,4,5}} is necessary and sufficient to make S a complete lattice under the partial order defined by set containment?
(GATE-2004) (2 Marks)

a) {1}

b) {1}, {2,3}

c) {1}, {1,3}

d) {1}, {1,3}, {1,2,3,4}, {1,2,3,5}

Break

Boolean algebra

- **Unbounded Lattice :-** If a lattice has infinite of elements then it is called Unbounded Lattice.

- **<u>Bounded Lattice</u> :-** If a lattice has finite number of elements then it is called Bounded lattice, there will be upper and lower bound in lattice.

- **<u>Complement of an element in a Lattice</u> :-** If two elements a and a^c, are complement of each other, then the following equations must always holds good.

  $a \vee a^c$ = Upper bound of lattice

  $a \wedge a^c$ = Lower bound of lattice

- **<u>Distributive Lattice</u> :-** A lattice is said to be distributed lattice. if for every element their exist at most one completemt(zero or one).

- **<u>Complement Lattice</u> :-** A Lattice is said to be Complement lattice. if for every element their exist at least one complement(one or more).

- **<u>Boolean Algebra</u> :-** A Lattice is said to be Boolean Algebra, if for every element their exist exactly one complement. Or if a lattice is both complemented and distributed then it is called Boolean Algebra.


f
e
d
c
b
a


f
b
c
d
e
a


f
d
e
b
c
a


g
e
f
d
b
c
a


f
b
c
d
e
a


f
d
e
b
c
a


g
e
f
d
b
c
a


d
c
b
a


i
g
h
b
f
c
d
e
a


d
c
e
b
a


g
e
f
d
b
c
a


d
b
c
a


h
e
f
g
b
c
d
a

Break

Q The following is the Hasse diagram of the Poset [{a, b, c, d, e}, ≤]
The Poset is **(GATE-2005) (1 Marks)**

(A) not a lattice

(B) a lattice but not a distributive lattice

(C) a distributive lattice but not a Boolean algebra

(D) a Boolean algebra



Q The complement(s) of the element 'a' in the lattice shown in below figure is (are) _____

(GATE-1988) (2 Marks)



Q Find which of the following is a lattice and Boolean Algebra?

(1) $[D_{10}, /]$

(2) $[D_{12}, /]$

(3) $[D_{30}, /]$

(4) $[D_{45}, /]$

(5) $[D_{64}, /]$

(6) $[D_{81}, /]$

(7) $[D_{91}, /]$

(8) $[D_{110}, /]$

Q Find which of the following is a lattice and Boolean Algebra?

(1) [{1,2,3,4,6,9}, /]

(2) [{2,3,4,6,12}, /]

(3) [{1,2,3,5,30}, /]

(4) [{1,2,3,6,9,18}, /]

(5) [{2,3,4,9,12,18}, /]

(6) [R, <=]

(7) [P(A), ⊑], A = {1,2,3}

Q Consider the following hasse diagram, find which of the following is true?

a) it is a lattice

b) subset {a, b, c , d} is a lattice

c) subset {b, c, d, e} is a lattice

d) subset {a, b, c, e} is a lattice



Q Consider the following hasse diagram, find which of the following is true?

a) subset {a, b, c, g} is a lattice

b) subset {a, b, f, g} is a lattice

c) subset {a, d, e, g} is a lattice

d) subset {a, c, e, g} is a lattice



Break

Q Suppose L = {p, q, r, s, t} is a lattice represented by the following Hasse diagram: For any x, y ∈ L, not necessarily distinct, x ∨ y and x ∧ y are join and meet of x, y respectively. Let L^3 = {(x, y, z): x, y, z ∈ L} be the set of all ordered triplets of the elements of L. Let P_r be the probability that an element (x, y, z) ∈ L^3chosen equiprobably satisfies x ∨ (y ∧ z) = (x ∨ y) ∧ (x ∨ z). Then **(GATE-2015) (2 Marks)**

(A) Pr = 0 **(B)** Pr = 1 **(C)** 0 < Pr ≤ 1/5 **(D)** 1/5 < Pr < 1



Q Consider the set S = {a, b, c, d}.

Consider the following 4 partitions π_1, π_2, π_3, π_4 on S: π_1={~~abcd~~}, π_2={~~ab~~, ~~cd~~}, π_3={~~abc~~, ~~d~~}, π_4={~~a~~, ~~b~~, ~~c~~, ~~d~~}

Let $\prec$ be the partial order on the set of partitions $S'=\{\pi_1,\pi_2,\pi_3,\pi_4\}$ defined as follows: $\pi_i \prec \pi_j$ if and only if π_i refines π_j. The Poset diagram for $(S', \prec)$ is: **(GATE-2007) (2 Marks)**