

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

Aclasta 5 mg solution for infusion

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each bottle with 100 ml of solution contains 5 mg zoledronic acid (as monohydrate).

Each ml of the solution contains 0.05 mg zoledronic acid (as monohydrate).

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Solution for infusion

Clear and colourless solution.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Treatment of osteoporosis

- in post-menopausal women
- in adult men

at increased risk of fracture, including those with a recent low-trauma hip fracture.

Treatment of osteoporosis associated with long-term systemic glucocorticoid therapy

- in post-menopausal women
- in adult men

at increased risk of fracture.

Treatment of Paget's disease of the bone in adults.

### **4.2 Posology and method of administration**

#### Posology

Patients must be appropriately hydrated prior to administration of Aclasta. This is especially important for the elderly ( $\geq 65$  years) and for patients receiving diuretic therapy.

Adequate calcium and vitamin D intake are recommended in association with Aclasta administration.

#### Osteoporosis

For the treatment of post-menopausal osteoporosis, osteoporosis in men and the treatment of osteoporosis associated with long-term systemic glucocorticoid therapy, the recommended dose is a single intravenous infusion of 5 mg Aclasta administered once a year.

The optimal duration of bisphosphonate treatment for osteoporosis has not been established. The need for continued treatment should be re-evaluated periodically based on the benefits and potential risks of Aclasta on an individual patient basis, particularly after 5 or more years of use.

In patients with a recent low-trauma hip fracture, it is recommended to give the Aclasta infusion at least two weeks after hip fracture repair (see section 5.1). In patients with a recent low-trauma hip

fracture, a loading dose of 50 000 to 125 000 IU of vitamin D given orally or via the intramuscular route is recommended prior to the first Aclasta infusion.

#### Paget's disease

For the treatment of Paget's disease, Aclasta should be prescribed only by physicians with experience in the treatment of Paget's disease of the bone. The recommended dose is a single intravenous infusion of 5 mg Aclasta. In patients with Paget's disease, it is strongly advised that adequate supplemental calcium corresponding to at least 500 mg elemental calcium twice daily is ensured for at least 10 days following Aclasta administration (see section 4.4).

Re-treatment of Paget's disease: After initial treatment with Aclasta in Paget's disease, an extended remission period is observed in responding patients. Re-treatment consists of an additional intravenous infusion of 5 mg Aclasta after an interval of one year or longer from initial treatment in patients who have relapsed. Limited data on re-treatment of Paget's disease are available (see section 5.1).

#### Special populations

##### *Patients with renal impairment*

Aclasta is contraindicated in patients with creatinine clearance < 35 ml/min (see sections 4.3 and 4.4).

No dose adjustment is necessary in patients with creatinine clearance  $\geq$  35 ml/min.

##### *Patients with hepatic impairment*

No dose adjustment is required (see section 5.2).

##### *Elderly ( $\geq$ 65 years)*

No dose adjustment is necessary since bioavailability, distribution and elimination were similar in elderly patients and younger subjects.

##### *Paediatric population*

Aclasta should not be used in children and adolescents below 18 years of age. There are no data available for children under 5 years of age. Currently available data for children aged 5 to 17 years are described in section 5.1.

#### Method of administration

Intravenous use.

Aclasta is administered via a vented infusion line and given slowly at a constant infusion rate. The infusion time must not be less than 15 minutes. For information on the infusion of Aclasta, see section 6.6.

Patients treated with Aclasta should be given the package leaflet and the patient reminder card.

### **4.3 Contraindications**

- Hypersensitivity to the active substance, to any bisphosphonates or to any of the excipients listed in section 6.1.
- Patients with hypocalcaemia (see section 4.4).
- Severe renal impairment with creatinine clearance < 35 ml/min (see section 4.4).
- Pregnancy and breast-feeding (see section 4.6).

## 4.4 Special warnings and precautions for use

### Renal function

The use of Aclasta in patients with severe renal impairment (creatinine clearance < 35 ml/min) is contraindicated due to an increased risk of renal failure in this population.

Renal impairment has been observed following the administration of Aclasta (see section 4.8), especially in patients with pre-existing renal dysfunction or other risks including advanced age, concomitant nephrotoxic medicinal products, concomitant diuretic therapy (see section 4.5), or dehydration occurring after Aclasta administration. Renal impairment has been observed in patients after a single administration. Renal failure requiring dialysis or with a fatal outcome has rarely occurred in patients with underlying renal impairment or with any of the risk factors described above.

The following precautions should be taken into account to minimise the risk of renal adverse reactions:

- Creatinine clearance should be calculated based on actual body weight using the Cockcroft-Gault formula before each Aclasta dose.
- Transient increase in serum creatinine may be greater in patients with underlying impaired renal function.
- Monitoring of serum creatinine should be considered in at-risk patients.
- Aclasta should be used with caution when concomitantly used with other medicinal products that could impact renal function (see section 4.5).
- Patients, especially elderly patients and those receiving diuretic therapy, should be appropriately hydrated prior to administration of Aclasta.
- A single dose of Aclasta should not exceed 5 mg and the duration of infusion should be at least 15 minutes (see section 4.2).

### Hypocalcaemia

Pre-existing hypocalcaemia must be treated by adequate intake of calcium and vitamin D before initiating therapy with Aclasta (see section 4.3). Other disturbances of mineral metabolism must also be effectively treated (e.g. diminished parathyroid reserve, intestinal calcium malabsorption). Physicians should consider clinical monitoring for these patients.

Elevated bone turnover is a characteristic of Paget's disease of the bone. Due to the rapid onset of effect of zoledronic acid on bone turnover, transient hypocalcaemia, sometimes symptomatic, may develop and is usually maximal within the first 10 days after infusion of Aclasta (see section 4.8).

Adequate calcium and vitamin D intake are recommended in association with Aclasta administration. In addition, in patients with Paget's disease, it is strongly advised that adequate supplemental calcium corresponding to at least 500 mg elemental calcium twice daily is ensured for at least 10 days following Aclasta administration (see section 4.2).

Patients should be informed about symptoms of hypocalcaemia and receive adequate clinical monitoring during the period of risk. Measurement of serum calcium before infusion of Aclasta is recommended for patients with Paget's disease.

Severe and occasionally incapacitating bone, joint and/or muscle pain have been infrequently reported in patients taking bisphosphonates, including zoledronic acid (see section 4.8).

### Osteonecrosis of the jaw (ONJ)

ONJ has been reported in the post-marketing setting in patients receiving Aclasta (zoledronic acid) for osteoporosis (see section 4.8).

The start of treatment or of a new course of treatment should be delayed in patients with unhealed

open soft tissue lesions in the mouth. A dental examination with preventive dentistry and an individual benefit-risk assessment is recommended prior to treatment with Aclasta in patients with concomitant risk factors.

The following should be considered when evaluating a patient's risk of developing ONJ:

- Potency of the medicinal product that inhibits bone resorption (higher risk for highly potent compounds), route of administration (higher risk for parenteral administration) and cumulative dose of bone resorption therapy.
- Cancer, co-morbid conditions (e.g. anaemia, coagulopathies, infection), smoking.
- Concomitant therapies: corticosteroids, chemotherapy, angiogenesis inhibitors, radiotherapy to head and neck.
- Poor oral hygiene, periodontal disease, poorly fitting dentures, history of dental disease, invasive dental procedures, e.g. tooth extractions.

All patients should be encouraged to maintain good oral hygiene, undergo routine dental check-ups, and immediately report any oral symptoms such as dental mobility, pain or swelling, non-healing of sores or discharge during treatment with zoledronic acid. While on treatment, invasive dental procedures should be performed with caution and avoided in close proximity to zoledronic acid treatment.

The management plan for patients who develop ONJ should be set up in close collaboration between the treating physician and a dentist or oral surgeon with expertise in ONJ. Temporary interruption of zoledronic acid treatment should be considered until the condition resolves and contributing risk factors are mitigated where possible.

#### Osteonecrosis of the external auditory canal

Osteonecrosis of the external auditory canal has been reported with bisphosphonates, mainly in association with long-term therapy. Possible risk factors for osteonecrosis of the external auditory canal include steroid use and chemotherapy and/or local risk factors such as infection or trauma. The possibility of osteonecrosis of the external auditory canal should be considered in patients receiving bisphosphonates who present with ear symptoms including chronic ear infections.

#### Atypical fractures of the femur

Atypical subtrochanteric and diaphyseal femoral fractures have been reported with bisphosphonate therapy, primarily in patients receiving long-term treatment for osteoporosis. These transverse or short oblique fractures can occur anywhere along the femur from just below the lesser trochanter to just above the supracondylar flare. These fractures occur after minimal or no trauma and some patients experience thigh or groin pain, often associated with imaging features of stress fractures, weeks to months before presenting with a completed femoral fracture. Fractures are often bilateral; therefore the contralateral femur should be examined in bisphosphonate-treated patients who have sustained a femoral shaft fracture. Poor healing of these fractures has also been reported. Discontinuation of bisphosphonate therapy in patients suspected to have an atypical femur fracture should be considered pending evaluation of the patient, based on an individual benefit risk assessment.

During bisphosphonate treatment patients should be advised to report any thigh, hip or groin pain and any patient presenting with such symptoms should be evaluated for an incomplete femur fracture.

#### Acute phase reactions

Acute phase reactions (APRs) or post-dose symptoms such as fever, myalgia, flu-like symptoms, arthralgia and headache have been observed, the majority of which occurred within three days following Aclasta administration.

APRs may sometimes be serious or prolonged in duration. The incidence of post-dose symptoms can be reduced with the administration of paracetamol or ibuprofen shortly following Aclasta

administration. It is also advisable to postpone treatment if the patient is clinically unstable due to an acute medical condition and an APR could be problematic (see section 4.8).

### General

Other products containing zoledronic acid as an active substance are available for oncology indications. Patients being treated with Aclasta should not be treated with such products or any other bisphosphonate concomitantly, since the combined effects of these agents are unknown.

This medicinal product contains less than 1 mmol sodium (23 mg) per 100 ml vial of Aclasta, i.e. essentially “sodium free”.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies with other medicinal products have been performed. Zoledronic acid is not systemically metabolised and does not affect human cytochrome P450 enzymes *in vitro* (see section 5.2). Zoledronic acid is not highly bound to plasma proteins (approximately 43-55% bound) and interactions resulting from displacement of highly protein-bound medicinal products are therefore unlikely.

Zoledronic acid is eliminated by renal excretion. Caution is indicated when zoledronic acid is administered in conjunction with medicinal products that can significantly impact renal function (e.g. aminoglycosides or diuretics that may cause dehydration) (see section 4.4).

In patients with renal impairment, the systemic exposure to concomitant medicinal products that are primarily excreted via the kidney may increase.

## **4.6 Fertility, pregnancy and lactation**

### Women of childbearing potential

Aclasta is not recommended in women of childbearing potential.

### Pregnancy

Aclasta is contraindicated during pregnancy (see section 4.3). There are no adequate data on the use of zoledronic acid in pregnant women. Studies in animals with zoledronic acid have shown reproductive toxicological effects including malformations (see section 5.3). The potential risk for humans is unknown.

### Breast-feeding

Aclasta is contraindicated during breast-feeding (see section 4.3). It is unknown whether zoledronic acid is excreted into human milk.

### Fertility

Zoledronic acid was evaluated in rats for potential adverse effects on fertility of the parental and F1 generation. This resulted in exaggerated pharmacological effects considered related to the compound's inhibition of skeletal calcium mobilisation, resulting in periparturient hypocalcaemia, a bisphosphonate class effect, dystocia and early termination of the study. Thus these results precluded determining a definitive effect of Aclasta on fertility in humans.

## **4.7 Effects on ability to drive and use machines**

Adverse reactions, such as dizziness, may affect the ability to drive or use machines.

## 4.8 Undesirable effects

### Summary of the safety profile

The overall percentage of patients who experienced adverse reactions were 44.7%, 16.7% and 10.2% after the first, second and third infusion, respectively. Incidence of individual adverse reactions following the first infusion was: pyrexia (17.1%), myalgia (7.8%), influenza-like illness (6.7%), arthralgia (4.8%) and headache (5.1%), see “acute phase reactions” below.

### Tabulated list of adverse reactions

Adverse reactions in Table 1 are listed according to MedDRA system organ class and frequency category. Frequency categories are defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

**Table 1**

<b><i>Infections and infestations</i></b>	<i>Uncommon</i>	Influenza, nasopharyngitis
<b><i>Blood and lymphatic system disorders</i></b>	<i>Uncommon</i>	Anaemia
<b><i>Immune system disorders</i></b>	<i>Not known**</i>	Hypersensitivity reactions including rare cases of bronchospasm, urticaria and angioedema, and very rare cases of anaphylactic reaction/shock
<b><i>Metabolism and nutrition disorders</i></b>	<i>Common</i>	Hypocalcaemia*
	<i>Uncommon</i>	Decreased appetite
	<i>Rare</i>	Hypophosphataemia
<b><i>Psychiatric disorders</i></b>	<i>Uncommon</i>	Insomnia
<b><i>Nervous system disorders</i></b>	<i>Common</i>	Headache, dizziness
	<i>Uncommon</i>	Lethargy, paraesthesia, somnolence, tremor, syncope, dysgeusia
<b><i>Eye disorders</i></b>	<i>Common</i>	Ocular hyperaemia
	<i>Uncommon</i>	Conjunctivitis, eye pain
	<i>Rare</i>	Uveitis, episcleritis, iritis
	<i>Not known**</i>	Scleritis and parophthalmia
<b><i>Ear and labyrinth disorders</i></b>	<i>Uncommon</i>	Vertigo
<b><i>Cardiac disorders</i></b>	<i>Common</i>	Atrial fibrillation
	<i>Uncommon</i>	Palpitations
<b><i>Vascular disorders</i></b>	<i>Uncommon</i>	Hypertension, flushing
	<i>Not known**</i>	Hypotension (some of the patients had underlying risk factors)
<b><i>Respiratory, thoracic and mediastinal disorders</i></b>	<i>Uncommon</i>	Cough, dyspnoea
<b><i>Gastrointestinal disorders</i></b>	<i>Common</i>	Nausea, vomiting, diarrhoea
	<i>Uncommon</i>	Dyspepsia, abdominal pain upper, abdominal pain, gastro-oesophageal reflux disease, constipation, dry mouth, oesophagitis, toothache, gastritis <sup>#</sup>
<b><i>Skin and subcutaneous tissue disorders</i></b>	<i>Uncommon</i>	Rash, hyperhidrosis, pruritus, erythema

<b><i>Musculoskeletal and connective tissue disorders</i></b>	<i>Common</i>	Myalgia, arthralgia, bone pain, back pain, pain in extremity
	<i>Uncommon</i>	Neck pain, musculoskeletal stiffness, joint swelling, muscle spasms, musculoskeletal chest pain, musculoskeletal pain, joint stiffness, arthritis, muscular weakness
	<i>Rare</i>	Atypical subtrochanteric and diaphyseal femoral fractures† (bisphosphonate class adverse reaction)
	<i>Very rare</i>	Osteonecrosis of the external auditory canal (bisphosphonate class adverse reaction)
	<i>Not known**</i>	Osteonecrosis of the jaw (see sections 4.4 and 4.8 Class effects)
<b><i>Renal and urinary disorders</i></b>	<i>Uncommon</i>	Blood creatinine increased, pollakiuria, proteinuria
	<i>Not known**</i>	Renal impairment. Rare cases of renal failure requiring dialysis and rare cases with a fatal outcome have been reported in patients with pre-existing renal dysfunction or other risk factors such as advanced age, concomitant nephrotoxic medicinal products, concomitant diuretic therapy, or dehydration in the post infusion period (see sections 4.4 and 4.8 Class effects)
<b><i>General disorders and administration site conditions</i></b>	<i>Very common</i>	Pyrexia
	<i>Common</i>	Influenza-like illness, chills, fatigue, asthenia, pain, malaise, infusion site reaction
	<i>Uncommon</i>	Peripheral oedema, thirst, acute phase reaction, non-cardiac chest pain
	<i>Not known**</i>	Dehydration secondary to acute phase reactions (post-dose symptoms such as pyrexia, vomiting and diarrhoea)
<b><i>Investigations</i></b>	<i>Common</i>	C-reactive protein increased
	<i>Uncommon</i>	Blood calcium decreased

# Observed in patients taking concomitant glucocorticosteroids.

\* Common in Paget's disease only.

\*\* Based on post-marketing reports. Frequency cannot be estimated from available data.

† Identified in post-marketing experience.

#### Description of selected adverse reactions

##### Atrial fibrillation

In the HORIZON – Pivotal Fracture Trial [PFT] (see section 5.1), the overall incidence of atrial fibrillation was 2.5% (96 out of 3,862) and 1.9% (75 out of 3,852) in patients receiving Aclasta and placebo, respectively. The rate of atrial fibrillation serious adverse events was increased in patients receiving Aclasta (1.3%) (51 out of 3,862) compared with patients receiving placebo (0.6%) (22 out of 3,852). The mechanism behind the increased incidence of atrial fibrillation is unknown. In the osteoporosis trials (PFT, HORIZON - Recurrent Fracture Trial [RFT]) the pooled atrial fibrillation incidences were comparable between Aclasta (2.6%) and placebo (2.1%). For atrial fibrillation serious adverse events the pooled incidences were 1.3% for Aclasta and 0.8% for placebo.



### Class effects

#### *Renal impairment*

Zoledronic acid has been associated with renal impairment manifested as deterioration in renal function (i.e. increased serum creatinine) and in rare cases acute renal failure. Renal impairment has been observed following the administration of zoledronic acid, especially in patients with pre-existing renal dysfunction or additional risk factors (e.g advanced age, oncology patients with chemotherapy, concomitant nephrotoxic medicinal products, concomitant diuretic therapy, severe dehydration), with the majority of them receiving a 4 mg dose every 3–4 weeks, but it has been observed in patients after a single administration.

In clinical trials in osteoporosis, the change in creatinine clearance (measured annually prior to dosing) and the incidence of renal failure and impairment was comparable for both the Aclasta and placebo treatment groups over three years. There was a transient increase in serum creatinine observed within 10 days in 1.8% of Aclasta-treated patients versus 0.8% of placebo-treated patients.

#### *Hypocalcaemia*

In clinical trials in osteoporosis, approximately 0.2% of patients had notable declines of serum calcium levels (less than 1.87 mmol/l) following Aclasta administration. No symptomatic cases of hypocalcaemia were observed.

In the Paget's disease trials, symptomatic hypocalcaemia was observed in approximately 1% of patients, in all of whom it resolved.

Based on laboratory assessment, transient asymptomatic calcium levels below the normal reference range (less than 2.10 mmol/l) occurred in 2.3% of Aclasta-treated patients in a large clinical trial compared to 21% of Aclasta-treated patients in the Paget's disease trials. The frequency of hypocalcaemia was much lower following subsequent infusions.

All patients received adequate supplementation with vitamin D and calcium in the post-menopausal osteoporosis trial, the prevention of clinical fractures after hip fracture trial, and the Paget's disease trials (see also section 4.2). In the trial for the prevention of clinical fractures following a recent hip fracture, vitamin D levels were not routinely measured but the majority of patients received a loading dose of vitamin D prior to Aclasta administration (see section 4.2).

#### *Local reactions*

In a large clinical trial, local reactions at the infusion site, such as redness, swelling and/or pain, were reported (0.7%) following the administration of zoledronic acid.

#### *Osteonecrosis of the jaw*

Cases of osteonecrosis of the jaw have been reported, predominantly in cancer patients treated with medicinal products that inhibit bone resorption, including zoledronic acid (see section 4.4). In a large clinical trial in 7,736 patients, osteonecrosis of the jaw has been reported in one patient treated with Aclasta and one patient treated with placebo. Cases of ONJ have been reported in the post-marketing setting for Aclasta.

#### *Acute phase reactions*

The overall percentage of patients who reported acute phase reactions or post-dose symptoms (including serious cases) after Aclasta administration is as follows (frequencies derived from the study in treatment of post-menopausal osteoporosis): fever (18.1%), myalgia (9.4%), flu-like symptoms (7.8%), arthralgia (6.8%) and headache (6.5%), the majority of which occurred within the first 3 days following Aclasta administration. The majority of these symptoms were mild to moderate in nature and resolved within 3 days of the event onset. The incidence of these symptoms decreased with subsequent annual doses of Aclasta. The percentage of patients who experienced adverse reactions was lower in a smaller study (19.5%, 10.4%, 10.7% after the first, second and third infusion, respectively), where prophylaxis against adverse reactions was used (see section 4.4).

## Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## **4.9 Overdose**

Clinical experience with acute overdose is limited. Patients who have received doses higher than those recommended should be carefully monitored. In the event of overdose leading to clinically significant hypocalcaemia, reversal may be achieved with supplemental oral calcium and/or an intravenous infusion of calcium gluconate.

# **5. PHARMACOLOGICAL PROPERTIES**

## **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for treatment of bone diseases, bisphosphonates, ATC code: M05BA08

### Mechanism of action

Zoledronic acid belongs to the class of nitrogen-containing bisphosphonates and acts primarily on bone. It is an inhibitor of osteoclast-mediated bone resorption.

### Pharmacodynamic effects

The selective action of bisphosphonates on bone is based on their high affinity for mineralised bone.

The main molecular target of zoledronic acid in the osteoclast is the enzyme farnesyl pyrophosphate synthase. The long duration of action of zoledronic acid is attributable to its high binding affinity for the active site of farnesyl pyrophosphate (FPP) synthase and its strong binding affinity to bone mineral.

Aclasta treatment rapidly reduced the rate of bone turnover from elevated post-menopausal levels with the nadir for resorption markers observed at 7 days, and for formation markers at 12 weeks. Thereafter bone markers stabilised within the pre-menopausal range. There was no progressive reduction of bone turnover markers with repeated annual dosing.

### Clinical efficacy in the treatment of post-menopausal osteoporosis (PFT)

The efficacy and safety of Aclasta 5 mg once a year for 3 consecutive years were demonstrated in post-menopausal women (7,736 women aged 65–89 years) with either: a femoral neck bone mineral density (BMD) with a T-score  $\leq -1.5$  and at least two mild or one moderate existing vertebral fracture(s); or a femoral neck BMD T-score  $\leq -2.5$  with or without evidence of existing vertebral fracture(s). 85% of patients were bisphosphonate-naïve. Women who were evaluated for the incidence of vertebral fractures did not receive concomitant osteoporosis therapy, which was allowed for women contributing to the hip and all clinical fracture evaluations. Concomitant osteoporosis therapy included: calcitonin, raloxifene, tamoxifen, hormone replacement therapy, tibolone; but excluded other bisphosphonates. All women received 1,000 to 1,500 mg elemental calcium and 400 to 1,200 IU of vitamin D supplements daily.

### Effect on morphometric vertebral fractures

Aclasta significantly decreased the incidence of one or more new vertebral fractures over three years and as early as the one year timepoint (see Table 2).

**Table 2 Summary of vertebral fracture efficacy at 12, 24 and 36 months**

Outcome	Aclasta (%)	Placebo (%)	Absolute reduction in fracture incidence % (CI)	Relative reduction in fracture incidence % (CI)
At least one new vertebral fracture (0–1 year)	1.5	3.7	2.2 (1.4, 3.1)	60 (43, 72)**
At least one new vertebral fracture (0–2 year)	2.2	7.7	5.5 (4.4, 6.6)	71 (62, 78)**
At least one new vertebral fracture (0–3 year)	3.3	10.9	7.6 (6.3, 9.0)	70 (62, 76)**
** p <0.0001				

Aclasta-treated patients aged 75 years and older exhibited a 60% reduction in the risk of vertebral fractures compared to placebo patients (p<0.0001).

#### Effect on hip fractures

Aclasta demonstrated a consistent effect over 3 years, resulting in a 41% reduction in the risk of hip fractures (95% CI, 17% to 58%). The hip fracture event rate was 1.44% for Aclasta-treated patients compared to 2.49% for placebo-treated patients. The risk reduction was 51% in bisphosphonate-naïve patients and 42% in patients allowed to take concomitant osteoporosis therapy.

#### Effect on all clinical fractures

All clinical fractures were verified based on the radiographic and/or clinical evidence. A summary of results is presented in Table 3.

**Table 3 Between treatment comparisons of the incidence of key clinical fracture variables over 3 years**

Outcome	Aclasta (N=3,875) event rate (%)	Placebo (N=3,861) event rate (%)	Absolute reduction in fracture event rate % (CI)	Relative risk reduction in fracture incidence % (CI)
Any clinical fracture (1)	8.4	12.8	4.4 (3.0, 5.8)	33 (23, 42)**
Clinical vertebral fracture (2)	0.5	2.6	2.1 (1.5, 2.7)	77 (63, 86)**
Non-vertebral fracture (1)	8.0	10.7	2.7 (1.4, 4.0)	25 (13, 36)*
*p-value <0.001, **p-value <0.0001				
(1) Excluding finger, toe and facial fractures				
(2) Including clinical thoracic and clinical lumbar vertebral fractures				

#### Effect on bone mineral density (BMD)

Aclasta significantly increased BMD at the lumbar spine, hip, and distal radius relative to treatment with placebo at all timepoints (6, 12, 24 and 36 months). Treatment with Aclasta resulted in a 6.7% increase in BMD at the lumbar spine, 6.0% at the total hip, 5.1% at the femoral neck, and 3.2% at the distal radius over 3 years as compared to placebo.

#### Bone histology

Bone biopsies were obtained from the iliac crest 1 year after the third annual dose in 152 post-menopausal patients with osteoporosis treated with Aclasta (N=82) or placebo (N=70). Histomorphometric analysis showed a 63% reduction in bone turnover. In patients treated with Aclasta, no osteomalacia, marrow fibrosis or woven bone formation was detected. Tetracycline label was detectable in all but one of 82 biopsies obtained from patients on Aclasta. Microcomputed tomography (μCT) analysis demonstrated increased trabecular bone volume and preservation of trabecular bone architecture in patients treated with Aclasta compared to placebo.

### Bone turnover markers

Bone specific alkaline phosphatase (BSAP), serum N-terminal propeptide of type I collagen (P1NP) and serum beta-C-telopeptides (b-CTx) were evaluated in subsets ranging from 517 to 1,246 patients at periodic intervals throughout the study. Treatment with a 5 mg annual dose of Aclasta significantly reduced BSAP by 30% relative to baseline at 12 months which was sustained at 28% below baseline levels at 36 months. P1NP was significantly reduced by 61% below baseline levels at 12 months and was sustained at 52% below baseline levels at 36 months. B-CTx was significantly reduced by 61% below baseline levels at 12 months and was sustained at 55% below baseline levels at 36 months. During this entire time period bone turnover markers were within the pre-menopausal range at the end of each year. Repeat dosing did not lead to further reduction of bone turnover markers.

### Effect on height

In the three-year osteoporosis study standing height was measured annually using a stadiometer. The Aclasta group revealed approximately 2.5 mm less height loss compared to placebo (95% CI: 1.6 mm, 3.5 mm) [ $p < 0.0001$ ].

### Days of disability

Aclasta significantly reduced the mean days of limited activity and the days of bed rest due to back pain by 17.9 days and 11.3 days respectively compared to placebo and significantly reduced the mean days of limited activity and the days of bed rest due to fractures by 2.9 days and 0.5 days respectively compared to placebo (all  $p < 0.01$ ).

### Clinical efficacy in the treatment of osteoporosis in patients at increased risk of fracture after a recent hip fracture (RFT)

The incidence of clinical fractures, including vertebral, non-vertebral and hip fractures, was evaluated in 2,127 men and women aged 50-95 years (mean age 74.5 years) with a recent (within 90 days) low-trauma hip fracture who were followed for an average of 2 years on study treatment (Aclasta). Approximately 42% of patients had a femoral neck BMD T-score below -2.5 and approximately 45% of the patients had a femoral neck BMD T-score above -2.5. Aclasta was administered once a year, until at least 211 patients in the study population had confirmed clinical fractures. Vitamin D levels were not routinely measured but a loading dose of vitamin D (50,000 to 125,000 IU orally or by intramuscular route) was given to the majority of patients 2 weeks prior to infusion. All participants received 1,000 to 1,500 mg of elemental calcium plus 800 to 1,200 IU of vitamin D supplementation per day. Ninety-five percent of the patients received their infusion two or more weeks after the hip fracture repair and the median timing of infusion was approximately six weeks after the hip fracture repair. The primary efficacy variable was the incidence of clinical fractures over the duration of the study.

### Effect on all clinical fractures

The incidence rates of key clinical fracture variables are presented in Table 4.

**Table 4 Between treatment comparisons of the incidence of key clinical fracture variables**

Outcome	Aclasta (N=1,065) event rate (%)	Placebo (N=1,062) event rate (%)	Absolute reduction in fracture event rate % (CI)	Relative risk reduction in fracture incidence % (CI)
Any clinical fracture (1)	8.6	13.9	5.3 (2.3, 8.3)	35 (16, 50)**
Clinical vertebral fracture (2)	1.7	3.8	2.1 (0.5, 3.7)	46 (8, 68)*
Non-vertebral fracture (1)	7.6	10.7	3.1 (0.3, 5.9)	27 (2, 45)*
*p-value <0.05, **p-value <0.01 (1) Excluding finger, toe and facial fractures (2) Including clinical thoracic and clinical lumbar vertebral fractures				

The study was not designed to measure significant differences in hip fracture, but a trend was seen

towards reduction in new hip fractures.

All cause mortality was 10% (101 patients) in the Aclasta-treated group compared to 13% (141 patients) in the placebo group. This corresponds to a 28% reduction in the risk of all cause mortality ( $p=0.01$ ).

The incidence of delayed hip fracture healing was comparable between Aclasta (34 [3.2%]) and placebo (29 [2.7%]).

#### Effect on bone mineral density (BMD)

In the HORIZON-RFT study Aclasta treatment significantly increased BMD at the total hip and femoral neck relative to treatment with placebo at all timepoints. Treatment with Aclasta resulted in an increase in BMD of 5.4% at the total hip and 4.3% at the femoral neck over 24 months as compared to placebo.

#### Clinical efficacy in men

In the HORIZON-RFT study 508 men were randomised into the study and 185 patients had BMD assessed at 24 months. At 24 months a similar significant increase of 3.6% in total hip BMD was observed for patients treated with Aclasta as compared to the effects observed in post-menopausal women in the HORIZON-PFT study. The study was not powered to show a reduction in clinical fractures in men; the incidence of clinical fractures was 7.5% in men treated with Aclasta versus 8.7% for placebo.

In another study in men (study CZOL446M2308) an annual infusion of Aclasta was non-inferior to weekly alendronate for the percentage change in lumbar spine BMD at month 24 relative to baseline.

#### Clinical efficacy in osteoporosis associated with long-term systemic glucocorticoid therapy

The efficacy and safety of Aclasta in the treatment and prevention of osteoporosis associated with long-term systemic glucocorticoid therapy were assessed in a randomised, multicentre, double-blind, stratified, active-controlled study of 833 men and women aged 18-85 years (mean age for men 56.4 years; for women 53.5 years) treated with  $> 7.5$  mg/day oral prednisone (or equivalent). Patients were stratified with respect to duration of glucocorticoid use prior to randomisation ( $\leq 3$  months versus  $> 3$  months). The duration of the trial was one year. Patients were randomised to either Aclasta 5 mg single infusion or to oral risedronate 5 mg daily for one year. All participants received 1,000 mg elemental calcium plus 400 to 1,000 IU vitamin D supplementation per day. Efficacy was demonstrated if non-inferiority to risedronate was shown sequentially with respect to the percentage change in lumbar spine BMD at 12 months relative to baseline in the treatment and prevention subpopulations, respectively. The majority of patients continued to receive glucocorticoids for the one year duration of the trial.

#### Effect on bone mineral density (BMD)

The increases in BMD were significantly greater in the Aclasta-treated group at the lumbar spine and femoral neck at 12 months compared to risedronate (all  $p<0.03$ ). In the subpopulation of patients receiving glucocorticoids for more than 3 months prior to randomisation, Aclasta increased lumbar spine BMD by 4.06% versus 2.71% for risedronate (mean difference: 1.36% ;  $p<0.001$ ). In the subpopulation of patients that had received glucocorticoids for 3 months or less prior to randomisation, Aclasta increased lumbar spine BMD by 2.60% versus 0.64% for risedronate (mean difference: 1.96% ;  $p<0.001$ ). The study was not powered to show a reduction in clinical fractures compared to risedronate. The incidence of fractures was 8 for Aclasta-treated patients versus 7 for risedronate-treated patients ( $p=0.8055$ ).

#### Clinical efficacy in the treatment of Paget's disease of the bone

Aclasta was studied in male and female patients aged above 30 years with primarily mild to moderate Paget's disease of the bone (median serum alkaline phosphatase level 2.6–3.0 times the upper limit of

the age-specific normal reference range at the time of study entry) confirmed by radiographic evidence.

The efficacy of one infusion of 5 mg zoledronic acid versus daily doses of 30 mg risedronate for 2 months was demonstrated in two 6-month comparative trials. After 6 months, Aclasta showed 96% (169/176) and 89% (156/176) response and serum alkaline phosphatase (SAP) normalisation rates compared to 74% (127/171) and 58% (99/171) for risedronate (all  $p < 0.001$ ).

In the pooled results, a similar decrease in pain severity and pain interference scores relative to baseline were observed over 6 months for Aclasta and risedronate.

Patients who were classified as responders at the end of the 6 month core study were eligible to enter an extended follow-up period. Of the 153 Aclasta-treated patients and 115 risedronate-treated patients who entered an extended observation study, after a mean duration of follow-up of 3.8 years from time of dosing, the proportion of patients ending the Extended Observation Period due to the need for re-treatment (clinical judgment) was higher for risedronate (48 patients, or 41.7%) compared with zoledronic acid (11 patients, or 7.2%). The mean time of ending the Extended Observation Period due to the need for Paget's re-treatment from the initial dose was longer for zoledronic acid (7.7 years) than for risedronate (5.1 years).

Six patients who achieved therapeutic response 6 months after treatment with Aclasta and later experienced disease relapse during the extended follow-up period were re-treated with Aclasta after a mean time of 6.5 years from initial treatment to re-treatment. Five of the 6 patients had SAP within the normal range at month 6 (Last Observation Carried Forward, LOCF).

Bone histology was evaluated in 7 patients with Paget's disease 6 months after treatment with 5 mg zoledronic acid. Bone biopsy results showed bone of normal quality with no evidence of impaired bone remodelling and no evidence of mineralisation defects. These results were consistent with biochemical marker evidence of normalisation of bone turnover.

### Paediatric population

A randomised, double-blind, placebo-controlled study was conducted in paediatric patients aged 5 to 17 years treated with glucocorticoids who had decreased bone mineral density (lumbar spine BMD Z-score of -0.5 or less) and a low impact/fragility fracture. The patient population randomised in this study (ITT population) included patients with several sub-types of rheumatic conditions, inflammatory bowel disease, or Duchenne muscular dystrophy. The study was planned to include 92 patients, however only 34 patients were enrolled and randomised to receive either a twice-yearly 0.05 mg/kg (max. 5 mg) intravenous zoledronic acid infusion or placebo for one year. All patients were required to receive background therapy of vitamin D and calcium.

Zoledronic acid infusion resulted in an increase in the lumbar spine BMD Z-score least square (LS) mean difference of 0.41 at month 12 relative to baseline compared to placebo (95% CI: 0.02, 0.81; 18 and 16 patients, respectively). No effect on lumbar spine BMD Z-score was evident after 6 months of treatment. At month 12, a statistically significant ( $p < 0.05$ ) reduction in three bone turnover markers (PINP, BSAP, NTX) was observed in the zoledronic acid group as compared to the placebo group. No statistically significant differences in total body bone mineral content were observed between patients treated with zoledronic acid versus placebo at 6 or 12 months. There is no clear evidence establishing a link between BMD changes and fracture prevention in children with growing skeletons.

No new vertebral fractures were observed in the zoledronic acid group as compared to two new fractures in the placebo group.

The most commonly reported adverse reactions after infusion of zoledronic acid were arthralgia (28%), pyrexia (22%), vomiting (22%), headache (22%), nausea (17%), myalgia (17%), pain (17%), diarrhoea (11%) and hypocalcaemia (11%).

More patients reported serious adverse events in the zoledronic acid group than in the placebo group (5 [27.8%] patients versus 1 [6.3%] patient).

In the 12-month open-label extension of the above-mentioned core study, no new clinical fractures were observed. However 2 patients, one in each of the core study treatment groups (zoledronic acid group: 1/9, 11.1% and placebo group: 1/14, 7.1%), had new morphometric vertebral fractures. There were no new safety findings.

Long-term safety data in this population cannot be established from these studies.

The European Medicines Agency has waived the obligation to submit the results of studies with Aclasta in all subsets of the paediatric population in Paget's disease of the bone, osteoporosis in post-menopausal women at an increased risk of fracture, osteoporosis in men at increased risk of fracture and prevention of clinical fractures after a hip fracture in men and women (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Single and multiple 5 and 15-minute infusions of 2, 4, 8 and 16 mg zoledronic acid in 64 patients yielded the following pharmacokinetic data, which were found to be dose independent.

### Distribution

After initiation of the zoledronic acid infusion, plasma concentrations of the active substance increased rapidly, achieving their peak at the end of the infusion period, followed by a rapid decline to < 10% of peak after 4 hours and < 1% of peak after 24 hours, with a subsequent prolonged period of very low concentrations not exceeding 0.1% of peak levels.

### Elimination

Intravenously administered zoledronic acid is eliminated by a triphasic process: rapid biphasic disappearance from the systemic circulation, with half-lives of  $t_{1/2\alpha}$  0.24 and  $t_{1/2\beta}$  1.87 hours, followed by a long elimination phase with a terminal elimination half-life of  $t_{1/2\gamma}$  146 hours. There was no accumulation of the active substance in plasma after multiple doses given every 28 days. The early disposition phases ( $\alpha$  and  $\beta$ , with  $t_{1/2}$  values above) presumably represent rapid uptake into bone and excretion via the kidneys.

Zoledronic acid is not metabolised and is excreted unchanged via the kidney. Over the first 24 hours,  $39 \pm 16\%$  of the administered dose is recovered in the urine, while the remainder is principally bound to bone tissue. This uptake into bone is common for all bisphosphonates and is presumably a consequence of the structural analogy to pyrophosphate. As with other bisphosphonates, the retention time of zoledronic acid in bones is very long. From the bone tissue it is released very slowly back into the systemic circulation and eliminated via the kidney. The total body clearance is  $5.04 \pm 2.5$  l/h, independent of dose, and unaffected by gender, age, race or body weight. The inter- and intra-subject variation for plasma clearance of zoledronic acid was shown to be 36% and 34%, respectively. Increasing the infusion time from 5 to 15 minutes caused a 30% decrease in zoledronic acid concentration at the end of the infusion, but had no effect on the area under the plasma concentration versus time curve.

### Pharmacokinetic/pharmacodynamic relationships

No interaction studies with other medicinal products have been performed with zoledronic acid. Since zoledronic acid is not metabolised in humans and the substance was found to have little or no capacity as a direct-acting and/or irreversible metabolism-dependent inhibitor of P450 enzymes, zoledronic acid is unlikely to reduce the metabolic clearance of substances which are metabolised via the cytochrome P450 enzyme systems. Zoledronic acid is not highly bound to plasma proteins (approximately 43-55% bound) and binding is concentration independent. Therefore, interactions

resulting from displacement of highly protein-bound medicinal products are unlikely.

#### Special populations (see section 4.2)

##### Renal impairment

The renal clearance of zoledronic acid was correlated with creatinine clearance, renal clearance representing  $75 \pm 33\%$  of the creatinine clearance, which showed a mean of  $84 \pm 29$  ml/min (range 22 to 143 ml/min) in the 64 patients studied. Small observed increases in  $AUC_{(0-24hr)}$ , by about 30% to 40% in mild to moderate renal impairment, compared to a patient with normal renal function, and lack of accumulation of drug with multiple doses irrespective of renal function, suggest that dose adjustments of zoledronic acid in mild ( $Cl_{cr} = 50-80$  ml/min) and moderate renal impairment down to a creatinine clearance of 35 ml/min are not necessary. The use of Aclasta in patients with severe renal impairment (creatinine clearance  $< 35$  ml/min) is contraindicated due to an increased risk of renal failure in this population.

### **5.3 Preclinical safety data**

#### Acute toxicity

The highest non-lethal single intravenous dose was 10 mg/kg body weight in mice and 0.6 mg/kg in rats. In the single-dose dog infusion studies, 1.0 mg/kg (6 fold the recommended human therapeutic exposure based on AUC) administered over 15 minutes was well tolerated with no renal effects.

#### Subchronic and chronic toxicity

In the intravenous infusion studies, renal tolerability of zoledronic acid was established in rats when given 0.6 mg/kg as 15-minute infusions at 3-day intervals, six times in total (for a cumulative dose that corresponded to AUC levels about 6 times the human therapeutic exposure) while five 15-minute infusions of 0.25 mg/kg administered at 2–3-week intervals (a cumulative dose that corresponded to 7 times the human therapeutic exposure) were well tolerated in dogs. In the intravenous bolus studies, the doses that were well-tolerated decreased with increasing study duration: 0.2 and 0.02 mg/kg daily was well tolerated for 4 weeks in rats and dogs, respectively but only 0.01 mg/kg and 0.005 mg/kg in rats and dogs, respectively, when given for 52 weeks.

Longer-term repeat administration at cumulative exposures sufficiently exceeding the maximum intended human exposure produced toxicological effects in other organs, including the gastrointestinal tract and liver, and at the site of intravenous administration. The clinical relevance of these findings is unknown. The most frequent finding in the repeat-dose studies consisted of increased primary spongiosa in the metaphyses of long bones in growing animals at nearly all doses, a finding that reflected the compound's pharmacological antiresorptive activity.

#### Reproduction toxicity

Teratology studies were performed in two species, both via subcutaneous administration. Teratogenicity was observed in rats at doses  $\geq 0.2$  mg/kg and was manifested by external, visceral and skeletal malformations. Dystocia was observed at the lowest dose (0.01 mg/kg body weight) tested in rats. No teratological or embryo/foetal effects were observed in rabbits, although maternal toxicity was marked at 0.1 mg/kg due to decreased serum calcium levels.

#### Mutagenicity and carcinogenic potential

Zoledronic acid was not mutagenic in the mutagenicity tests performed and carcinogenicity testing did not provide any evidence of carcinogenic potential.



## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Mannitol  
Sodium citrate  
Water for injections

### **6.2 Incompatibilities**

This medicinal product must not be allowed to come into contact with any calcium-containing solutions. Aclasta must not be mixed or given intravenously with any other medicinal products.

### **6.3 Shelf life**

Unopened bottle: 3 years

After opening: 24 hours at 2°C - 8°C

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.  
For storage conditions after first opening of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

100 ml solution in a transparent plastic (cycloolefinic polymer) bottle closed with a fluoro-polymer coated bromobutyl rubber stopper and an aluminium/polypropylene cap with a flip component.

Aclasta is supplied in packs containing one bottle as unit pack, or in multipacks comprising five packs, each containing one bottle.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

For single use only.  
Only clear solution free from particles and discoloration should be used.

If refrigerated, allow the refrigerated solution to reach room temperature before administration. Aseptic techniques must be followed during the preparation of the infusion.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/05/308/001

EU/1/05/308/002

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 15 April 2005

Date of latest renewal: 19 January 2015

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**

### Name and address of the manufacturer responsible for batch release

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

The MAH shall ensure that the educational programme implemented for the authorised indications of treatment of osteoporosis in post-menopausal women and in men at increased risk of fracture, including those with a recent low-trauma hip fracture, and treatment of osteoporosis associated with long-term systemic glucocorticoid therapy in post-menopausal women and in men at increased risk of fracture is updated. The following documents should be included in the patient information pack:

- Package leaflet
- Patient reminder card on osteonecrosis of the jaw

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****CARTON (WITH BLUE BOX) FOR UNIT PACK****1. NAME OF THE MEDICINAL PRODUCT**

Aclasta 5 mg solution for infusion  
zoledronic acid

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each bottle of 100 ml contains 5 mg zoledronic acid (as monohydrate).

**3. LIST OF EXCIPIENTS**

Mannitol, sodium citrate and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for infusion

1 bottle of 100 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For single use only.  
Read the package leaflet before use.  
Intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
After opening: 24 hours at 2°C - 8°C.

**9. SPECIAL STORAGE CONDITIONS**

<b>10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE</b>
--

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

<b>11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER</b>
---

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

<b>12. MARKETING AUTHORISATION NUMBER(S)</b>
--

EU/1/05/308/001

<b>13. BATCH NUMBER</b>
-------------------------

Lot

<b>14. GENERAL CLASSIFICATION FOR SUPPLY</b>
--

<b>15. INSTRUCTIONS ON USE</b>
--------------------------------

<b>16. INFORMATION IN BRAILLE</b>
-----------------------------------

Justification for not including braille accepted.

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
---

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
--

PC  
SN  
NN



**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Aclasta 5 mg solution for infusion  
zoledronic acid

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

1 bottle contains 5 mg zoledronic acid (as monohydrate).

**3. LIST OF EXCIPIENTS**

Mannitol, sodium citrate and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for infusion

100 ml

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For single use only.  
Read the package leaflet before use.  
Intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
After opening: 24 hours at 2°C - 8°C.

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/05/308/001

Unit pack

EU/1/05/308/002

Multipack

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Justification for not including braille accepted.

**PARTICULARS TO APPEAR ON THE INTERMEDIATE PACKAGING****CARTON FOR INTERMEDIATE PACK (WITHOUT BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Aclasta 5 mg solution for infusion  
zoledronic acid

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each bottle of 100 ml contains 5 mg zoledronic acid (as monohydrate).

**3. LIST OF EXCIPIENTS**

Mannitol, sodium citrate and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for infusion

1 bottle of 100 ml  
Component of a multipack. Not to be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For single use only.  
Read the package leaflet before use.  
Intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP  
After opening: 24 hours at 2°C - 8°C.

**9. SPECIAL STORAGE CONDITIONS**

<b>10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE</b>
--

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

<b>11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER</b>
---

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

<b>12. MARKETING AUTHORISATION NUMBER(S)</b>
--

EU/1/05/308/002

<b>13. BATCH NUMBER</b>
-------------------------

Lot

<b>14. GENERAL CLASSIFICATION FOR SUPPLY</b>
--

<b>15. INSTRUCTIONS ON USE</b>
--------------------------------

<b>16. INFORMATION IN BRAILLE</b>
-----------------------------------

Justification for not including braille accepted.

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
---

<b>18. UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
--

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**PARTICULARS TO APPEAR ON THE OUTER CARTON OF MULTIPACK (INCLUDING BLUE BOX)**

**1. NAME OF THE MEDICINAL PRODUCT**

Aclasta 5 mg solution for infusion  
zoledronic acid

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each bottle of 100 ml contains 5 mg zoledronic acid (as monohydrate).

**3. LIST OF EXCIPIENTS**

Mannitol, sodium citrate and water for injections.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Solution for infusion

Multipack: 5 bottles, each bottle of 100 ml.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For single use only.  
Read the package leaflet before use.  
Intravenous use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP  
After opening: 24 hours at 2°C - 8°C.

**9. SPECIAL STORAGE CONDITIONS**

<b>10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE</b>
--

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

<b>11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER</b>
---

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

<b>12. MARKETING AUTHORISATION NUMBER(S)</b>
--

EU/1/05/308/002

<b>13. BATCH NUMBER</b>
-------------------------

Lot

<b>14. GENERAL CLASSIFICATION FOR SUPPLY</b>
--

<b>15. INSTRUCTIONS ON USE</b>
--------------------------------

<b>16. INFORMATION IN BRAILLE</b>
-----------------------------------

Justification for not including braille accepted.

<b>17. UNIQUE IDENTIFIER – 2D BARCODE</b>
---

2D barcode carrying the unique identifier included.

<b>18. UNIQUE IDENTIFIER - HUMAN READABLE DATA</b>
--

PC  
SN  
NN

## **B. PACKAGE LEAFLET**

## **Package leaflet: Information for the user**

### **Aclasta 5 mg solution for infusion** zoledronic acid

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Aclasta is and what it is used for
2. What you need to know before you are given Aclasta
3. How Aclasta is given
4. Possible side effects
5. How to store Aclasta
6. Contents of the pack and other information

#### **1. What Aclasta is and what it is used for**

Aclasta contains the active substance zoledronic acid. It belongs to a group of medicines called bisphosphonates and is used to treat post-menopausal women and adult men with osteoporosis or osteoporosis caused by treatment with corticosteroids used to treat inflammation, and Paget's disease of the bone in adults.

##### **Osteoporosis**

Osteoporosis is a disease that involves the thinning and weakening of the bones and is common in women after the menopause, but can also occur in men. At the menopause, a woman's ovaries stop producing the female hormone oestrogen, which helps keep bones healthy. Following the menopause bone loss occurs, bones become weaker and break more easily. Osteoporosis could also occur in men and women because of the long term use of steroids, which can affect the strength of bones. Many patients with osteoporosis have no symptoms but they are still at risk of breaking bones because osteoporosis has made their bones weaker. Decreased circulating levels of sex hormones, mainly oestrogens converted from androgens, also play a role in the more gradual bone loss observed in men. In both women and men, Aclasta strengthens the bone and therefore makes it less likely to break. Aclasta is also used in patients who have recently broken their hip in a minor trauma such as a fall and therefore are at risk of subsequent bone breaks.

##### **Paget's disease of the bone**

It is normal that old bone is removed and is replaced with new bone material. This process is called remodelling. In Paget's disease, bone remodelling is too rapid and new bone is formed in a disordered fashion, which makes it weaker than normal. If the disease is not treated, bones may become deformed and painful, and may break. Aclasta works by returning the bone remodelling process to normal, securing formation of normal bone, thus restoring strength to the bone.

#### **2. What you need to know before you are given Aclasta**

Follow all instructions given to you by your doctor, pharmacist or nurse carefully before you are given Aclasta.

##### **You must not be given Aclasta:**

- if you are allergic to zoledronic acid, other bisphosphonates or any of the other ingredients of



- this medicine (listed in section 6).
- if you have hypocalcaemia (this means that the levels of calcium in your blood are too low).
- if you have severe kidney problems.
- if you are pregnant.
- if you are breast-feeding.

### **Warnings and precautions**

Talk to your doctor before you are given Aclasta:

- if you are being treated with any medicine containing zoledronic acid, which is also the active substance of Aclasta (zoledronic acid is used in adult patients with certain types of cancer to prevent bone complications or to reduce the amount of calcium).
- if you have a kidney problem, or used to have one.
- if you are unable to take daily calcium supplements.
- if you have had some or all of the parathyroid glands in your neck surgically removed.
- if you have had sections of your intestine removed.

A side effect called osteonecrosis of the jaw (ONJ) (bone damage in the jaw) has been reported in the post-marketing setting in patients receiving Aclasta (zoledronic acid) for osteoporosis. ONJ can also occur after stopping treatment.

It is important to try and prevent ONJ developing as it is a painful condition that can be difficult to treat. In order to reduce the risk of developing osteonecrosis of the jaw, there are some precautions you should take.

Before receiving Aclasta treatment, tell your doctor, pharmacist or nurse if

- you have any problems with your mouth or teeth such as poor dental health, gum disease, or a planned tooth extraction;
- you do not receive routine dental care or have not had a dental check-up for a long time;
- you are a smoker (as this may increase the risk of dental problems);
- you have previously been treated with a bisphosphonate (used to treat or prevent bone disorders);
- you are taking medicines called corticosteroids (such as prednisolone or dexamethasone)
- you have cancer.

Your doctor may ask you to undergo a dental examination before you start treatment with Aclasta.

While being treated with Aclasta, you should maintain good oral hygiene (including regular teeth brushing) and receive routine dental check-ups. If you wear dentures you should make sure these fit properly. If you are under dental treatment or are due to undergo dental surgery (e.g. tooth extractions), inform your doctor about your dental treatment and tell your dentist that you are being treated with Aclasta. Contact your doctor and dentist immediately if you experience any problems with your mouth or teeth such as loose teeth, pain or swelling, or non-healing of sores or discharge, as these could be signs of osteonecrosis of the jaw.

### **Monitoring test**

Your doctor should do a blood test to check your kidney function (levels of creatinine) before each dose of Aclasta. It is important for you to drink at least 2 glasses of fluid (such as water), within a few hours before receiving Aclasta, as directed by your healthcare provider.

### **Children and adolescents**

Aclasta is not recommended for anyone under 18 years of age.

### **Other medicines and Aclasta**

Tell your doctor, pharmacist or nurse if you are taking, have recently taken or might take any other medicines.

It is important for your doctor to know all the medicines you are taking, especially if you are taking

any medicines known to be harmful to your kidneys (e.g. aminoglycosides) or diuretics (“waterpills”) that may cause dehydration.

### **Pregnancy and breast-feeding**

You must not be given Aclasta if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby.

Ask your doctor, pharmacist or nurse for advice before taking this medicine.

### **Driving and using machines**

If you feel dizzy while taking Aclasta, do not drive or use machines until you feel better.

### **Aclasta contains sodium**

This medicinal product contains less than 1 mmol sodium (23 mg) per 100 ml vial of Aclasta, i.e., essentially “sodium free”.

## **3. How Aclasta is given**

Follow carefully all instructions given to you by your doctor or nurse. Check with your doctor or nurse if you are not sure.

### **Osteoporosis**

The usual dose is 5 mg given as one infusion per year into a vein by your doctor or nurse. The infusion will take at least 15 minutes.

In case you recently broke your hip, it is recommended that Aclasta is administered two or more weeks after your hip repair surgery.

It is important to take calcium and vitamin D supplements (for example tablets) as directed by your doctor.

For osteoporosis, Aclasta works for one year. Your doctor will let you know when to return for your next dose.

### **Paget’s disease**

For the treatment of Paget’s disease, Aclasta should be prescribed only by physicians with experience in the treatment of Paget’s disease of the bone.

The usual dose is 5 mg, given to you as one initial infusion into a vein by your doctor or nurse. The infusion will take at least 15 minutes. Aclasta may work for longer than one year, and your doctor will let you know if you need to be treated again.

Your doctor may advise you to take calcium and vitamin D supplements (e.g. tablets) for at least the first ten days after being given Aclasta. It is important that you follow this advice carefully so that the level of calcium in your blood does not become too low in the period after the infusion. Your doctor will inform you regarding the symptoms associated with hypocalcaemia.

### **Aclasta with food and drink**

Make sure you drink enough fluids (at least one or two glasses) before and after the treatment with Aclasta, as directed by your doctor. This will help to prevent dehydration. You may eat normally on the day you are treated with Aclasta. This is especially important in patients who take diuretics (“water pills”) and in elderly patients (age 65 years or over).

### **If you missed a dose of Aclasta**

Contact your doctor or hospital as soon as possible to re-schedule your appointment.

### **Before stopping Aclasta therapy**

If you are considering stopping Aclasta treatment, please go to your next appointment and discuss this with your doctor. Your doctor will advise you and decide how long you should be treated with Aclasta.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Side effects related to the first infusion are very common (occurring in more than 30% of patients) but are less common following subsequent infusions. The majority of the side effects, such as fever and chills, pain in the muscles or joints, and headache, occur within the first three days following the dose of Aclasta. The symptoms are usually mild to moderate and go away within three days. Your doctor can recommend a mild pain reliever such as ibuprofen or paracetamol to reduce these side effects. The chance of experiencing these side effects decreases with subsequent doses of Aclasta.

### **Some side effects could be serious**

#### Common (may affect up to 1 in 10 people)

Irregular heart rhythm (atrial fibrillation) has been seen in patients receiving Aclasta for the treatment of postmenopausal osteoporosis. It is currently unclear whether Aclasta causes this irregular heart rhythm but you should report it to your doctor if you experience such symptoms after you have received Aclasta.

#### Uncommon (may affect up to 1 in 100 people)

Swelling, redness, pain and itching to the eyes or eye sensitivity to light.

#### Very rare (may affect up to 1 in 10,000 people)

Talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear.

#### Not known (frequency cannot be estimated from the available data)

Pain in the mouth and/or jaw, swelling or non-healing sores in the mouth or jaw, discharge, numbness or a feeling of heaviness in the jaw, or loosening of a tooth; these could be signs of bone damage in the jaw (osteonecrosis). Tell your doctor and dentist immediately if you experience such symptoms while being treated with Aclasta or after stopping treatment.

Kidney disorders (e.g. decreased urine output) may occur. Your doctor should do a blood test to check your kidney function before each dose of Aclasta. It is important for you to drink at least 2 glasses of fluid (such as water), within a few hours before receiving Aclasta, as directed by your healthcare provider.

If you experience any of the above side effects, you should contact your doctor immediately.

### **Aclasta may also cause other side effects**

#### Very common (may affect more than 1 in 10 people)

Fever

#### Common (may affect up to 1 in 10 people)

Headache, dizziness, sickness, vomiting, diarrhoea, pain in the muscles, pain in the bones and/or joints, pain in the back, arms or legs, flu-like symptoms (e.g. tiredness, chills, joint and muscle pain), chills, feeling of tiredness and lack of interest, weakness, pain, feeling unwell, swelling and/or pain at the infusion site.

In patients with Paget's disease, symptoms due to low blood calcium, such as muscle spasms, or

numbness, or a tingling sensation especially in the area around the mouth have been reported.

Uncommon (may affect up to 1 in 100 people)

Flu, upper respiratory tract infections, decreased red cell count, loss of appetite, sleeplessness, sleepiness which may include reduced alertness and awareness, tingling sensation or numbness, extreme tiredness, trembling, temporary loss of consciousness, eye infection or irritation or inflammation with pain and redness, spinning sensation, increased blood pressure, flushing, cough, shortness of breath, upset stomach, abdominal pain, constipation, dry mouth, heartburn, skin rash, excessive sweating, itching, skin reddening, neck pain, stiffness in muscles, bones and/or joints, joint swelling, muscle spasms, shoulder pain, pain in your chest muscles and rib cage, joint inflammation, muscular weakness, abnormal kidney test results, abnormal frequent urination, swelling of hands, ankles or feet, thirst, toothache, taste disturbances.

Rare (may affect up to 1 in 1,000 people)

Unusual fracture of the thigh bone particularly in patients on long-term treatment for osteoporosis may occur rarely. Contact your doctor if you experience pain, weakness or discomfort in your thigh, hip or groin as this may be an early indication of a possible fracture of the thigh bone. Low levels of phosphate in the blood.

Not known (frequency cannot be estimated from the available data)

Severe allergic reactions including dizziness and difficulty breathing, swelling mainly of the face and throat, decreased blood pressure, dehydration secondary to acute phase reactions (post-dose symptoms such as fever, vomiting and diarrhoea).

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Aclasta**

Your doctor, pharmacist or nurse knows how to store Aclasta properly.

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and bottle after EXP.
- The unopened bottle does not require any special storage conditions.
- After opening the bottle, the product should be used immediately in order to avoid microbial contamination. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C – 8°C. Allow the refrigerated solution to reach room temperature before administration.

**6. Contents of the pack and other information**

**What Aclasta contains**

- The active substance is zoledronic acid. Each bottle with 100 ml of solution contains 5 mg zoledronic acid (as monohydrate).  
One ml solution contains 0.05 mg zoledronic acid (as monohydrate).
- The other ingredients are mannitol, sodium citrate and water for injections.

**What Aclasta looks like and contents of the pack**

Aclasta is a clear and colourless solution. It comes in 100 ml plastic bottles as a ready-to-use solution for infusion. It is supplied in packs containing one bottle as unit pack, or in multipacks comprising five packs, each containing one bottle. Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

**Manufacturer**

Novartis Pharma GmbH  
Roonstraße 25  
D-90429 Nuremberg  
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**

Sandoz N.V.  
Telecom Gardens  
Medialaan 40  
B-1800 Vilvoorde  
Tél/Tel: +32 (0)2 722 97 97

**Lietuva**

Sandoz Pharmaceuticals d.d  
Branch Office Lithuania  
Seimyniskiu 3A  
LT – 09312 Vilnius  
Tel: +370 5 2636 037

**България**

КЧТ Сандоз България  
Тел.: +359 2 970 47 47

**Luxembourg/Luxemburg**

Sandoz N.V.  
Telecom Gardens  
Medialaan 40  
B-1800 Vilvoorde  
Tél/Tel: +32 (0)2 722 97 97

**Česká republika**

Sandoz s.r.o.  
Na Pankráci 1724/129  
CZ-140 00, Praha 4  
Tel: +420 225 775 111  
[office.cz@sandoz.com](mailto:office.cz@sandoz.com)

**Magyarország**

Sandoz Hungária Kft.  
Bartók Béla út 43-47  
H-1114 Budapest  
Tel: +36 1 430 2890  
[Info.hungary@sandoz.com](mailto:Info.hungary@sandoz.com)

**Danmark**

Sandoz A/S  
Edvard Thomsens Vej 14  
DK-2300 København S  
Tlf: +45 6395 1000  
[info.danmark@sandoz.com](mailto:info.danmark@sandoz.com)

**Malta**

Sandoz Pharmaceuticals d.d.  
Verovškova 57  
SI-1000 Ljubljana  
Slovenia  
Tel: +356 21222872

**Deutschland**

Hexal AG  
Industriestr. 25  
D-83607 Holzkirchen  
Tel: +49 8024 908-0  
[service@hexal.com](mailto:service@hexal.com)

**Nederland**

Sandoz B.V.  
Veluwezoom 22  
NL-1327 AH Almere  
Tel: +31 (0)36 5241600  
[info.sandoz-nl@sandoz.com](mailto:info.sandoz-nl@sandoz.com)

**Eesti**

Sandoz d.d. Eesti filiaal  
Pärnu mnt 105  
EE – 11312 Tallinn  
Tel: +372 6652405

**Ελλάδα**

SANDOZ HELLAS  
ΜΟΝΟΠΡΟΣΩΠΗ Α.Ε.  
Τηλ: +30 216 600 5000

**España**

Bexal Farmacéutica, S.A.  
Centro Empresarial Parque Norte  
Edificio Roble  
C/ Serrano Galvache, 56  
28033 Madrid  
Tel: +34 900 456 856

**France**

Sandoz SAS  
49, avenue Georges Pompidou  
F-92300 Levallois-Perret  
Tél: +33 1 49 64 48 00

**Hrvatska**

Sandoz d.o.o.  
Maksimirska 120  
10 000 Zagreb  
Tel : +385 1 235 3111  
[upit.croatia@sandoz.com](mailto:upit.croatia@sandoz.com)

**Ireland**

Sandoz Pharmaceuticals d.d.  
Verovškova ulica 57  
1000 Ljubljana  
Slovenia

**Ísland**

Sandoz A/S  
Edvard Thomsens Vej 14  
DK-2300 Kaupmaannahöfn S  
Danmörk  
Tlf: +45 6395 1000  
[info.danmark@sandoz.com](mailto:info.danmark@sandoz.com)

**Norge**

Sandoz A/S  
Edvard Thomsens Vej 14  
DK-2300 København S  
Danmark  
Tlf: +45 6395 1000  
[info.norge@sandoz.com](mailto:info.norge@sandoz.com)

**Österreich**

Sandoz GmbH  
Biochemiestr. 10  
A-6250 Kundl  
Tel: +43(0)1 86659-0

**Polska**

Sandoz Polska Sp. z o.o.  
ul. Domaniewska 50 C  
02 672 Warszawa  
Tel.: +48 22 209 7000  
[maintenance.pl@sandoz.com](mailto:maintenance.pl@sandoz.com)

**Portugal**

Sandoz Farmacêutica Lda.  
Avenida Professor Doutor Cavaco Silva, n.º10E  
Taguspark  
P-2740–255 Porto Salvo  
Tel: +351 21 196 40 42  
[regaff.portugal@sandoz.com](mailto:regaff.portugal@sandoz.com)

**România**

Sandoz S.R.L.  
Strada Livezeni Nr. 7a  
540472 Târgu Mureş  
Tel: +40 21 407 51 60

**Slovenija**

Lek farmacevtska družba d.d.  
Verovškova 57  
SI-1526 Ljubljana  
Tel: +386 1 580 21 11  
[Info.lek@sandoz.com](mailto:Info.lek@sandoz.com)

**Slovenská republika**

Sandoz d.d. - organizačná zložka  
Žižkova 22B  
811 02 Bratislava  
Tel: +421 2 48 200 600  
[sk.regulatory@sandoz.com](mailto:sk.regulatory@sandoz.com)

**Italia**

Sandoz S.p.A.  
Largo Umberto Boccioni, 1  
I-21040 Origgio / VA  
Tel: +39 02 96 54 1  
regaff.italy@sandoz.com

**Κύπρος**

Sandoz Pharmaceuticals d.d.  
Verovskova 57  
SI-1000 Ljubljana  
Σλοβενία  
Τηλ: +357 22 69 0690

**Latvija**

Sandoz d.d. Latvia filiāle  
K.Valdemāra 33 – 29  
LV-1010 Rīga  
Tel: +371 67892006

**Suomi/Finland**

Sandoz A/S  
Edvard Thomsens Vej 14  
DK-2300 Kööpenhamina S  
Tanska  
Puh: + 358 010 6133 400  
info.suomi@sandoz.com

**Sverige**

Sandoz A/S  
Edvard Thomsens Vej 14  
DK-2300 Köpenhamn S  
Danmark  
Tel: +45 6395 1000  
info.sverige@sandoz.com

**United Kingdom (Northern Ireland)**

Sandoz Pharmaceuticals d.d.  
Verovskova 57  
SI-1000 Ljubljana  
Slovenia  
Tel: +43 5338 2000

**This leaflet was last revised in**

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>

## **INFORMATION FOR THE HEALTHCARE PROFESSIONAL**

The following information is intended for healthcare professionals only (see section 3):

### **How to prepare and administer Aclasta**

- Aclasta 5 mg solution for infusion is ready for use.

For single use only. Any unused solution should be discarded. Only clear solution free from particles and discoloration should be used. Aclasta must not be mixed or given intravenously with any other medicinal product and must be given through a separate vented infusion line at a constant infusion rate. The infusion time must not be less than 15 minutes. Aclasta must not be allowed to come into contact with any calcium-containing solutions. If refrigerated, allow the refrigerated solution to reach room temperature before administration. Aseptic techniques must be followed during preparation of the infusion. The infusion must be conducted according to standard medical practice.

### **How to store Aclasta**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and bottle after EXP.
- The unopened bottle does not require any special storage conditions.
- After opening the bottle, the product should be used immediately in order to avoid microbial contamination. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C - 8°C. Allow the refrigerated solution to reach room temperature before administration.