

DOI:10.61818/78570407

ISSN: 2965-7857



Climate monitoring bulletin for the major river basins: The Amazon Basin

Volume 4, Number 07

Manaus, February 18, 2026



MINISTÉRIO DA
CIÊNCIA, TECNOLOGIA
E INOVAÇÃO



Climate monitoring bulletin for the major river basins: The Amazon Basin

Editor in Chief Renato Cruz Senna

Meteorologist

Researcher - CODAM, INPA

Publishing Renato Cruz Senna

Tainá Sampaio Xavier Conchy Rocha

Adriano Nobre Arcos

Periodicity Weekly

Proofreading and Layout Inácio de Oliveira Lima Neto

Contact us Instituto Nacional de Pesquisas da Amazônia - INPA

Av. André Araújo, 2.936 - Petrópolis

CEP 69067-375 - Manaus -AM, Brasil

E-mail: renato.senna@inpa.gov.br

clima.amazonia@inpa.gov.br

Phone: +55 92 3643 3154 / 3643-3170

www.instagram.com/clima.amazonia



This research was supported as part of the Large-Scale Biosphere-Atmosphere Program in the Amazon (LBA), coordinated by the National Amazon Research Institute (INPA), funded by the Ministry of Science, Technology and Innovation (MCTI), a direct administration entity of the Brazilian Federal Government.




This work is licensed under a Creative Commons Attribution-Non Commercial - No Derivatives 4.0 International License.

Contents

Current conditions	1
Branco River Basin	2
Negro River Basin	2
Marañon River Basin	2
Ucayali River Basin	3
Napo River Basin	3
The main course of the Amazon River (Peru)	3
Javari River Basin	4
Putumayo and Içá River Basins	4
Jutaí River Basin	4
Juruá River Basin	5
Caquetá and Japurá River Basins	5
Tefé River Basin	5
Coari River Basin	6
Purus River Basin	6
The main course of the Solimões River	6
Madre de Dios and Beni River Basins	7
Mamoré River Basin	7
Guaporé and Iténez Rivers Basins	7
Ji-Paraná River Basin	8
Aripuanã River Basin	8
Madeira River Basin	8
Basins on the left bank of the Amazon River (Amazonas State)	9
Abacaxis River Basin	9
Juruena River Basin	9
Teles Pires River Basin	10
Tapajós River Basin	10
Basins on the left bank of the Amazon River (northwest of the Pará State)	10
Curuá Una River Basin	11
Basins on the left bank of the Amazon River (north-eastern of the Pará State)	11
Iriri River Basin	11
Xingu River Basin	12
The main course of the Amazon River (Brazil)	12
Multi-model sub-seasonal forecast	13
Reference Values	15
Categorizing rainfall anomalies	16
Behavior of the 07 and 30 days anomalies observed in previous weeks	17
Single-line diagram of the basins represented	20

Current conditions


Maps of observed precipitation conditions, individual graphics by basin are produced from MERGE/GPM data generated by INPE/CPTEC, considering the period from de 2000 a 2025. **Between January 20 to February 18, 2026, below-average rainfall caused precipitation deficits along the main course of the Amazon River in Brazilian territory, the watersheds of the Abacaxis, Branco, Coari, Curuá Una, Guaporé, Iriti, Jutai, Mamoré, the left bank basins of the Amazon River in the northeast and northwest of the state of Pará, Purus, Tapajós, Xingu, and the main course of the Solimões River; rainfall above climatological norms was recorded on the main course of the Amazon River in Peruvian territory and in the basins of the Içá, Javari, Marañon, Napo, and Ucayali rivers; rainfall close to normal recorded over the Aripuanã, Beni, Japurá, Ji-Paraná, Juruá, Juruena, and Madeira river basins, basins on the left bank of the Amazon River in the northeast of the state of Amazonas, and the Negro, Tefé, and Teles Pires rivers. The multimodel indicates below-normal rainfall for the coming weeks on the main course of the Amazon River in Brazilian territory, the Abacaxis, Aripuanã, Beni, Branco, Coari, Curuá Una, Guaporé, Iriti, Ji-Paraná, Juruena, Madeira, Mamoré, basins on the left bank of the Amazon River in the northeast and northwest of the state of Pará, Purus, Tapajós, Ucayali, and Xingu; rainfall above climatology is forecast for the Japurá and Negro river basins in the first week.**



1 Abacaxis	9 Guaporé	17 Jutai	25 Negro
2 Amazonas (BR)	10 Içá	18 Madeira	26 Purus
3 Amazonas (PE)	11 Iriti	19 Mamoré	27 Solimões
4 Aripuanã	12 Japurá	20 Marañon	28 Tapajós
5 Beni	13 Javari	21 Marg Esq (AM)	29 Tefé
6 Branco	14 Ji-Paraná	22 Marg Esq (PA) NE	30 Teles Pires
7 Coari	15 Juruá	23 Marg Esq (PA) NW	31 Ucayali
8 Curuá Una	16 Juruena	24 Napo	32 Xingu


Individual analysis by river basin

Branco River Basin




The climatology of the period under analysis indicates rainfall with records varying between **36 and 48 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 30 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.9**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Negro River Basin




The climatology of the period under analysis indicates rainfall with records varying between **171 e 198 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 186 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.0**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Marañon River Basin




The climatology of the period under analysis indicates rainfall with records varying between **158 e 180 mm** s being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 257 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **1.9**, classifies the basin as **tendency to very rainy condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **very rainy behavior or a tending to be very rainy**.

Ucayali River Basin




The climatology of the period under analysis indicates rainfall with records varying between **181 e 203 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 271 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **1.7**, classifies the basin as **tendency to very rainy condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

Napo River Basin




The climatology of the period under analysis indicates rainfall with records varying between **185 e 212 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 241 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.8**, classifies the basin as **tendency to rainy condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

The main course of the Amazon River (Peru)




The climatology of the period under analysis indicates rainfall with records varying between **282 e 310 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 396 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **1.9**, classifies the basin as **tendency to very rainy condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very rainy behavior or a tending to be very rainy**.

Javari River Basin




The climatology of the period under analysis indicates rainfall with records varying between **258 e 284 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 338 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **1.1**, classifies the basin as **rainy condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

Putumayo and Içá Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **228 e 254 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 278 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.8**, classifies the basin as **tendency to rainy condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

Jutaí River Basin




The climatology of the period under analysis indicates rainfall with records varying between **305 e 337 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 245 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.5**, classifies the basin as **tendency to very dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Juruá River Basin




The climatology of the period under analysis indicates rainfall with records varying between **254 e 282 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 246 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.4**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality**.

Caquetá and Japurá Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **180 e 202 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 198 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.1**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be rainy**.

Tefé River Basin




The climatology of the period under analysis indicates rainfall with records varying between **252 e 286 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 290 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.4**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality**.

Coari River Basin




The climatology of the period under analysis indicates rainfall with records varying between **253 e 278 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 236 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.7**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Purus River Basin




The climatology of the period under analysis indicates rainfall with records varying between **274 e 302 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 244 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.0**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

The main course of the Solimões River




The climatology of the period under analysis indicates rainfall with records varying between **251 e 283 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 242 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.6**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Madre de Dios and Beni Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **250 e 277 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 267 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.1**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Mamoré River Basin




The climatology of the period under analysis indicates rainfall with records varying between **219 e 248 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 146 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.0**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be very dry**.

Guaporé and Itenéz Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **193 e 217 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 144 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.5**, classifies the basin as **tendency to very dry contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Ji-Paraná River Basin




The climatology of the period under analysis indicates rainfall with records varying between **244 e 273 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 245 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.2**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Aripuanã River Basin




The climatology of the period under analysis indicates rainfall with records varying between **266 e 296 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 301 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.3**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality**.

Madeira River Basin




The climatology of the period under analysis indicates rainfall with records varying between **251 e 280 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 253 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.2**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality**.

Basins on the left bank of the Amazon River (Amazonas State)




The climatology of the period under analysis indicates rainfall with records varying between **200 e 224 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 209 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.1**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Abacaxis River Basin




The climatology of the period under analysis indicates rainfall with records varying between **253 e 277 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 232 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.8**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Juruena River Basin




The climatology of the period under analysis indicates rainfall with records varying between **262 e 289 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 270 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.1**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Teles Pires River Basin




The climatology of the period under analysis indicates rainfall with records varying between **273 e 300 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 267 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.4**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Tapajós River Basin




The climatology of the period under analysis indicates rainfall with records varying between **268 e 294 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 233 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.1**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Basins on the left bank of the Amazon River (northwest of the Pará State)




The climatology of the period under analysis indicates rainfall with records varying between **169 e 191 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 132 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.5**, classifies the basin as **tendency to very dry contition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Curuá Una River Basin




The climatology of the period under analysis indicates rainfall with records varying between **206 e 227 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 195 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.6**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Basins on the left bank of the Amazon River (north-estern of the Pará State)




The climatology of the period under analysis indicates rainfall with records varying between **196 e 218 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 167 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.2**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Iri River Basin




The climatology of the period under analysis indicates rainfall with records varying between **269 e 295 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 229 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.1**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Xingu River Basin



The climatology of the period under analysis indicates rainfall with records varying between **244 e 272 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 199 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.3**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.


The main course of the Amazon River (Brazil)




The climatology of the period under analysis indicates rainfall with records varying between **278 e 304 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **February 18, 2026, 250 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.0**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

A multi-model sub-seasonal forecast CPTEC/INPE-FUNCEME produced on 02/17/2026 for the next 7 and 14 days.

The calibrated sub-seasonal multi-model forecast CPTEC/INPE-FUNCEME is generated through scientific cooperation between CPTEC/INPE and FUNCEME, and comes from a set of 4 global models (a Brazilian model, BAM-1.2/CPTEC, and three US models, CFSv2/NCEP, GEFSv12/NCEP and ESRL/NOAA, the latter three from the SubX project). Brazilian model, BAM-1.2/CPTEC, and three US models, CFSv2/NCEP, GEFSv12/NCEP, and ESRL/NOAA, the latter three from the SubX project). The forecast precipitation anomalies are determined in relation to the climatological period from 1999 to 2016. The outputs for the 7 and 14-day forecast intervals are shown below, detailing the forecast behavior of the basins of interest.



The figure above shows the forecast for the 7-day interval between 02/18/2026 e 02/24/2026, prediction of a predominance of precipitation deficit (orange) in the south and northeast of the monitored region, over the main course of the Amazon River in Brazilian territory, the Abacaxis, Aripuanã, Beni, Branco, Coari, Curuá Una, Guaporé, Iriti, Ji-Paraná, Juruena, Madeira, Mamoré, basins on the left bank of the Amazon River in the northeast and northwest of the state of Pará, Purus, Tapajós, Teles Pires, Ucayali, and Xingu. Forecast of positive precipitation anomalies (blue) over the Japurá and Negro river basins. Forecast of rainfall close to climatology (white) over the other monitored basins.



The figure above shows the prognosis for the 14-day interval between 02/18/2026 e 03/03/2026, forecast of a predominance of precipitation deficit (orange) in almost the entire monitored region, over the main course of the Amazon River in Brazilian territory, the basins of the Abacaxis, Aripuanã, Beni, Branco, Coari, Curuá Una, Guaporé, Iriti, Japurá, Ji-Paraná, Juruena, Madeira, Mamoré, Marañon, basins on the left bank of the Amazon River in the northeast of the state of Amazonas and in the northeast and northwest of the state of Pará, Napo, Negro, Purus, Tapajós, Tefé, Ucayali, Xingu, and the main course of the Solimões River. No positive precipitation anomalies (blue) are forecast for the monitored region. Rainfall forecast close to climatology (white) over the main course of the Amazon River in Peruvian territory, the Javari, Juruá, Jutaí, and Teles Pires river basins.

Reference Values for 30-day accumulated precipitation on the date of analysis.

Table 1 shows the average accumulated precipitation values (mm of rain) per basin, based on precipitation estimates using satellite images, a product called MERGE/GPM, made available by the National Institute for Space Research, for the period 2000 - 2022, taking into account the geographical limits of the Amazon basin's watersheds. To this end, the quantile technique was used, as it is a suitable and accurate tool for categorizing precipitation and anomalies of discrete variables. The following thresholds were adopted: 5%, 12.5%, 20%, 27.5%, 35%, 42.5%, 57.5%, 65%, 72.5%, 80%, 87.5%, and 95% in order to stratify the technique and allow for a more detailed categorization of the conditions in each monitored basin.

18/02/2026	Quantiles for categorizing precipitation anomalies											
	5.0%	12.5%	20.0%	27.5%	35.0%	42.5%	57.5%	65.0%	72.5%	80.0%	87.5%	95.0%
Abacaxis	117	159	192	216	235	253	277	295	314	335	365	419
Amazonas (BR)	167	200	221	240	259	278	304	323	344	366	395	455
Amazonas (PE)	151	190	217	240	260	282	310	331	356	384	420	478
Aripuanã	135	177	206	229	248	266	296	319	345	373	404	456
Beni	147	178	198	216	233	250	277	297	319	349	389	454
Branco	6	12	17	23	29	36	48	59	74	94	117	150
Coari	153	175	196	216	235	253	278	296	314	339	371	420
Curuá Una	120	145	163	179	192	206	227	244	264	289	319	360
Guaporé	101	131	148	163	178	193	217	234	253	277	308	360
Içá	112	142	170	192	212	228	254	274	295	321	360	418
Iriri	136	180	207	230	250	269	295	314	335	362	395	457
Japurá	84	113	135	151	165	180	202	217	235	256	285	342
Javari	151	184	207	226	242	258	284	304	326	353	386	440
Ji-Paraná	110	161	189	208	226	244	273	294	317	341	372	420
Juruá	136	166	191	212	233	254	282	301	323	349	383	435
Juruena	143	179	203	224	243	262	289	308	330	355	390	448
Jutaí	169	202	231	259	284	305	337	359	380	403	430	478
Madeira	126	164	189	212	232	251	280	300	320	342	370	414
Mamoré	120	148	167	185	201	219	248	269	295	326	368	444
Marañon	73	96	113	129	144	158	180	195	212	231	257	297
Marg Esq (AM)	63	97	132	161	183	200	224	239	257	279	304	344
Marg Esq (PA) NE	100	137	155	170	184	196	218	233	251	273	299	339
Marg Esq (PA) NW	77	105	124	141	156	169	191	208	227	248	278	342
Napo	74	107	127	146	166	185	212	233	255	284	325	388
Negro	68	95	115	134	153	171	198	218	243	270	303	354
Purus	155	189	213	235	255	274	302	321	342	365	394	438
Solimões	130	165	191	212	233	251	283	306	332	360	389	431
Tapajós	130	175	208	232	252	268	294	311	332	356	386	443
Tefé	152	183	199	215	232	252	286	312	337	362	392	443
Teles Pires	154	187	215	237	256	273	300	319	339	363	395	456
Ucayali	104	123	138	153	167	181	203	219	236	257	285	334
Xingu	131	163	186	207	225	244	272	293	315	343	377	433

Table 1. Accumulated precipitation quantiles (mm) in 30 days (20 de January a 18 de February),

Climatology of the period (2000 - 2025) MERGE/GPM – INPE/CPTEC.

Categorizing rainfall anomalies

Using the values in the table above, it is possible to categorize the rainfall observed in the current year in relation to the values observed in the previous records since the beginning of the available series, so the observed values below the 5% quantile characterize the basin in an extremely dry condition, between 5 and 12.5% in an extremely dry tendency condition, between 12.5 and 20% very dry condition, between 20 and 27.5% very dry tendency, between 27.5 and 35% dry condition, between 35 and 42.5 dry tendency condition, values between 42.5 and 57.5 define normal condition, values between 57.5 and 65% rainy tendency condition, between 65 and 72.5% rainy condition, between 72.5 and 80% very rainy tendency, between 80 and 87.5 very rainy condition, between 87.5 and 95% indicate extremely rainy tendency and finally values above 95% define the basin in extremely rainy condition, as per the legend below.

QUANTILE	5.0%	12.5%	20.0%	27.5%	35.0%	42.5%		57.5%	65.0%	72.5%	80.0%	87.5%	95.0%
INDEX	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0.0	0.5	1.0	1.5	2.0	2.5	3.0
CATEGORY	EXTREMELY DRY	EXTREMELY DRY TENDENCY	VERY DRY	VERY DRY TENDENCY	DRY	DRY TENDENCY	NORMAL	RAINY TENDENCY	RAINY	VERY RAINY TENDENCY	VERY RAINY	EXTREMELY RAINY TENDENCY	EXTREMELY RAINY

The tables below show (Table 2A) the average precipitation observed (mm) in each basin, taking as reference the precipitation estimates by satellite using the MERGE technique, available at <http://ftp.cptec.inpe.br/modelos/tempo/MERGE/GPM/DAILY/> accumulated in 30 days on the indicated dates, the average values of the categorized anomalies (Table 2B) were estimated based on the anomaly value of each pixel in the monitored watershed area, calculated according to the methodology described in the previous item, on the same dates of precipitation monitoring, the color scale of the anomalies follows the legend described.

	Average accumulated rainfall in the basin (mm)				
	21/01/2026	28/01/2026	04/02/2026	11/02/2026	18/02/2026
Abacaxis	189	217	205	223	232
Amazonas (BR)	103	130	144	209	250
Amazonas (PE)	214	260	320	311	396
Aripuanã	232	247	226	264	301
Beni	309	290	300	274	267
Branco	27	30	40	32	30
Coari	188	226	236	236	236
Curuá Una	113	108	98	119	195
Guaporé	143	120	129	123	144
Içá	217	223	264	256	278
Iriri	202	225	201	192	229
Japurá	166	169	198	174	198
Javari	184	248	288	274	338
Ji-Paraná	201	254	249	233	245
Juruá	252	256	255	239	246
Juruena	242	236	206	236	270
Jutai	258	222	241	243	245
Madeira	255	266	273	254	253
Mamoré	195	159	162	130	146
Marañon	172	198	238	216	257
Marg Esq (AM)	169	186	176	185	209
Marg Esq (PA) NE	80	88	93	132	167
Marg Esq (PA) NW	79	95	92	126	132
Napo	157	208	258	231	241
Negro	176	206	219	185	186
Purus	277	282	287	244	244
Solimões	240	233	242	235	242
Tapajós	179	204	189	205	233
Tefé	254	277	306	312	290
Teles Pires	191	213	222	241	267
Ucayali	187	215	260	249	271
Xingu	165	182	174	180	199









Table 2A. Accumulated rainfall over 30 days (mm)
data MERGE/GPM – INPE/CPTEC.

	Average categorized anomaly in the basin				
	21/01/2026	28/01/2026	04/02/2026	11/02/2026	18/02/2026
-1.0	-1.0	-0.5	-0.9	-0.8	-0.8
-2.3	-2.3	-2.1	-2.2	-1.3	-1.0
-1.7	-1.7	-0.7	0.7	0.5	1.9
-0.7	-0.7	-0.5	-1.0	-0.3	0.3
0.9	0.9	0.5	0.4	0.0	-0.1
-1.3	-1.3	-0.9	-0.1	-0.8	-0.9
-2.0	-2.0	-0.9	-1.0	-1.0	-0.7
-1.2	-1.2	-1.8	-2.3	-2.2	-0.6
-1.6	-1.6	-2.2	-1.6	-1.8	-1.5
-0.6	-0.6	-0.4	0.6	0.5	0.8
-0.9	-0.9	-0.7	-1.4	-1.6	-1.1
-0.8	-0.8	-0.3	0.3	-0.5	0.1
-1.9	-1.9	-0.6	0.3	0.1	1.1
-1.1	-1.1	0.1	-0.1	-0.4	-0.2
-0.2	-0.2	0.1	-0.1	-0.4	-0.4
-0.5	-0.5	-0.7	-1.3	-0.7	-0.1
-1.1	-1.1	-1.3	-1.3	-1.5	-1.5
0.1	0.1	0.4	0.4	0.0	-0.2
-0.9	-0.9	-1.7	-1.5	-2.1	-2.0
0.5	0.5	1.1	1.8	1.3	1.9
-0.3	-0.3	0.1	-0.2	-0.3	-0.1
-2.2	-2.2	-2.2	-2.4	-1.8	-1.2
-1.8	-1.8	-1.6	-1.8	-1.3	-1.5
-1.3	-1.3	0.2	1.1	0.6	0.8
-0.4	-0.4	0.6	0.8	-0.2	0.0
0.0	0.0	0.1	0.1	-0.9	-1.0
-0.7	-0.7	-0.5	-0.5	-0.8	-0.6
-1.4	-1.4	-1.1	-1.6	-1.5	-1.1
-0.9	-0.9	-0.2	0.4	0.3	0.4
-1.8	-1.8	-1.3	-1.1	-0.7	-0.4
0.0	0.0	0.8	1.7	1.4	1.7
-1.6	-1.6	-1.4	-1.7	-1.7	-1.3


Table 2B. Categorized precipitation
anomaly by quantiles

Behavior of the 07 and 30 days anomalies observed in previous weeks


The following graphs illustrate the behavior of the precipitation anomaly index in recent weeks, red lines show the behavior for periods of 30 days and blue lines the behavior for periods of 7 days, updated weekly.




Guaporé and Iténez rivers



Içá and Putumayo rivers




Iriri River



Caqueta and Japurá rivers




Javari River




Ji-Paraná River




Juruá River




Juruena River




Jutaí River




Madeira River




Mamoré River




Marañón River




Left bank of the Amazon River AM




Left bank of the Amazon River NE-PA




Left bank of the Amazon River NW-PA




Napo River




Negro River




Purus River







Main course of the Solimões River




Tapajós River





Single-line diagram of the basins represented



Renato Cruz Senna

Pesquisador - CODAM

Meteorologista, CREA-AM 2880-D

Registro Nacional 040459935-4

Fone de contato +55 92 3643 3170

