

DOI:10.61818/78570414

ISSN: 2965-7857



Climate monitoring bulletin for the major river basins: The Amazon Basin

Volume 4, Number 14

Manaus, April 8, 2026



MINISTÉRIO DA
CIÊNCIA, TECNOLOGIA
E INOVAÇÃO



Climate monitoring bulletin for the major river basins: The Amazon Basin

Editor in Chief Renato Cruz Senna

Meteorologist

Researcher - CODAM, INPA

Publishing Renato Cruz Senna

Tainá Sampaio Xavier Conchy Rocha

Inácio de Oliveira Lima Neto

Adriano Nobre Arcos

Periodicity Weekly

Proofreading and Layout Inácio de Oliveira Lima Neto

Contact us Instituto Nacional de Pesquisas da Amazônia - INPA

Av. André Araújo, 2.936 - Petrópolis

CEP 69067-375 - Manaus -AM, Brasil

E-mail: renato.senna@inpa.gov.br

clima.amazonia@inpa.gov.br

Phone: +55 92 3643 3154 / 3643-3170

www.instagram.com/clima.amazonia



This research was supported as part of the Large-Scale Biosphere-Atmosphere Program in the Amazon (LBA), coordinated by the National Amazon Research Institute (INPA), funded by the Ministry of Science, Technology and Innovation (MCTI), a direct administration entity of the Brazilian Federal Government.



This work is licensed under a Creative Commons Attribution-Non Commercial - No Derivatives 4.0 International License.



MINISTÉRIO DA
CIÊNCIA, TECNOLOGIA
E INOVAÇÃO




Contents

Current conditions	1
Branco River Basin	2
Negro River Basin	2
Marañon River Basin	2
Ucayali River Basin	3
Napo River Basin	3
The main course of the Amazon River (Peru)	3
Javari River Basin	4
Putumayo and Içá River Basins	4
Jutaí River Basin	4
Juruá River Basin	5
Caquetá and Japurá River Basins	5
Tefé River Basin	5
Coari River Basin	6
Purus River Basin	6
The main course of the Solimões River	6
Madre de Dios and Beni River Basins	7
Mamoré River Basin	7
Guaporé and Itenéz Rivers Basins	7
Ji-Paraná River Basin	8
Aripuanã River Basin	8
Madeira River Basin	8
Basins on the left bank of the Amazon River (Amazonas State)	9
Abacaxis River Basin	9
Juruena River Basin	9
Teles Pires River Basin	10
Tapajós River Basin	10
Basins on the left bank of the Amazon River (northwest of the Pará State)	10
Curuá Una River Basin	11
Basins on the left bank of the Amazon River (north-eastern of the Pará State)	11
Iriri River Basin	11
Xingu River Basin	12
The main course of the Amazon River (Brazil)	12
Multi-model sub-seasonal forecast	13
Reference Values	15
Categorizing rainfall anomalies	16
Behavior of the 07 and 30 days anomalies observed in previous weeks	17
Single-line diagram of the basins represented	20


Current conditions

Maps of observed precipitation conditions, individual graphics by basin are produced from MERGE/GPM data generated by INPE/CPTEC, considering the period from 2000 to 2025. **Between March 10 to April 8, 2026, below-average rainfall resulted in a precipitation deficit along the main course of the Amazon River in Brazilian and Peruvian territories, as well as in the watersheds of the Abacaxis, Aripuanã, Coari, Curuá Una, Içá, Iriri, Japurá, Javari, Ji-Paraná, Juruá, Juruena, Jutai, Madeira, the left-bank basins of the Amazon River in northeastern Amazonas State and northeastern Pará State, Napo, Negro, Purus, Tapajós, Tefé, Teles Pires, Xingu, and the main course of the Solimões River; above-normal rainfall recorded in the Branco and Ucayali river basins; near-normal rainfall recorded in the Beni, Guaporé, Mamoré, and Marañon river basins and the left-bank basins of the Amazon River in the northwest of the state of Pará.** The multimodel indicates, for the period from March 4, 2026, to March 10, 2026, **a forecast of rainfall below normal levels concentrated in the northeast of the monitored region, over the main course of the Amazon River in Brazilian territory, the basins of the Branco, Coari, Curuá Una, Iriri, Jutai, Mamoré, and Marañon rivers, the basins on the left bank of the Amazon River in the northeast of the state of Amazonas and in the northeast and northwest of the state of Pará, the Napo, Negro, Tapajós, and Tefé rivers, and the main course of the Solimões River; no rainfall above the climatological average is forecast for the monitored region.**




Individual analysis by river basin

Branco River Basin




The climatology of the period under analysis indicates rainfall with records varying between **71 and 94 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 119 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.7**, classifies the basin as **tendency to rainy condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

Negro River Basin




The climatology of the period under analysis indicates rainfall with records varying between **234 e 258 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 187 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.4**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Marañon River Basin




The climatology of the period under analysis indicates rainfall with records varying between **188 e 208 mm** s being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 205 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.4**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality**.

Ucayali River Basin




The climatology of the period under analysis indicates rainfall with records varying between **150 e 169 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 188 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **1.0**, classifies the basin as **rainy condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **rainy behavior or a tending to be rainy**.

Napo River Basin




The climatology of the period under analysis indicates rainfall with records varying between **279 e 307 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 250 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.0**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

The main course of the Amazon River (Peru)




The climatology of the period under analysis indicates rainfall with records varying between **339 e 371 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 223 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.3**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be extremely dry**.

Javari River Basin




The climatology of the period under analysis indicates rainfall with records varying between **283 e 308 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 200 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.0**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Putumayo and Içá Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **299 e 329 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 222 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.9**, classifies the basin as **tendency to very dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be very dry**.

Jutaí River Basin




The climatology of the period under analysis indicates rainfall with records varying between **273 e 299 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 185 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.3**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be very dry**.

Juruá River Basin




The climatology of the period under analysis indicates rainfall with records varying between **241 e 264 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 199 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.3**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Caquetá and Japurá Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **268 e 293 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 199 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.8**, classifies the basin as **tendency to very dry contition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Tefé River Basin




The climatology of the period under analysis indicates rainfall with records varying between **262 e 279 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 166 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.8**, classifies the basin as **tendency to extremely dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be extremely dry**.

Coari River Basin




The climatology of the period under analysis indicates rainfall with records varying between **273 e 292 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 146 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-3.0**, classifies the basin as **extremely dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **very dry behavior or a tending to be extremely dry**.

Purus River Basin




The climatology of the period under analysis indicates rainfall with records varying between **237 e 262 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 183 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.4**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

The main course of the Solimões River




The climatology of the period under analysis indicates rainfall with records varying between **270 e 295 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 186 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.0**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Madre de Dios and Beni Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **186 e 209 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 183 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.4**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Mamoré River Basin




The climatology of the period under analysis indicates rainfall with records varying between **155 e 179 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 172 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.2**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Guaporé and Itenéz Rivers Basins




The climatology of the period under analysis indicates rainfall with records varying between **143 e 164 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 152 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.1**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Ji-Paraná River Basin




The climatology of the period under analysis indicates rainfall with records varying between **207 e 231 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 185 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.8**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be rainy**.

Aripuanã River Basin




The climatology of the period under analysis indicates rainfall with records varying between **219 e 244 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 185 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.2**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Madeira River Basin




The climatology of the period under analysis indicates rainfall with records varying between **229 e 255 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 198 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.1**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Basins on the left bank of the Amazon River (Amazonas State)




The climatology of the period under analysis indicates rainfall with records varying between **250 e 280 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026**, **220 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.9**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Abacaxis River Basin




The climatology of the period under analysis indicates rainfall with records varying between **253 e 278 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026**, **235 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.7**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Juruena River Basin




The climatology of the period under analysis indicates rainfall with records varying between **204 e 229 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026**, **173 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.3**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Teles Pires River Basin




The climatology of the period under analysis indicates rainfall with records varying between **225 e 248 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 153 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-2.0**, classifies the basin as **very dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Tapajós River Basin




The climatology of the period under analysis indicates rainfall with records varying between **257 e 287 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 230 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.1**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.

Basins on the left bank of the Amazon River (northwest of the Pará State)




The climatology of the period under analysis indicates rainfall with records varying between **231 e 262 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 252 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **0.2**, classifies the basin as **normal contition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

Curuá Una River Basin




The climatology of the period under analysis indicates rainfall with records varying between **250 e 274 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 233 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-0.9**, classifies the basin as **tendency to dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Basins on the left bank of the Amazon River (north-estern of the Pará State)




The climatology of the period under analysis indicates rainfall with records varying between **248 e 274 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 202 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.5**, classifies the basin as **tendency to very dry contition**. In the coming weeks, climate behavior indicates a **increase** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Iriri River Basin




The climatology of the period under analysis indicates rainfall with records varying between **243 e 265 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 178 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.9**, classifies the basin as **tendency to very dry contition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be very dry**.

Xingu River Basin



The climatology of the period under analysis indicates rainfall with records varying between **223 e 245 mm** being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 167 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.8**, classifies the basin as **tendency to very dry condition**. In the coming weeks, climate behavior indicates a **decrease** in rainfall volumes, the sub-seasonal forecast model suggests **dry behavior or a tending to be dry**.


The main course of the Amazon River (Brazil)




The climatology of the period under analysis indicates rainfall with records varying between **317 e 342 mm** sendo being considered normal (reference quantiles 42.5% and 57.5%). On **April 8, 2026, 282 mm** of average accumulated precipitation was observed over the basin in 30 days, the calculation of the average of the categorized anomaly index in the basin area the value of **-1.0**, classifies the basin as **dry condition**. In the coming weeks, climate behavior indicates a **maintenance** in rainfall volumes, the sub-seasonal forecast model suggests **behavior close to normality or a tending to be dry**.

A multi-model sub-seasonal forecast CPTEC/INPE-FUNCEME produced on 04/07/2026 for the next 7 and 14 days.

The calibrated sub-seasonal multi-model forecast CPTEC/INPE-FUNCEME is generated through scientific cooperation between CPTEC/INPE and FUNCEME, and comes from a set of 4 global models (a Brazilian model, BAM-1.2/CPTEC, and three US models, CFSv2/NCEP, GEFSv12/NCEP and ESRL/NOAA, the latter three from the SubX project). Brazilian model, BAM-1.2/CPTEC, and three US models, CFSv2/NCEP, GEFSv12/NCEP, and ESRL/NOAA, the latter three from the SubX project). The forecast precipitation anomalies are determined in relation to the climatological period from 1999 to 2016. The following are the outputs for the forecast interval from 1 to 7 and from 8 to 14 days, detailing the predicted behavior over the basins of interest



The figure above shows the forecast for the **7-day interval between 04/08/2026 and 04/14/2026**, forecast of a predominance of precipitation deficits (orange) in the northeastern part of the monitored region, along the main course of the Amazon River in Brazilian territory, and in the basins of the Branco, Coari, Curuá Una, Iriri, Jutaí, Mamoré, and Marañon rivers; the basins on the left bank of the Amazon River in the northeast of the state of Amazonas and in the northeast and northwest of the state of Pará; the Napo, Negro, Tapajós, and Tefé rivers; and the main course of the Solimões River. There is no forecast of predominantly positive precipitation anomalies (blue) over the monitored region. Forecast of rainfall close to climatological norms (white) over the remaining monitored basins.



The figure above shows the prognosis for the **7-day interval between 04/15/2026 and 04/21/2026**, forecast of a predominance of positive precipitation anomalies (blue) over the main course of the Amazon River in Peruvian territory, the basins of the Beni, Branco, Curuá Una, Içá, Iriri, Japurá, Javari, Juruena, Jutaí, Mamoré, Marañon, the left-bank basins of the Amazon River in northeastern Amazonas State and northwestern Pará State, Napo, Negro, Purus, Teles Pires, and Xingu. Forecast of a predominance of precipitation deficit (orange) over the basins on the left bank of the Amazon River in the northeast of the state of Pará. Forecast of rainfall close to climatological norms (white) over the other monitored basins.

Reference Values for 30-day accumulated precipitation on the date of analysis.

Table 1 shows the average accumulated precipitation values (mm of rain) per basin, based on precipitation estimates using satellite images, a product called MERGE/GPM, made available by the National Institute for Space Research, for the period 2000 - 2025, taking into account the geographical limits of the Amazon basin's watersheds. To this end, the quantile technique was used, as it is a suitable and accurate tool for categorizing precipitation and anomalies of discrete variables. The following thresholds were adopted: 5%, 12.5%, 20%, 27.5%, 35%, 42.5%, 57.5%, 65%, 72.5%, 80%, 87.5%, and 95% in order to stratify the technique and allow for a more detailed categorization of the conditions in each monitored basin.

08/04/2026	Quantiles for categorizing precipitation anomalies											
	5.0%	12.5%	20.0%	27.5%	35.0%	42.5%	57.5%	65.0%	72.5%	80.0%	87.5%	95.0%
Abacaxis	127	157	187	212	233	253	278	295	314	340	372	435
Amazonas (BR)	205	237	264	283	300	317	342	360	380	404	440	496
Amazonas (PE)	191	234	270	296	317	339	371	394	422	453	496	561
Aripuanã	103	137	165	186	203	219	244	262	283	305	337	393
Beni	92	117	138	155	171	186	209	225	242	263	290	336
Branco	18	32	41	51	60	71	94	116	143	173	206	248
Coari	188	210	228	245	261	273	292	307	325	346	372	418
Curuá Una	133	168	202	220	235	250	274	289	306	327	352	399
Guaporé	71	90	104	117	130	143	164	180	197	219	246	294
Içá	183	212	237	258	279	299	329	350	373	398	429	475
Iriri	133	167	190	210	227	243	265	283	302	328	365	424
Japurá	166	195	217	235	252	268	293	310	331	355	388	442
Javari	178	209	232	251	267	283	308	327	348	373	406	459
Ji-Paraná	91	122	148	171	190	207	231	250	269	289	314	354
Juruá	134	170	194	211	226	241	264	279	296	317	347	396
Juruena	109	138	157	174	189	204	229	246	265	288	318	365
Jutaí	160	191	217	237	255	273	299	319	338	360	388	434
Madeira	125	157	178	196	213	229	255	273	292	315	347	399
Mamoré	74	94	110	124	140	155	179	197	217	241	272	322
Marañon	107	129	146	161	175	188	208	224	242	263	290	338
Marg Esq (AM)	121	160	186	209	229	250	280	301	326	354	388	440
Marg Esq (PA) NE	127	165	192	214	232	248	274	294	316	341	371	419
Marg Esq (PA) NW	111	146	169	189	211	231	262	283	305	334	374	453
Napo	174	199	219	238	259	279	307	328	351	378	413	470
Negro	131	162	183	202	218	234	258	276	297	322	355	405
Purus	135	165	187	205	221	237	262	280	300	325	359	414
Solimões	171	195	216	236	253	270	295	313	334	357	388	437
Tapajós	136	174	198	220	239	257	287	308	331	357	390	444
Tefé	178	201	220	237	250	262	279	291	306	325	348	409
Teles Pires	129	157	176	194	209	225	248	265	284	308	344	403
Ucayali	83	101	115	126	138	150	169	183	199	219	248	297
Xingu	126	155	175	192	207	223	245	260	278	300	329	376

Table 1. Accumulated precipitation quantiles (mm) in 30 days (March 10 to April 8),

Climatology of the period (2000 - 2025) MERGE/GPM – INPE/CPTEC.

Categorizing rainfall anomalies

Using the values in the table above, it is possible to categorize the rainfall observed in the current year in relation to the values observed in the previous records since the beginning of the available series, so the observed values below the 5% quantile characterize the basin in an extremely dry condition, between 5 and 12.5% in an extremely dry tendency condition, between 12.5 and 20% very dry condition, between 20 and 27.5% very dry tendency, between 27.5 and 35% dry condition, between 35 and 42.5% dry tendency condition, values between 42.5 and 57.5 define normal condition, values between 57.5 and 65% rainy tendency condition, between 65 and 72.5% rainy condition, between 72.5 and 80% very rainy tendency, between 80 and 87.5% very rainy condition, between 87.5 and 95% indicate extremely rainy tendency and finally values above 95% define the basin in extremely rainy condition, as per the legend below.

QUANTILE	5.0%	12.5%	20.0%	27.5%	35.0%	42.5%		57.5%	65.0%	72.5%	80.0%	87.5%	95.0%
INDEX	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0.0	0.5	1.0	1.5	2.0	2.5	3.0
CATEGORY	EXTREMELY DRY	EXTREMELY DRY TENDENCY	VERY DRY	VERY DRY TENDENCY	DRY	DRY TENDENCY	NORMAL	RAINY TENDENCY	RAINY	VERY RAINY TENDENCY	VERY RAINY	EXTREMELY RAINY TENDENCY	EXTREMELY RAINY

The tables below show (Table 2A) the average precipitation observed (mm) in each basin, taking as reference the precipitation estimates by satellite using the MERGE technique, available at <http://ftp.cptec.inpe.br/modelos/tempo/MERGE/GPM/DAILY/> accumulated in 30 days on the indicated dates, the average values of the categorized anomalies (Table 2B) were estimated based on the anomaly value of each pixel in the monitored watershed area, calculated according to the methodology described in the previous item, on the same dates of precipitation monitoring, the color scale of the anomalies follows the legend described.

	Average accumulated rainfall in the basin (mm)				
	11/03/2026	18/03/2026	25/03/2026	01/04/2026	08/04/2026
Abacaxis	241	226	237	257	235
Amazonas (BR)	282	294	283	278	282
Amazonas (PE)	250	213	187	205	223
Aripuanã	228	174	190	180	185
Beni	219	197	196	204	183
Branco	7	12	15	74	119
Coari	156	124	133	159	146
Curuá Una	254	276	251	244	233
Guaporé	200	188	180	168	152
Içá	184	189	175	199	222
Iriti	336	324	316	254	178
Japurá	123	125	114	167	199
Javari	194	167	166	180	200
Ji-Paraná	180	147	156	189	185
Juruá	173	154	153	181	199
Juruena	257	218	240	217	173
Jutai	188	149	126	169	185
Madeira	203	183	194	204	198
Mamoré	218	221	234	205	172
Marañon	180	174	190	194	205
Marg Esq (AM)	224	234	188	228	220
Marg Esq (PA) NE	171	190	160	204	202
Marg Esq (PA) NW	143	190	178	261	252
Napo	172	211	206	233	250
Negro	100	90	90	156	187
Purus	202	195	184	217	183
Solimões	137	128	125	168	186
Tapajós	339	318	311	282	230
Tefé	148	117	111	143	166
Teles Pires	240	216	205	184	153
Ucayali	174	160	172	175	188
Xingu	225	207	197	177	167









Table 2A. Accumulated rainfall over 30 days (mm)
data MERGE/GPM – INPE/CPTEC.

	Average categorized anomaly in the basin				
	11/03/2026	18/03/2026	25/03/2026	01/04/2026	08/04/2026
	-1.0	-1.1	-0.9	-0.4	-0.7
	-1.2	-0.8	-1.0	-1.4	-1.0
	-1.4	-1.9	-2.4	-2.4	-2.3
	-0.9	-1.7	-1.3	-1.5	-1.2
	-1.1	-1.0	-0.6	-0.1	-0.4
	-2.7	-2.2	-2.2	0.0	0.7
	-2.7	-3.0	-2.9	-2.9	-3.0
	0.2	0.6	-0.1	-0.6	-0.9
	-0.1	0.1	0.2	0.0	-0.1
	-2.0	-2.1	-2.5	-2.2	-1.9
	0.5	0.6	0.4	-0.7	-1.9
	-2.5	-2.6	-2.8	-2.3	-1.8
	-2.1	-2.5	-2.4	-2.4	-2.0
	-1.4	-2.0	-1.6	-1.1	-0.8
	-2.2	-2.3	-2.1	-1.8	-1.3
	-0.5	-0.9	0.0	-0.3	-1.3
	-2.4	-2.8	-2.9	-2.4	-2.3
	-1.4	-1.5	-1.3	-1.1	-1.1
	-0.5	0.2	0.8	0.5	0.2
	0.1	-0.2	0.0	0.1	0.4
	-0.6	-0.4	-1.5	-0.9	-0.9
	-1.9	-1.4	-1.9	-1.2	-1.5
	-1.9	-1.0	-1.0	0.4	0.2
	-1.8	-1.3	-1.6	-1.2	-1.0
	-2.6	-2.7	-2.7	-1.9	-1.4
	-1.9	-1.8	-1.9	-1.0	-1.4
	-2.7	-2.7	-2.8	-2.3	-2.0
	0.5	0.3	0.3	-0.3	-1.1
	-2.8	-2.9	-3.0	-2.9	-2.8
	-1.0	-1.1	-1.2	-1.5	-2.0
	-0.8	-1.0	-0.3	0.3	1.0
	-0.7	-1.2	-1.4	-1.8	-1.8


Table 2B. Categorized precipitation
anomaly by quantiles

Behavior of the 07 and 30 days anomalies observed in previous weeks


The following graphs illustrate the behavior of the precipitation anomaly index in recent weeks, red lines show the behavior for periods of 30 days and blue lines the behavior for periods of 7 days, updated weekly.




Guaporé and Itenéz rivers




Içá and Putumayo rivers




Irirí River




Caqueta and Japurá rivers




Javari River




Ji-Paraná River




Juruá River




Juruena River




Jutaí River




Madeira River




Mamoré River




Marañón River




Left bank of the Amazon River AM




Left bank of the Amazon River NE-PA




Left bank of the Amazon River NW-PA




Napo River




Negro River




Purus River




Main course of the Solimões River




Tapajós River





Single-line diagram of the basins represented



Renato Cruz Senna

Pesquisador - CODAM

Meteorologista, CREA-AM 2880-D

Registro Nacional 040459935-4

Fone de contato +55 92 3643 3170

