

Second Order ODEs

The **general form** of a second order ODE is

$$\frac{d^2y}{dt^2} = f(t, y, \frac{dy}{dt})$$

for some given function f .

1 Linear Homogeneous Equations

Definition 1.1 (Linear Equation).

A second order ODE is called **linear** if it can be written in the form

$$a(t) \cdot y'' + b(t) \cdot y' + c(t) \cdot y = g(t)$$

If $a(t) \neq 0$, we can divide both sides by $a(t)$ to get $y'' + p(t) \cdot y' + q(t) \cdot y = r(t)$.

Definition 1.2 (Homogeneous Equation).

A linear equation is called **homogeneous** if it can be written in the form

$$a(t) \cdot y'' + b(t) \cdot y' + c(t) \cdot y = 0$$

Otherwise, it is called **non-homogeneous**.

Theorem 1.1 (Existence and Uniqueness Theorem).

Consider the IVP:

$$y'' + p(t) \cdot y' + q(t) \cdot y = r(t), \quad y(t_0) = y_0, \quad y'(t_0) = y_1$$

if there exists an open interval $I \subseteq \mathbb{R}$ such that $t_0 \in I$ and p , q and r are **continuous** on I , then there is a **exact one** solution $y = \phi(t)$, and it exists **throughout** I .

Theorem 1.2 (Principle of Superposition).

If y_1 and y_2 are two solutions of the **homogeneous equation**

$$a(t) \cdot y'' + b(t) \cdot y' + c(t) \cdot y = 0$$

then any linear combination $c_1 y_1(t) + c_2 y_2(t)$ is also a solution of the ODE.

Definition 1.3 (Wronskian).

The **Wronskian**, or **Wronskian determinant**, of two functions y_1 and y_2 is defined as

$$W(y_1, y_2)(t) := \begin{vmatrix} y_1(t) & y_2(t) \\ y_1'(t) & y_2'(t) \end{vmatrix} = y_1(t) \cdot y_2'(t) - y_2(t) \cdot y_1'(t)$$

Theorem 1.3.

Suppose $p(t)$ and $q(t)$ are continuous on an open interval I , and y_1 and y_2 are two linearly independent solutions of the homogeneous equation

$$y'' + p(t) \cdot y' + q(t) \cdot y = 0 \quad \text{for } t \in I \quad (1)$$

Let $\phi(t)$ be the **unique** solution (by Theorem 1.1) of the IVP:

$$y'' + p(t) \cdot y' + q(t) \cdot y = 0 \quad y(t_0) = m, \quad y'(t_0) = n \quad \text{for } t \in I \quad \text{and } t_0 \in I \quad (2)$$

Then, for $t \in I$,

$$\exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \phi(t) = c_1 y_1(t) + c_2 y_2(t) \quad \Longleftrightarrow \quad W(y_1, y_2)(t_0) \neq 0$$

Proof.

Sufficiency. Since $\phi(t_0) = m$ and $\phi'(t_0) = n$, we have

$$\begin{aligned} \exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \phi(t) = c_1 y_1(t) + c_2 y_2(t) &\implies \exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \begin{bmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} m \\ n \end{bmatrix} \\ &\implies \exists b_1, b_2 \in \mathbb{R}, \exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \begin{bmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} \\ &\Longleftrightarrow W(y_1, y_2)(t_0) \neq 0 \end{aligned}$$

Necessity.

$$W(y_1, y_2)(t_0) \neq 0 \implies \exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \begin{bmatrix} y_1(t_0) & y_2(t_0) \\ y_1'(t_0) & y_2'(t_0) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} = \begin{bmatrix} m \\ n \end{bmatrix}$$

Let $\varphi(t) := c_1 y_1(t) + c_2 y_2(t), t \in I$. Then we have $\varphi(t_0) = m$ and $\varphi'(t_0) = n$.

Theorem 1.2 (Principle of Superposition) $\implies \varphi(t)$ is a solution of the ODE eq. (1).

Theorem 1.1 (Existence and Uniqueness Theorem) $\implies \varphi(t)$ is the **only** solution of the IVP in eq. (2).

Thus, $\varphi(t) = \phi(t)$ and we have

$$W(y_1, y_2)(t_0) \neq 0 \implies \exists! c_1, c_2 \in \mathbb{R} \text{ s.t. } \phi(t) = \varphi(t) = c_1 y_1(t) + c_2 y_2(t)$$

□

Theorem 1.4.

Suppose $p(t)$ and $q(t)$ are continuous on an open interval I , and y_1 and y_2 are two linearly independent solutions of the homogeneous equation

$$y'' + p(t) \cdot y' + q(t) \cdot y = 0$$

for $t \in I$. Then,

The **general solution** is given by $\phi(t) = c_1 y_1(t) + c_2 y_2(t)$. $\Longleftrightarrow \exists t_0 \in I \text{ s.t. } W(y_1, y_2)(t_0) \neq 0$.

y_1, y_2 are said to form a **fundamental set of solutions (FSS)** of the ODE, if their Wronskian is non-zero.

Proof.

Let S denote the set of all solutions of the ODE. We aim to prove that

$$\forall y \in S, \exists c_1, c_2 \in \mathbb{R} \text{ s.t. } y(t) = c_1 y_1(t) + c_2 y_2(t) \quad \Longleftrightarrow \quad \exists t_0 \in I \text{ s.t. } W(y_1, y_2)(t_0) \neq 0$$

Let $\mathcal{S}_{\text{IVP}}[t^*, m, n]$ denote the solution of the following IVP (by Theorem 1.1, there is exactly one solution):

$$y'' + p(t) \cdot y' + q(t) \cdot y = 0 \quad y(t^*) = m, \quad y'(t^*) = n \quad \text{for } t \in I \quad \text{and } t^* \in I$$

By Theorem 1.3, we have

$$\forall t_0 \in I \left[\left(W(y_1, y_2)(t_0) \neq 0 \right) \leftrightarrow \forall m, n \left(\exists c_1, c_2 \in \mathbb{R} \text{ s.t. } \mathcal{S}_{\text{IVP}}[t_0, m, n] = c_1 y_1 + c_2 y_2 \right) \right]$$

Instantiate m, n with $y(t_0), y'(t_0)$ for all $y \in S$:

$$\forall t_0 \in I \left[\left(W(y_1, y_2)(t_0) \neq 0 \right) \leftrightarrow \forall y \in S \left(\exists c_1, c_2 \in \mathbb{R} \text{ s.t. } \mathcal{S}_{\text{IVP}}[t_0, y(t_0), y'(t_0)] = c_1 y_1 + c_2 y_2 \right) \right]$$

Since y satisfies the ODE and initial conditions itself, $\mathcal{S}_{\text{IVP}}[t_0, y(t_0), y'(t_0)]$ is identical to y . Thus we obtain

$$\forall t_0 \in I \left[\left(W(y_1, y_2)(t_0) \neq 0 \right) \leftrightarrow \forall y \in S \left(\exists c_1, c_2 \in \mathbb{R} \text{ s.t. } y = c_1 y_1 + c_2 y_2 \right) \right] \quad (3)$$

Now we can apply the following corollary from first-order formal logic:

$$\left(\exists x. R(x) \right) \wedge \forall x. \left(R(x) \rightarrow (P(x) \leftrightarrow Q) \right) \implies \left(\exists x. (R(x) \wedge P(x)) \right) \leftrightarrow Q$$

(Note that Q does not depend on x). Identifying $R(x)$ with $x \in I$, $P(x)$ with $W(y_1, y_2)(x) \neq 0$, and noting I is non-empty so that $\exists t_0 \in I$, we can transform eq. (3) into

$$\left(\exists t_0 \in I, W(y_1, y_2)(t_0) \neq 0 \right) \leftrightarrow \forall y \in S \left(\exists c_1, c_2 \in \mathbb{R} \text{ s.t. } y = c_1 y_1 + c_2 y_2 \right)$$

This completes the proof. □