

First Order ODEs

The **general form** of a first order ODE is

$$\frac{dy}{dt} = f(t, y)$$

for some given function f .

1 Linear

The **general form** of a first order linear ODE is

$$\frac{dy}{dt} + p(t) \cdot y = q(t)$$

for some given functions p and q .

1.1 Method of Integrating Factors

For ODEs of the form

$$\frac{dy}{dt} + p(t) \cdot y = q(t)$$

the **general solution** is

$$y = \frac{1}{\mu(t)} \int \mu(t) \cdot q(t) dt$$

where the **integrating factor** is any $\mu(t)$ that satisfies

$$\ln |\mu(t)| = \int p(t) dt$$

Remark. Indefinite integrals yield a family of functions differing by a constant. Each indefinite integral results in an independent constant, which represents a degree of freedom. However, the constant from $\mu(t)$ cancels out in the final solution for y .

1.2 Variation of Parameters

For ODEs of the form

$$\frac{dy}{dt} + p(t) \cdot y = q(t)$$

let $h(t)$ be a function such that

$$h(t) = \int p(t) dt$$

Then the **general solution** is given by

$$y = e^{-h(t)} \int q(t) \cdot e^{h(t)} dt$$

1.3

For ODEs of the form

$$\frac{dy}{dt} + p(t) \cdot y = q(t)$$

the **general solution** is given by

$$y = v \int \frac{q(t)}{v} dt$$

where v satisfies

$$\frac{dv}{dt} + p(t) \cdot v = 0$$

Proof.

We want to solve eq. (1). Suppose v satisfies eq. (2), which is separable and can be solved easily.

$$y' + p(t) \cdot y = q(t) \quad (1)$$

$$v' + p(t) \cdot v = 0 \quad (2)$$

Take the derivative of y/v :

$$\begin{aligned} \frac{d}{dt} \left(\frac{y}{v} \right) &= \frac{y'v - v'y}{v^2} \\ &= \frac{y'v + (p(t) \cdot v)y}{v^2} && \text{Plug in eq. (2).} \\ &= \frac{q(t)}{v} && \text{Plug in eq. (1).} \end{aligned}$$

Thus we have

$$\frac{y}{v} = \int \frac{q(t)}{v} dt$$

Since v can be solved from eq. (2), we obtain the solution for y . □

2 Non-Linear

2.1 Separable Equations

A first order ODE is **separable** if it can be written in the form

$$M(t) + N(y) \cdot \frac{dy}{dt} = 0$$

for some functions M and N . Then the solution satisfies

$$\int N(y) dy = - \int M(t) dt$$

2.2 Transformation Methods

Some complicated non-linear ODEs can be transformed into ODEs that are solvable by suitable substitutions.

2.2.1 Bernoulli Equations

Bernoulli equations can be transformed into linear ODEs by the substitution $v = y^{1-r}$.

Definition 2.1 (Bernoulli Equations).

A first order ODE is called a **Bernoulli equation** if it can be written in the form

$$\frac{dy}{dt} + p(t) \cdot y = q(t) \cdot y^r \quad (3)$$

for some given functions p, q and a real number $r \in \mathbb{R} \setminus \{0, 1\}$.

The **general solution** of a Bernoulli equation (3) is given by

$$y^{1-r} = \frac{1-r}{\mu(t)} \int \mu(t) \cdot q(t) dt$$

where $\mu(t)$ satisfies

$$\ln |\mu(t)| = (1-r) \int p(t) dt$$

Note: If $r > 0$, then $y = 0$ is also a solution, which is not captured by the above formula.

Proof.

Substitute $v = y^{1-r}$, and then solve with the method of integrating factors.

$$\begin{aligned} \frac{dy}{dt} + p(t) \cdot y &= q(t) \cdot y^r &\implies & y^{-r} \frac{dy}{dt} + p(t) \cdot y^{1-r} = q(t) \quad \text{for } y \neq 0 \\ &&\implies & \frac{d}{dt}(y^{1-r}) + (1-r)p(t) \cdot y^{1-r} = (1-r)q(t) \end{aligned}$$

□

2.2.2 Homogeneous Equations

Homogeneous equations can be transformed into separable equations.

If a first-order ODE can be expressed in the following form, then it is said to be **homogeneous**:

$$\frac{dy}{dt} = F(y/t)$$

for some given function F .

Then it can be transformed into a separable equation using $v = y/t$:

$$\frac{1}{F(v) - v} dv = \frac{1}{t} dt$$

Remark. The homogeneous equations considered here have nothing to do with the homogeneous LCCDEs.

2.3 Exact Equations

Definition 2.2 (Exact Equations).

A first order ODE $M(t, y) dt + N(t, y) dy = 0$ is said to be **exact** if there exists a function $\Psi(t, y)$ such that

$$\frac{\partial \Psi(t, y)}{\partial t} = M(t, y) \quad \text{and} \quad \frac{\partial \Psi(t, y)}{\partial y} = N(t, y)$$

Then the general solution is given implicitly as

$$\Psi(t, y) = C \quad \text{for some constant } C \in \mathbb{R}$$

Theorem 2.1.

Let R be a simply connected region in the ty -plane. If $M(t, y)$, $N(t, y)$, $M_y(t, y)$, and $N_t(t, y)$ are continuous in R , then

$$M(t, y) + N(t, y) \frac{dy}{dt} = 0 \text{ is exact in } R \quad \iff \quad M_y(t, y) = N_t(t, y) \quad \text{for all } (t, y) \in R$$

where $M_y(t, y) := \partial M(t, y) / \partial y$ and $N_t(t, y) := \partial N(t, y) / \partial t$.

To solve the first order ODE $M(t, y) dt + N(t, y) dy = 0$, we can follow these steps:

1. Verify that the equation is exact by checking if $M_y(t, y) = N_t(t, y)$.
2. Integrate $M(t, y)$ with respect to t to find $\Psi(t, y)$:

$$\Psi(t, y) = \int M(t, y) dt + h(y)$$

where $h(y)$ is an arbitrary function of y .

3. Differentiate $\Psi(t, y)$ with respect to y and set it equal to $N(t, y)$:

$$\frac{\partial \Psi(t, y)}{\partial y} = N(t, y)$$

This will allow us to solve for $h(y)$.

4. Substitute $h(y)$ back into $\Psi(t, y)$ to obtain the implicit solution:

$$\Psi(t, y) = C$$

for some constant C .

2.4 Exact Equations and Integrating Factors

It is sometimes possible to convert an ODE that is not exact into an exact equation with a suitable integrating factor. Suppose the following first order ODE is exact:

$$\mu M(t, y) + \mu N(t, y) \frac{dy}{dt} = 0$$

Then the integrating factor μ must satisfy

$$\frac{\partial(\mu M)}{\partial y} = \frac{\partial(\mu N)}{\partial t}$$

On one hand, assuming that $\mu = \mu(t)$, we have

$$\frac{d\mu}{dt} = \frac{M_y - N_t}{N} \cdot \mu$$

If $(M_y - N_t)/N$ depends on t alone, then we can solve for $\mu(t)$.

On the other hand, assuming that $\mu = \mu(y)$, we have

$$\frac{d\mu}{dy} = \frac{N_t - M_y}{M} \cdot \mu$$

If $(N_t - M_y)/M$ depends on y alone, then we can solve for $\mu(y)$.

If neither case applies, then try some substitutions to transform the ODE into a solvable form.

3 The Existence and Uniqueness Theorem

3.1 The Theorem

Theorem 3.1 (Existence and Uniqueness Theorem for First Order Linear ODEs).

For the IVP:

$$\frac{dy}{dt} = p(t) \cdot y + q(t) \quad \text{and} \quad y(t_0) = y_0$$

if there exists an interval $I \subseteq \mathbb{R}$ such that $t_0 \in I$ and $p(t)$, $q(t)$ are **continuous** on I , then there exists a **unique** solution $y(t)$ defined on I that satisfies the IVP.

Remark. The solution is defined on the entire interval I .

To make it simpler, we let the initial point be the origin. If some other initial point is given, we can always make a preliminary change of variables.

Theorem 3.2 (Existence and Uniqueness Theorem).

For the IVP:

$$\frac{dy}{dt} = f(t, y) \quad \text{and} \quad y(0) = 0$$

if there exists a rectangle $R : |t| \leq a, |y| \leq b$ such that $f(t, y)$ and $\partial f(t, y)/\partial y$ are **continuous** on R , then there exists a **unique** solution $y = \phi(t)$ defined for $|t| \leq h$, where $0 < \min\{b/M, a\} \leq h \leq a$ and $M = \max_{(t,y) \in R} |f(t, y)| < +\infty$.

Remark. The solution is guaranteed to be defined at least on the interval $|t| \leq \min\{b/M, a\}$, where M is the bound of $f(t, y)$ on the rectangle R .

Remark. If the conditions of the theorem are not satisfied, then the solution may not be unique or may not exist.

3.2 The Preliminary before the Proof

Definition 3.1 (Pointwise Convergence).

A sequence of functions $\{f_n\}$ defined on a set $\mathcal{D} \subseteq \mathbb{R}$ is said to **converge pointwise** to a function f if:

$$\forall x \in \mathcal{D}, \lim_{n \rightarrow \infty} f_n(x) = f(x)$$

The **rate of convergence** can be **different** for different points in $\mathcal{D}$.

Definition 3.2 (Uniform Convergence).

A sequence of functions $\{f_n\}$ defined on a set $\mathcal{D} \subseteq \mathbb{R}$ is said to **converge uniformly** to a function f if:

$$\forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n > N, \forall x \in \mathcal{D}, |f_n(x) - f(x)| < \varepsilon$$

The **rate of convergence** is the **same** for all points in $\mathcal{D}$.

Uniform convergence implies pointwise convergence.

Proof.

The proof is given in first-order formal logic. The inference rules used are the following:

- $\alpha \rightarrow \beta \iff \neg\alpha \vee \beta$
- $\forall x(\alpha \vee P(x)) \iff \alpha \vee \forall x P(x)$ and $\exists x(\alpha \vee P(x)) \iff \alpha \vee \exists x P(x)$
- $\forall x \forall y P(x, y) \iff \forall y \forall x P(x, y)$ and $\exists x \forall y P(x, y) \implies \forall y \exists x P(x, y)$

To make it easier, we suppose all the variables are restricted to real numbers. Rewrite the definition of uniform convergence into first-order formal logic:

$$\begin{aligned} \text{Uniform Convergence} &\iff \forall \varepsilon \left(\varepsilon > 0 \rightarrow \exists N \left[N \in \mathbb{N} \rightarrow \forall n \left(n > N \rightarrow \forall x \left(x \in \mathcal{D} \rightarrow |f_n(x) - f(x)| < \varepsilon \right) \right) \right] \right) \\ &\iff \forall \varepsilon \left(\varepsilon \leq 0 \vee \exists N \left[N \notin \mathbb{N} \vee \forall n \left(n \leq N \vee \forall x \left(x \notin \mathcal{D} \vee |f_n(x) - f(x)| < \varepsilon \right) \right) \right] \right) \\ &\implies \forall x \forall \varepsilon \left(\varepsilon \leq 0 \vee \exists N \left[N \notin \mathbb{N} \vee \forall n \left(n \leq N \vee x \notin \mathcal{D} \vee |f_n(x) - f(x)| < \varepsilon \right) \right] \right) \\ &\iff \forall x \left[x \notin \mathcal{D} \vee \forall \varepsilon \left(\varepsilon \leq 0 \vee \exists N \left[N \notin \mathbb{N} \vee \forall n \left(n \leq N \vee |f_n(x) - f(x)| < \varepsilon \right) \right] \right) \right] \\ &\iff \forall x \left[x \in \mathcal{D} \rightarrow \forall \varepsilon \left(\varepsilon > 0 \rightarrow \exists N \left[N \in \mathbb{N} \rightarrow \forall n \left(n > N \rightarrow |f_n(x) - f(x)| < \varepsilon \right) \right] \right) \right] \\ &\iff \forall x \in \mathcal{D}, \forall \varepsilon > 0, \exists N \in \mathbb{N}, \forall n > N, |f_n(x) - f(x)| < \varepsilon \\ &\iff \text{Pointwise Convergence} \end{aligned}$$

□

Theorem 3.3 (Weierstrass M-test).

Let $\{\phi_n\}$ be a sequence of functions defined on a set $\mathcal{D} \subseteq \mathbb{R}$. If there exists a sequence of positive numbers $\{M_n\}$ such that

$$\forall n \in \mathbb{N}, \forall x \in \mathcal{D}, |\phi_n(x)| < M_n \quad \text{and} \quad \sum_{n=1}^{\infty} M_n \text{ converges}$$

then the new sequence $\{\Phi_n(x)\}$, where $\Phi_n(x) := \sum_{k=1}^n \phi_k(x)$, **converges uniformly** on $\mathcal{D}$.

Theorem 3.4.

Suppose $\{\phi_n(x)\}$ is a sequence of functions defined on an interval $[a, b] \subseteq \mathbb{R}$ and is **uniformly convergent**, then the operations of limit and integration can be interchanged:

$$\lim_{n \rightarrow \infty} \int_a^b \phi_n(x) \, dx = \int_a^b \lim_{n \rightarrow \infty} \phi_n(x) \, dx$$

3.3 The Proof