

Systems of First Order Linear ODEs

1 First order ODE systems

The general system of first order equations involving n dependent variables defined on an interval $I \subseteq \mathbb{R}$ can be written as

$$\begin{cases} y_1'(t) = F_1(t, y_1, \dots, y_n) \\ y_2'(t) = F_2(t, y_1, \dots, y_n) \\ \vdots \\ y_n'(t) = F_n(t, y_1, \dots, y_n) \end{cases} \quad \text{for } t \in I. \quad (1)$$

with initial conditions $y_1(t_0) = x_1, \quad y_2(t_0) = x_2, \quad \dots, \quad y_n(t_0) = x_n,$

The solution to the system can be expressed in vector form as

$$\mathbf{y}(t) = \begin{pmatrix} y_1(t) \\ y_2(t) \\ \vdots \\ y_n(t) \end{pmatrix}$$

Theorem 1.1 (Existence and Uniqueness).

For the system of first order ODEs (1), if the functions $F_1, F_2, \dots, F_n$ and the partial derivatives $\frac{\partial F_i}{\partial y_j}$ for all pairs $i, j = 1, 2, \dots, n$ are continuous on a region R defined by

$$R := [t_0 - a, t_0 + a] \times [x_1 - b, x_1 + b] \times \dots \times [x_n - b, x_n + b] \subseteq \mathbb{R}^{n+1},$$

where constants $a, b > 0$, then there exists a unique solution $\mathbf{y}(t)$ for $t \in (t_0 - h, t_0 + h)$ to the IVP, where

$$h = \min\{a, b/M\} \quad \text{and} \quad M = \max_{\mathbf{v} \in R} \{|F_1(\mathbf{v})|, |F_2(\mathbf{v})|, \dots, |F_n(\mathbf{v})|\}.$$

Any n -th order ODE can be rewritten as a system of n first order ODEs. Given an n -th order ODE

$$y^{(n)} = F(t, y, y', \dots, y^{(n-1)}),$$

setting $z_1 = y, z_2 = y', \dots, z_n = y^{(n-1)}$, we obtain the system

$$\begin{cases} z_1' = z_2 \\ z_2' = z_3 \\ \vdots \\ z_{n-1}' = z_n \\ z_n' = F(t, z_1, z_2, \dots, z_n) \end{cases}$$

which is a system of n first order ODEs.

Definition 1.1 (First-Order Linear System).

A system of first order ODEs is called **linear** if it can be expressed in the form

$$\mathbf{y}'(t) = \mathbf{P}(t)\mathbf{y}(t) + \mathbf{g}(t) \quad \text{for } t \in I, \quad (2)$$

where

$$\mathbf{y}(t) = \begin{pmatrix} y_1(t) \\ y_2(t) \\ \vdots \\ y_n(t) \end{pmatrix} \quad \mathbf{g}(t) = \begin{pmatrix} g_1(t) \\ g_2(t) \\ \vdots \\ g_n(t) \end{pmatrix} \quad \mathbf{P}(t) = \begin{pmatrix} p_{11}(t) & p_{12}(t) & \cdots & p_{1n}(t) \\ p_{21}(t) & p_{22}(t) & \cdots & p_{2n}(t) \\ \vdots & \vdots & \ddots & \vdots \\ p_{n1}(t) & p_{n2}(t) & \cdots & p_{nn}(t) \end{pmatrix}$$

Definition 1.2 (Homogeneous First-Order Linear System).

A first order linear system is called **homogeneous** if $\mathbf{g}(t) = \mathbf{0}_n$ for all $t \in I$, i.e., it can be expressed as

$$\mathbf{y}'(t) = \mathbf{P}(t)\mathbf{y}(t) \quad \text{for } t \in I.$$

Theorem 1.2 (Existence and Uniqueness for First-Order Linear Systems).

For the first-order linear system (2), if all entries of $\mathbf{P}(t)$ and $\mathbf{g}(t)$ are continuous on an interval I , then for any $t_0 \in I$ and $\mathbf{y}_0 \in \mathbb{R}^n$, there exists a unique solution $\mathbf{y}(t)$ to the IVP

$$\begin{cases} \mathbf{y}'(t) = \mathbf{P}(t)\mathbf{y}(t) + \mathbf{g}(t) \\ \mathbf{y}(t_0) = \mathbf{a} \in \mathbb{R}^n \end{cases} \quad \text{for } t \in I.$$

In the rest parts, we will focus on the following homogeneous system defined on an open interval $I \subseteq \mathbb{R}$:

$$\mathbf{y}'(t) = \mathbf{P}(t)\mathbf{y}(t) \quad \text{for } t \in I, \quad (3)$$

where $\mathbf{y} \in C^1(I)^n$ and $\mathbf{P} \in C(I)^{n \times n}$, i.e., all entries of $\mathbf{P}$ and $\mathbf{y}$ are continuous on I .

Theorem 1.3 (Principle of Superposition).

If $\mathbf{y}_1(t)$ and $\mathbf{y}_2(t)$ are solutions to the homogeneous first-order linear system (3), then the linear combination

$$\mathbf{y}(t) = c_1\mathbf{y}_1(t) + c_2\mathbf{y}_2(t)$$

is also a solution for any constants c_1, c_2 .

Definition 1.3 (Wronskian).

The **Wronskian** of n vector functions $\mathbf{f}_1(t), \mathbf{f}_2(t), \dots, \mathbf{f}_n(t) \in C(I)^n$ is defined as

$$W(\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)(t) := \det \left(\begin{bmatrix} | & & | \\ \mathbf{f}_1(t) & \cdots & \mathbf{f}_n(t) \\ | & & | \end{bmatrix} \right) \quad \text{for } t \in I.$$

Remark. The Wronskian for vector functions is different from the Wronskian for scalar functions. There is no derivative involved in the definition of the Wronskian for vector functions.

For any $t \in I$,

$$W(\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n)(t) \neq 0 \iff \mathbf{f}_1(t), \mathbf{f}_2(t), \dots, \mathbf{f}_n(t) \text{ are linearly independent.}$$

Theorem 1.4 (Liouville's Formula).

Let $\mathbf{y}^{(1)}(t), \mathbf{y}^{(2)}(t), \dots, \mathbf{y}^{(n)}(t)$ be n solutions to the homogeneous system (3). Then the Wronskian of these solutions is given by

$$W(\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(n)})(t) = c \cdot \exp\left(\int \operatorname{tr}(\mathbf{P}(t)) dt\right) \quad \text{for } t \in I,$$

where c is a constant not depending on t .

Remark. The **trace** of a square matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is defined as the sum of its diagonal entries:

$$\operatorname{tr}(\mathbf{A}) := a_{11} + a_{22} + \dots + a_{nn}.$$

Proof (Liouville's Formula).

Before the proof, we need to introduce several theorems from linear algebra:

1. For any square matrix $\mathbf{A}$,

$$\frac{d}{dt}(\det(\mathbf{A})) = \operatorname{tr}\left(\operatorname{adj}(\mathbf{A}) \frac{d\mathbf{A}}{dt}\right). \quad (4)$$

2. The trace remains unchanged under cycle permutations:

$$\operatorname{tr}(\mathbf{ABC}) = \operatorname{tr}(\mathbf{BCA}). \quad (5)$$

3. Property of the adjugate matrix:

$$\operatorname{adj}(\mathbf{A})\mathbf{A} = \mathbf{A} \operatorname{adj}(\mathbf{A}) = \det(\mathbf{A})\mathbf{I}. \quad (6)$$

Now, let $\mathbf{Y} := \begin{bmatrix} \mathbf{y}^{(1)} & \mathbf{y}^{(2)} & \dots & \mathbf{y}^{(n)} \end{bmatrix} \in C^1(I)^{n \times n}$ and $W := \det(\mathbf{Y})$ be the Wronskian. Then we have

$$\begin{aligned} W' &= \operatorname{tr}(\operatorname{adj}(\mathbf{Y})\mathbf{Y}') \\ &= \operatorname{tr}(\operatorname{adj}(\mathbf{Y})\mathbf{PY}) \quad \text{since } \mathbf{y}' = \mathbf{Py} \\ &= \operatorname{tr}(\mathbf{Y} \operatorname{adj}(\mathbf{Y})\mathbf{P}) \\ &= \operatorname{tr}(\mathbf{WIP}) \\ &= W \operatorname{tr}(\mathbf{P}). \end{aligned}$$

Hence the Liouville's formula. □

Theorem 1.5.

If $\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(n)}$ are n solutions to a homogeneous system (3), then

$$\forall t \in I, \quad W(\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(n)})(t) = 0 \iff \exists t_0 \in I \text{ s.t. } W(\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(n)})(t_0) = 0.$$

Proof.

Proven by Liouville's formula. □

Definition 1.4 (Fundamental Set of Solutions).

A set of n **linearly independent** solutions $\{\mathbf{y}^{(1)}(t), \mathbf{y}^{(2)}(t), \dots, \mathbf{y}^{(n)}(t)\}$ to the homogeneous system (3) is called a **fundamental set of solutions**.

Theorem 1.6.

If the n solutions $\mathbf{y}^{(1)}(t), \mathbf{y}^{(2)}(t), \dots, \mathbf{y}^{(n)}(t)$ of the homogeneous system (3) are **linearly independent** for each $t \in I$, then every solution of the system can be expressed as a linear combination:

$$\mathbf{y}(t) = c_1 \mathbf{y}^{(1)}(t) + c_2 \mathbf{y}^{(2)}(t) + \dots + c_n \mathbf{y}^{(n)}(t) \quad \text{for } t \in I$$

in **exactly one way**.

Remark. The Principle of Superposition indicates the solutions form a vector space, thus there exist bases. Therefore, the essence of this theorem is to show that the dimension of the solution space is exactly n .

Proof.

Let $\phi(t)$ be any solution to the homogeneous system and let $\mathbf{Y} := [\mathbf{y}^{(1)} \quad \mathbf{y}^{(2)} \quad \dots \quad \mathbf{y}^{(n)}]$. We pick some $t_0 \in I$ and define the constant vector $\xi := \phi(t_0) \in \mathbb{R}^n$. Since $\mathbf{y}^{(1)}(t), \dots, \mathbf{y}^{(n)}(t)$ are linearly independent for any $t \in I$, we have $\det(\mathbf{Y})(t_0) \neq 0$. Then the following equation has a unique solution $\mathbf{c} \in \mathbb{R}^n$:

$$\mathbf{Y}(t_0) \cdot \mathbf{c} = \xi \quad \text{for } \mathbf{c} \in \mathbb{R}^n.$$

Now we define the linear combination of $\mathbf{y}^{(1)}, \mathbf{y}^{(2)}, \dots, \mathbf{y}^{(n)}$ as a new vector function:

$$\eta(t) := \mathbf{Y}(t) \cdot \mathbf{c} \quad \text{for } t \in I.$$

Construct the following IVP: the homogeneous system (3) with the initial condition $\mathbf{y}(t_0) = \xi$. Since $\phi(t)$ and $\eta(t)$ are both solutions to this IVP, by the uniqueness of solutions (Theorem 1.2), we have $\phi(t) = \eta(t)$ for all $t \in I$. Thus any solution $\phi(t)$ can be expressed as a linear combination of the fundamental solutions. □