Solutions Manual to Mechanics

Hiromichi Inawashiro

May 12, 2025

Contents

1 The Equations of Motion

1

1 The Equations of Motion

1.1

Let the field be a uniform gravitational one with the acceleration g. Let the system be a coplanar double pendulum of the strings with lengths l_1 and l_2 and the particles with masses m_1 and m_2 and with the angles ϕ_1 and ϕ_2 from the y axis. Then,

$$\mathbf{r}_{1} = \begin{bmatrix} l_{1} \sin \phi_{1} \\ l_{1} \cos \phi_{1} \end{bmatrix}, \mathbf{r}_{2} = \begin{bmatrix} l_{1} \sin \phi_{1} + l_{2} \sin \phi_{2} \\ l_{1} \cos \phi_{1} + l_{2} \cos \phi_{2} \end{bmatrix}, \tag{1.1}$$

so that

$$\mathbf{v}_{1} = \begin{bmatrix} l_{1}\dot{\phi}_{1}\cos\phi_{1} \\ -l_{1}\dot{\phi}_{1}\sin\phi_{1} \end{bmatrix}, \mathbf{v}_{2} = \begin{bmatrix} l_{1}\dot{\phi}_{1}\cos\phi_{1} + l_{2}\dot{\phi}_{2}\cos\phi_{2} \\ -l_{1}\dot{\phi}_{1}\sin\phi_{1} - l_{2}\dot{\phi}_{2}\sin\phi_{2} \end{bmatrix}.$$
(1.2)

Then, the Lagrangian is given by

$$L = \frac{1}{2}m_1 \|\mathbf{v}_1\|^2 + mgr_{1y} + \frac{1}{2}m_2 \|\mathbf{v}_2\|^2 + mgr_{2y}.$$
 (1.3)

The right hand side can be written as

$$\frac{1}{2}m_1l_1^2\dot{\phi_1}^2 + m_1gl_1\cos\phi_1 + \frac{1}{2}m_2\left(l_1^2\dot{\phi_1}^2 + l_2^2\dot{\phi_2}^2 + 2l_1l_2\dot{\phi_1}\dot{\phi_2}\cos(\phi_1 - \phi_2)\right) + m_2g\left(l_1\cos\phi_1 + l_2\cos\phi_2\right).$$
(1.4)

Therefore,

$$L = \frac{1}{2}(m_1 + m_2)l_1^2 \dot{\phi_1}^2 + \frac{1}{2}m_2 l_2^2 \dot{\phi_2}^2 + m_2 l_1 l_2 \dot{\phi_1} \dot{\phi_2} \cos(\phi_1 - \phi_2) + (m_1 + m_2)g l_1 \cos \phi_1 + m_2 g l_2 \cos \phi_2.$$
(1.5)

1.2 (Incomplete)

Let the field be a uniform gravitational one with the acceleration g. Let the system be a pendulum of the string with the length l and the particles with the mass m_1 at the point of support which can move on a horisontal line and the mass m_2 at the end of the string with the angle ϕ from the y axis.