

Solutions Manual to Pattern Recognition and Machine Learning

Hiromichi Inawashiro

January 30, 2025

Contents

1	Introduction	1
2	Probability Distributions	39
3	Linear Models for Regression	93
4	Linear Models for Classification	117

1 Introduction

1.1

Let

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N (y(x_n, \mathbf{w}) - t_n)^2. \quad (1.1)$$

Setting the derivative with respect to \mathbf{w} to zero gives

$$\mathbf{0} = \sum_{n=1}^N \frac{\partial y(x_n, \mathbf{w})}{\partial \mathbf{w}} (y(x_n, \mathbf{w}) - t_n). \quad (1.2)$$

If

$$y(x_n, \mathbf{w}) = \mathbf{w}^\top \boldsymbol{\phi}(x_n), \quad (1.3)$$

then

$$\mathbf{0} = \sum_{n=1}^N \boldsymbol{\phi}(x_n) (\mathbf{w}^\top \boldsymbol{\phi}(x_n) - t_n). \quad (1.4)$$

Therefore,

$$\left(\sum_{n=1}^N \boldsymbol{\phi}(x_n) \boldsymbol{\phi}(x_n)^\top \right) \mathbf{w} = \sum_{n=1}^N t_n \boldsymbol{\phi}(x_n). \quad (1.5)$$

Thus,

$$\underset{\mathbf{w}}{\operatorname{argmin}} E(\mathbf{w}) = \mathbf{A}^{-1} \mathbf{v}, \quad (1.6)$$

where

$$\begin{aligned} \mathbf{A} &= \sum_{n=1}^N \boldsymbol{\phi}(x_n) \boldsymbol{\phi}(x_n)^\top, \\ \mathbf{v} &= \sum_{n=1}^N t_n \boldsymbol{\phi}(x_n). \end{aligned} \quad (1.7)$$

If

$$\boldsymbol{\phi}(x_n) = \begin{bmatrix} 1 \\ x_n \\ \vdots \\ x_n^M \end{bmatrix},$$

then

$$\begin{aligned} A_{mm'} &= \sum_{n=1}^N x_n^{m+m'}, \\ v_m &= \sum_{n=1}^N t_n x_n^m. \end{aligned} \tag{1.8}$$

1.2

Let

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N (y(x_n, \mathbf{w}) - t_n)^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2. \tag{1.9}$$

Setting the derivative with respect to \mathbf{w} to zero gives

$$\mathbf{0} = \sum_{n=1}^N \frac{\partial y(x_n, \mathbf{w})}{\partial \mathbf{w}} (y(x_n, \mathbf{w}) - t_n) + \lambda \mathbf{w}. \tag{1.10}$$

If

$$y(x_n, \mathbf{w}) = \mathbf{w}^\top \phi(x_n), \tag{1.11}$$

then

$$\mathbf{0} = \sum_{n=1}^N \phi(x_n) (\mathbf{w}^\top \phi(x_n) - t_n) + \lambda \mathbf{w}. \tag{1.12}$$

Therefore,

$$\left(\sum_{n=1}^N \phi(x_n) \phi(x_n)^\top + \lambda \mathbf{I} \right) \mathbf{w} = \sum_{n=1}^N t_n \phi(x_n). \tag{1.13}$$

Thus,

$$\underset{\mathbf{w}}{\operatorname{argmin}} E(\mathbf{w}) = \mathbf{A}^{-1} \mathbf{v}, \tag{1.14}$$

where

$$\begin{aligned} \mathbf{A} &= \sum_{n=1}^N \phi(x_n) \phi(x_n)^\top + \lambda \mathbf{I}, \\ \mathbf{v} &= \sum_{n=1}^N t_n \phi(x_n). \end{aligned} \tag{1.15}$$

If

$$\phi(x_n) = \begin{bmatrix} 1 \\ x_n \\ \vdots \\ x_n^M \end{bmatrix},$$

then

$$\begin{aligned} A_{mm'} &= \sum_{n=1}^N x_n^{m+m'} + \lambda I_{mm'}, \\ v_m &= \sum_{n=1}^N t_n x_n^m. \end{aligned} \tag{1.16}$$

1.3

Let a , o and l be the events where an apple, orange and lime are selected respectively. The probability that an apple is selected is given by

$$p(a) = p(a|r)p(r) + p(a|b)p(b) + p(a|g)p(g). \tag{1.17}$$

Substituting $p(a|r) = \frac{3}{10}$, $p(r) = \frac{1}{5}$, $p(a|g) = \frac{1}{2}$, $p(r) = \frac{1}{5}$, $p(a|g) = \frac{3}{10}$ and $p(g) = \frac{3}{5}$ gives

$$p(a) = \frac{17}{50}. \tag{1.18}$$

If an orange is selected, the probability that it came from the green box is given by

$$p(g|o) = \frac{p(g,o)}{p(o)}. \tag{1.19}$$

Here,

$$\begin{aligned} p(g,o) &= p(o|g)p(g), \\ p(o) &= p(o|r)p(r) + p(o|b)p(b) + p(o|g)p(g). \end{aligned} \tag{1.20}$$

Substituting $p(o|r) = \frac{2}{5}$, $p(r) = \frac{1}{5}$, $p(o|b) = \frac{1}{2}$, $p(b) = \frac{1}{5}$, $p(o|g) = \frac{3}{10}$ and $p(g) = \frac{3}{5}$ gives $p(g,o) = \frac{9}{50}$ and $p(o) = \frac{9}{25}$. Therefore,

$$p(g|o) = \frac{1}{2}. \tag{1.21}$$

1.4

Let

$$x = g(y) \tag{1.22}$$

and \hat{x} and \hat{y} be the locations of the maximum of $p_x(x)$ and $p_y(y)$ respectively. Let us assume that there exists $\epsilon > 0$ such that $g'(y) \neq 0$ for $|y - \hat{y}| < \epsilon$. Then, Taking the derivative of the transformation

$$p_y(y) = p_x(g(y)) |g'(y)| \tag{1.23}$$

and substituting $y = \hat{y}$ gives

$$0 = g'(\hat{y}) p'_x(g(\hat{y})) + p_x(g(\hat{y})) g''(\hat{y}). \tag{1.24}$$

Therefore, in general,

$$\hat{x} \neq g(\hat{y}). \tag{1.25}$$

Here, let us assume that

$$g(y) = ay + b. \tag{1.26}$$

Then, Taking the derivative of the transformation and substituting $y = \hat{y}$ gives

$$0 = p'_x(g(\hat{y})). \tag{1.27}$$

Therefore,

$$\hat{x} = g(\hat{y}). \tag{1.28}$$

1.5

By the definition,

$$\text{var } f(x) = E(f(x) - E f(x))^2. \tag{1.29}$$

The right hand side can be written as

$$E((f(x))^2 - 2f(x) E f(x) + (E f(x))^2) = E(f(x))^2 - (E f(x))^2. \tag{1.30}$$

Therefore,

$$\text{var } f(x) = E(f(x))^2 - (E f(x))^2. \tag{1.31}$$

1.6

By the definition,

$$\text{cov}(x, y) = E((x - E x)(y - E y)). \quad (1.32)$$

The right hand side can be written as

$$E xy - E(x E y) - E(y E x) + E(E x E y) = E xy - E x E y. \quad (1.33)$$

The right hand side can be written as

$$\int xyp(x, y)dxdy - \int xp(x)dx \int yp(y)dy. \quad (1.34)$$

If x and y are independent, by the definition,

$$f(x, y) = f(x)f(y). \quad (1.35)$$

Then,

$$\int xyp(x, y)dxdy = \int p(x)dx \int p(y)dy. \quad (1.36)$$

Therefore,

$$\text{cov}(x, y) = 0. \quad (1.37)$$

1.7

Let

$$I = \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}x^2\right) dx. \quad (1.38)$$

Then

$$I^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x^2 + y^2)\right) dxdy. \quad (1.39)$$

By the transformation from Cartesian coordinates (x, y) to polar coordinates (r, θ) , the right hand side can be written as

$$\int_0^{\infty} \int_0^{2\pi} \exp\left(-\frac{1}{2\sigma^2}r^2\right) \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} drd\theta = 2\pi \int_0^{\infty} \exp\left(-\frac{1}{2\sigma^2}r^2\right) r dr. \quad (1.40)$$

By the transformation $s = \frac{r}{\sigma}$, the right hand side can be written as

$$2\pi\sigma^2 \int_0^\infty \exp\left(-\frac{1}{2}s^2\right) s ds = 2\pi\sigma^2 \left[-\exp\left(-\frac{1}{2}s^2\right)\right]_0^\infty. \quad (1.41)$$

Therefore,

$$I = (2\pi\sigma^2)^{\frac{1}{2}}. \quad (1.42)$$

By the definition,

$$\mathcal{N}(x|\mu, \sigma^2) = (2\pi\sigma^2)^{-\frac{1}{2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right). \quad (1.43)$$

Then

$$\int_{-\infty}^\infty \mathcal{N}(x|\mu, \sigma^2) dx = (2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^\infty \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx. \quad (1.44)$$

By the transformation $t = x - \mu$, the right hand side can be written as

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^\infty \exp\left(-\frac{1}{2\sigma^2}t^2\right) dt = (2\pi\sigma^2)^{-\frac{1}{2}} I. \quad (1.45)$$

Therefore,

$$\int_{-\infty}^\infty \mathcal{N}(x|\mu, \sigma^2) dx = 1. \quad (1.46)$$

1.8

Let x be a variable such that

$$p(x) = \mathcal{N}(x|\mu, \sigma^2). \quad (1.47)$$

Then

$$\mathbb{E} x = \int_{-\infty}^\infty x \mathcal{N}(x|\mu, \sigma^2) dx. \quad (1.48)$$

By the definition, the right hand side can be written as

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^\infty x \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx. \quad (1.49)$$

By the transformation $y = x - \mu$, it can be written as

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^{\infty} (y + \mu) \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy. \quad (1.50)$$

Since

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^{\infty} y \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy = 0, \quad (1.51)$$

and

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^{\infty} \mu \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy = \mu \int_{-\infty}^{\infty} \mathcal{N}(y|\mu, \sigma^2) dy, \quad (1.52)$$

we have

$$\mathbb{E} x = \mu. \quad (1.53)$$

By the definition,

$$\int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) dx = 1 \quad (1.54)$$

can be written as

$$(2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx = 1. \quad (1.55)$$

Taking the derivative with respect to σ^2 gives

$$\begin{aligned} & (2\pi)^{-\frac{1}{2}} \left(-\frac{1}{2}\right) (\sigma^2)^{-\frac{3}{2}} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx \\ & + (2\pi\sigma^2)^{-\frac{1}{2}} \int_{-\infty}^{\infty} \frac{1}{2} (\sigma^2)^{-2} (x - \mu)^2 \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) dx = 0. \end{aligned} \quad (1.56)$$

The left hand side can be written as

$$\begin{aligned} & -\frac{1}{2} (\sigma^2)^{-1} \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) dx + \frac{1}{2} (\sigma^2)^{-2} \int_{-\infty}^{\infty} (x - \mu)^2 \mathcal{N}(x|\mu, \sigma^2) dx \\ & = -\frac{1}{2} (\sigma^2)^{-1} + \frac{1}{2} (\sigma^2)^{-2} \text{var } x. \end{aligned} \quad (1.57)$$

Therefore,

$$\text{var } x = \sigma^2. \quad (1.58)$$

1.9

Let

$$\mathcal{N}(x|\mu, \sigma^2) = (2\pi\sigma^2)^{-\frac{1}{2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right). \quad (1.59)$$

Setting its derivative with respect to x to zero gives

$$0 = (2\pi\sigma^2)^{-\frac{1}{2}} \left(-\frac{1}{\sigma^2}(x - \mu)\right) \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right). \quad (1.60)$$

Therefore, the mode is given by μ .

Similarly, let

$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{D}{2}} (\det \boldsymbol{\Sigma})^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right). \quad (1.61)$$

Setting its derivative with respect to \mathbf{x} to zero gives

$$\mathbf{0} = -(2\pi)^{-\frac{D}{2}} (\det \boldsymbol{\Sigma})^{-\frac{1}{2}} (\boldsymbol{\Sigma}^{-1} + (\boldsymbol{\Sigma}^{-1})^\top) (\mathbf{x} - \boldsymbol{\mu}) \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right). \quad (1.62)$$

Therefore, the mode is given by $\boldsymbol{\mu}$.

1.10

By the definition,

$$\mathbb{E}(x + y) = \int \int (x + y)p(x, y)dx dy. \quad (1.63)$$

The right hand side can be written as

$$\int x \left(\int p(x, y)dy \right) dx + \int y \left(\int p(x, y)dx \right) dy = \int xp(x)dx + \int yp(y)dy. \quad (1.64)$$

By the definition, the right hand side can be written as

$$\mathbb{E} x + \mathbb{E} y. \quad (1.65)$$

Therefore,

$$\mathbb{E}(x + y) = \mathbb{E} x + \mathbb{E} y. \quad (1.66)$$

Similarly, by the definition,

$$\text{var}(x + y) = \text{E} (x + y - \text{E}(x + y))^2 \quad (1.67)$$

By the result above and the definition, the right hand side can be written as

$$\begin{aligned} & \text{E} (x - \text{E} x)^2 + 2 \text{E} ((x - \text{E} x) (y - \text{E} y)) + \text{E} (y - \text{E} y)^2 \\ &= \text{var} x + 2 \text{cov}(x, y) + \text{var} y. \end{aligned} \quad (1.68)$$

If x and y are independent, then

$$\text{cov}(x, y) = 0, \quad (1.69)$$

by 1.6. Therefore,

$$\text{var}(x + y) = \text{var} x + \text{var} y. \quad (1.70)$$

1.11

Let x_1, \dots, x_N be variables such that

$$p(x_n) = \mathcal{N}(x_n | \mu, \sigma^2). \quad (1.71)$$

Then

$$\ln p(\mathbf{x}) = -\frac{N}{2} \ln(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu)^2. \quad (1.72)$$

To maximise it with respect to μ and σ^2 , setting the partial derivatives to zero gives

$$\begin{aligned} 0 &= \frac{1}{\sigma^2} \sum_{n=1}^N (x_n - \mu), \\ 0 &= -\frac{N}{2\sigma^2} + \frac{1}{2(\sigma^2)^2} \sum_{n=1}^N (x_n - \mu)^2. \end{aligned} \quad (1.73)$$

Therefore,

$$\begin{aligned} \mu_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N x_n, \\ \sigma_{\text{ML}}^2 &= \frac{1}{N} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2. \end{aligned} \quad (1.74)$$

1.12

Let x_m and x_n be independent variables. Then

$$\mathbb{E} x_m x_n = \mathbb{E} x_m \mathbb{E} x_n. \quad (1.75)$$

If they are samples from the Gaussian distribution with mean μ and variance σ^2 , the right hand side is given by μ^2 . On the other hand, by the definition,

$$\mathbb{E} x_n^2 = \text{var } x_n + (\mathbb{E} x_n)^2. \quad (1.76)$$

If x_n is a sample from the Gaussian distribution with mean μ and variance σ^2 , the right hand side is given by $\sigma^2 + \mu^2$. Therefore,

$$\mathbb{E} x_m x_n = \mu^2 + \delta_{mn} \sigma^2. \quad (1.77)$$

Here, since

$$\mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N x_n, \quad (1.78)$$

we have

$$\mathbb{E} \mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N \mathbb{E} x_n. \quad (1.79)$$

Therefore,

$$\mathbb{E} \mu_{\text{ML}} = \mu. \quad (1.80)$$

Similarly, since

$$\sigma_{\text{ML}}^2 = \frac{1}{N} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2, \quad (1.81)$$

we have

$$\mathbb{E} \sigma_{\text{ML}}^2 = \frac{1}{N} \sum_{n=1}^N \mathbb{E} (x_n - \mu_{\text{ML}})^2. \quad (1.82)$$

The right hand side can be written as

$$\frac{1}{N} \sum_{n=1}^N \mathbb{E} (x_n^2 - 2\mu_{\text{ML}} x_n + \mu_{\text{ML}}^2) = \frac{1}{N} \sum_{n=1}^N \mathbb{E} x_n^2 - \frac{2}{N} \mathbb{E} \left(\mu_{\text{ML}} \left(\sum_{n=1}^N x_n \right) \right) + \mathbb{E} \mu_{\text{ML}}^2. \quad (1.83)$$

The first term of the right hand side can be written as

$$\frac{1}{N} \sum_{n=1}^N (\mu^2 + \sigma^2) = \mu^2 + \sigma^2, \quad (1.84)$$

while the second and third terms can be written as

$$-2 \mathbb{E} \mu_{\text{ML}}^2 + \mathbb{E} \mu_{\text{ML}}^2 = -\mathbb{E} \mu_{\text{ML}}^2. \quad (1.85)$$

Here,

$$\mathbb{E} \mu_{\text{ML}}^2 = \mathbb{E} \left(\frac{1}{N} \sum_{n=1}^N x_n \right)^2. \quad (1.86)$$

The right hand side can be written as

$$\frac{1}{N^2} \sum_{n=1}^N \mathbb{E} x_n^2 + \frac{2}{N^2} \sum_{1 \leq m < n \leq N} \mathbb{E} x_m x_n = \frac{1}{N} (\mu^2 + \sigma^2) + \frac{N-1}{N} \mu^2. \quad (1.87)$$

Therefore,

$$\mathbb{E} \mu_{\text{ML}}^2 = \mu^2 + \frac{1}{N} \sigma^2. \quad (1.88)$$

Thus,

$$\mathbb{E} \sigma_{\text{ML}}^2 = \frac{N-1}{N} \sigma^2. \quad (1.89)$$

1.13

Let x_1, \dots, x_N be a set of variables whose mean is μ and variance is σ^2 . Then

$$\mathbb{E} \left(\frac{1}{N} \sum_{n=1}^N (x_n - \mu)^2 \right) = \frac{1}{N} \sum_{n=1}^N \mathbb{E} (x_n - \mu)^2. \quad (1.90)$$

The right hand side can be written as

$$\frac{1}{N} \sum_{n=1}^N \mathbb{E} (x_n^2 - 2\mu x_n + \mu^2) = \frac{1}{N} \sum_{n=1}^N \mathbb{E} x_n^2 - \frac{2\mu}{N} \sum_{n=1}^N \mathbb{E} x_n + \mu^2. \quad (1.91)$$

The first term of the right hand side can be written as

$$\frac{1}{N} \sum_{n=1}^N (\mu^2 + \sigma^2) = \mu^2 + \sigma^2, \quad (1.92)$$

while the second term can be written as

$$-\frac{2\mu}{N} \sum_{n=1}^N \mu = -2\mu^2. \quad (1.93)$$

Therefore,

$$\mathbb{E} \left(\frac{1}{N} \sum_{n=1}^N (x_n - \mu)^2 \right) = \sigma^2. \quad (1.94)$$

1.14

Let

$$\begin{aligned} w_{ij}^S &= \frac{1}{2}(w_{ij} + w_{ji}), \\ w_{ij}^A &= \frac{1}{2}(w_{ij} - w_{ji}). \end{aligned} \quad (1.95)$$

Then

$$\begin{aligned} w_{ij} &= w_{ij}^S + w_{ij}^A, \\ w_{ij}^S &= w_{ji}^S, \\ w_{ij}^A &= -w_{ji}^A. \end{aligned} \quad (1.96)$$

Here,

$$\sum_{i=1}^D \sum_{j=1}^D w_{ij}^A x_i x_j = \frac{1}{2} \sum_{i=1}^D \sum_{j=1}^D (w_{ij} - w_{ji}) x_i x_j. \quad (1.97)$$

The right hand side can be written as

$$\frac{1}{2} \left(\sum_{i=1}^D \sum_{j=1}^D w_{ij} x_i x_j - \sum_{i=1}^D \sum_{j=1}^D w_{ji} x_i x_j \right) = 0. \quad (1.98)$$

Therefore,

$$\sum_{i=1}^D \sum_{j=1}^D w_{ij}^A x_i x_j = 0. \quad (1.99)$$

Additionally,

$$\sum_{i=1}^D \sum_{j=1}^D w_{ij} x_i x_j = \sum_{i=1}^D \sum_{j=1}^D (w_{ij}^S + w_{ij}^A) x_i x_j. \quad (1.100)$$

The right hand side can be written as

$$\sum_{i=1}^D \sum_{j=1}^D w_{ij}^S x_i x_j + \sum_{i=1}^D \sum_{j=1}^D w_{ij}^A x_i x_j = \sum_{i=1}^D \sum_{j=1}^D w_{ij}^S x_i x_j, \quad (1.101)$$

where the result above is used. Therefore,

$$\sum_{i=1}^D \sum_{j=1}^D w_{ij} x_i x_j = \sum_{i=1}^D \sum_{j=1}^D w_{ij}^S x_i x_j. \quad (1.102)$$

Finally, since the matrix w_{ij}^S is a $D \times D$ symmetric matrix, its number of independent parameters is $\frac{D(D+1)}{2}$.

1.15

Let $n(D, M)$ be the number of independent parameters of a polynomial in D dimensions and M orders. Then

$$n(1, M) = n(1, M - 1) = 1. \quad (1.103)$$

Let us assume that

$$n(D, M) = \sum_{i=1}^D n(i, M - 1). \quad (1.104)$$

The independent terms of a polynomial in $D + 1$ dimensions and M orders can be split into 1. the ones of a polynomial in D dimensions and M orders and 2. the ones generated by multiplying the ones in $D + 1$ dimensions and M orders by the $D + 1$ th variable. Therefore,

$$n(D + 1, M) = n(D, M) + n(D + 1, M - 1). \quad (1.105)$$

Thus,

$$n(D + 1, M) = \sum_{i=1}^{D+1} n(i, M - 1). \quad (1.106)$$

Hence, the assumption is proved by induction on D .

Additionally,

$$\sum_{i=1}^1 \frac{(i + M - 2)!}{(i - 1)!(M - 1)!} = 1. \quad (1.107)$$

Let us assume that

$$\sum_{i=1}^D \frac{(i+M-2)!}{(i-1)!(M-1)!} = \frac{(D+M-1)!}{(D-1)!M!}. \quad (1.108)$$

Then

$$\sum_{i=1}^{D+1} \frac{(i+M-2)!}{(i-1)!(M-1)!} = \frac{(D+M-1)!}{(D-1)!M!} + \frac{(D+M-1)!}{D!(M-1)!}. \quad (1.109)$$

The right hand side can be written as

$$\frac{D(D+M-1)! + M(D+M-1)!}{D!M!} = \frac{(D+M)!}{D!M!}. \quad (1.110)$$

Therefore, the assumption is proved by induction on D .

Finally, by 1.14,

$$n(D, 2) = \frac{D(D+1)}{2}. \quad (1.111)$$

Let us assume that

$$n(D, M) = \frac{(D+M-1)!}{(D-1)!M!}. \quad (1.112)$$

Then, by the result above,

$$n(D, M+1) = \sum_{i=1}^D n(i, M). \quad (1.113)$$

By the assumption and result above, the right hand side can be written as

$$\sum_{i=1}^D \frac{(i+M-1)!}{(i-1)!M!} = \frac{(D+M)!}{(D-1)!(M+1)!}. \quad (1.114)$$

Therefore, the assumption is proved by induction on M .

1.16

Let $N(D, M)$ be the number of independent parameters in all of the terms up to and including the ones of D dimensions and M orders. Then, by 1.15,

$$N(D, M) = \sum_{m=0}^M n(D, m), \quad (1.115)$$

where

$$n(D, m) = \frac{(D + m - 1)!}{(D - 1)!m!}. \quad (1.116)$$

Additionally,

$$N(D, 0) = 1. \quad (1.117)$$

Let us assume that

$$\sum_{m=0}^M n(D, m) = \frac{(D + M)!}{D!M!}. \quad (1.118)$$

Then

$$\sum_{m=0}^{M+1} n(D, m) = \frac{(D + M)!}{D!M!} + \frac{(D + M)!}{(D - 1)!(M + 1)!}. \quad (1.119)$$

The right hand side can be written as

$$\frac{(M + 1)(D + M)! + D(D + M)!}{D!(M + 1)!} = \frac{(D + M + 1)!}{D!(M + 1)!}. \quad (1.120)$$

Therefore, the assumption is proved by induction on M . Thus,

$$N(D, M) = \frac{(D + M)!}{D!M!}. \quad (1.121)$$

Additionally, by the approximation

$$n! \simeq n^n \exp(-n), \quad (1.122)$$

the right hand side can be approximated as

$$\frac{(D + M)^{D+M} \exp(-(D + M))}{D^D \exp(-D) M^M \exp(-M)} = \frac{(D + M)^{D+M}}{D^D M^M}. \quad (1.123)$$

The right hand side can be written as

$$D^M \left(1 + \frac{M}{D}\right)^D \left(\frac{1}{M} + \frac{1}{D}\right)^M = M^D \left(1 + \frac{D}{M}\right)^M \left(\frac{1}{D} + \frac{1}{M}\right)^D. \quad (1.124)$$

Therefore, $N(D, M)$ can be approximated as D^M for $D \gg M$ and as M^D for $M \gg D$.

Finally, by the result above,

$$\begin{aligned} N(10, 3) &= 286, \\ N(100, 3) &= 176851, \\ N(1000, 3) &= 167668501. \end{aligned} \quad (1.125)$$

1.17

Let

$$\Gamma(x) = \int_0^\infty u^{x-1} \exp(-u) du. \quad (1.126)$$

Then

$$\Gamma(x+1) = \int_0^\infty u^x \exp(-u) du. \quad (1.127)$$

The right hand side can be written as

$$[-u^x \exp(-u)]_{u=0}^{u=\infty} + \int_0^\infty x u^{x-1} \exp(-u) du = x \Gamma(x). \quad (1.128)$$

Therefore,

$$\Gamma(x+1) = x \Gamma(x). \quad (1.129)$$

Since

$$\Gamma(1) = \int_0^1 \exp(-u) du, \quad (1.130)$$

and the right hand side can be written as 1,

$$\Gamma(1) = 0!. \quad (1.131)$$

For a positive integer x , let us assume that

$$\Gamma(x) = (x-1)!. \quad (1.132)$$

Then,

$$\Gamma(x+1) = x \Gamma(x), \quad (1.133)$$

where the right hand side can be written as $x!$. Therefore,

$$\Gamma(x+1) = x!. \quad (1.134)$$

Thus, the assumption is proved by induction on x .

1.18

Let us consider the transformation from Cartesian to polar coordinates

$$\prod_{i=1}^D \int_{-\infty}^{\infty} \exp(-x_i^2) dx_i = S_D \int_0^{\infty} \exp(-r^2) r^{D-1} dr, \quad (1.135)$$

where S_D is the surface area of a sphere of unit radius in D dimensions. By 1.7, the left hand side can be written as $\pi^{\frac{D}{2}}$. By the transformation $s = r^2$, the right hand side can be written as

$$\frac{S_D}{2} \int_0^{\infty} \exp(-s) s^{\frac{D-1}{2}} s^{-\frac{1}{2}} ds = \frac{S_D}{2} \Gamma\left(\frac{D}{2}\right). \quad (1.136)$$

Therefore,

$$S_D = \frac{2\pi^{\frac{D}{2}}}{\Gamma\left(\frac{D}{2}\right)}. \quad (1.137)$$

Additionally, the volume of the sphere can be written as

$$V_D = S_D \int_0^1 r^{D-1} dr. \quad (1.138)$$

The right hand side can be written as

$$S_D \left[\frac{r^D}{D} \right]_{r=0}^{r=1} = \frac{S_D}{D}. \quad (1.139)$$

Therefore,

$$V_D = \frac{S_D}{D}. \quad (1.140)$$

Finally, the results above reduce to

$$\begin{aligned} S_2 &= \frac{2\pi}{\Gamma(1)}, \\ V_2 &= \frac{S_2}{2}. \end{aligned} \quad (1.141)$$

Therefore,

$$\begin{aligned} S_2 &= 2\pi, \\ V_2 &= \pi. \end{aligned} \quad (1.142)$$

Similarly,

$$\begin{aligned} S_3 &= \frac{2\pi^{\frac{3}{2}}}{\Gamma\left(\frac{3}{2}\right)}, \\ V_3 &= \frac{S_3}{3}. \end{aligned} \tag{1.143}$$

Therefore,

$$\begin{aligned} S_3 &= 4\pi, \\ V_3 &= \frac{4}{3}\pi. \end{aligned} \tag{1.144}$$

1.19

The volume of a cube of side 2 in D dimensions is 2^D . Therefore, the ratio of the volume of the cocentric sphere of radius 1 divided by the volume of the cube is given by

$$\frac{V_D}{2^D} = \frac{\pi^{\frac{D}{2}}}{D2^{D-1}\Gamma\left(\frac{D}{2}\right)}, \tag{1.145}$$

by 1.18.

Additionally, by Sterling's formula

$$\Gamma(x+1) \simeq (2\pi)^{\frac{1}{2}} \exp(-x) x^{\frac{x+1}{2}}, \tag{1.146}$$

the ratio can be approximated as

$$\frac{V_D}{2^D} \simeq \frac{\pi^{\frac{D}{2}}}{D2^{D-1}(2\pi)^{\frac{1}{2}} \exp\left(1 - \frac{D}{2}\right) \left(\frac{D}{2} - 1\right)^{\frac{D}{4}}}. \tag{1.147}$$

The right hand side can be written as

$$\frac{1}{2e(2\pi)^{\frac{1}{2}}} \frac{1}{D} \left(\frac{e^2\pi^2}{8D-16} \right)^{\frac{D}{4}}. \tag{1.148}$$

Therefore, the ratio goes to zero as $D \rightarrow \infty$.

Finally, the ratio of the distance from the center of the cube to one of the corners divided by the perpendicular distance to one of the sides is given by

$$\frac{\sqrt{\sum_{i=1}^D 1^2}}{1} = \sqrt{D}. \tag{1.149}$$

Therefore, the ration goes to ∞ as $D \rightarrow \infty$.

1.20

For a vector \mathbf{x} in D dimensions, let

$$p(\mathbf{x}) = (2\pi\sigma^2)^{-\frac{D}{2}} \exp\left(-\frac{\|\mathbf{x}\|^2}{2\sigma^2}\right). \quad (1.150)$$

Then

$$\int_{r \leq \|\mathbf{x}\| \leq r+\epsilon} p(\mathbf{x}) d\mathbf{x} = \int_r^{r+\epsilon} \int (2\pi\sigma^2)^{-\frac{D}{2}} \exp\left(-\frac{r'^2}{2\sigma^2}\right) J dr' d\phi, \quad (1.151)$$

where ϕ is the vector of the angular components of the polar coordinate and J is the Jacobian of the transformation from the Cartesian to polar coordinate. For a sufficiently small ϵ , the right hand side can be approximated as

$$\begin{aligned} & (2\pi\sigma^2)^{-\frac{D}{2}} \exp\left(-\frac{r^2}{2\sigma^2}\right) \int_r^{r+\epsilon} \int J dr' d\phi \\ &= (2\pi\sigma^2)^{-\frac{D}{2}} \exp\left(-\frac{r^2}{2\sigma^2}\right) \int_{r \leq \|\mathbf{x}\| \leq r+\epsilon} d\mathbf{x}. \end{aligned} \quad (1.152)$$

Therefore,

$$\int_{r \leq \|\mathbf{x}\| \leq r+\epsilon} p(\mathbf{x}) d\mathbf{x} \simeq p(r)\epsilon, \quad (1.153)$$

where

$$p(r) = (2\pi\sigma^2)^{-\frac{D}{2}} S_D r^{D-1} \exp\left(-\frac{r^2}{2\sigma^2}\right), \quad (1.154)$$

and S_D is the surface area of a unit sphere in D dimensions.

Additionally, setting the derivative of $p(r)$ to zero gives

$$0 = (2\pi\sigma^2)^{-\frac{D}{2}} S_D \left((D-1)r^{D-2} - \frac{r^D}{\sigma^2} \right) \exp\left(-\frac{r^2}{2\sigma^2}\right). \quad (1.155)$$

Therefore, $p(r)$ is maximised at a single stationary point

$$\hat{r} = \sqrt{D-1}\sigma. \quad (1.156)$$

Additionally, by the expression of $p(r)$ above,

$$\frac{p(\hat{r} + \epsilon)}{p(\hat{r})} = \left(\frac{\hat{r} + \epsilon}{\hat{r}} \right)^{D-1} \exp\left(-\frac{2\hat{r}\epsilon + \epsilon^2}{2\sigma^2}\right). \quad (1.157)$$

Using the expression of \hat{r} above, the right hand side can be written as

$$\begin{aligned} & \exp \left((D-1) \ln \left(1 + \frac{\epsilon}{\hat{r}} \right) - \frac{2\hat{r}\epsilon + \epsilon^2}{2\sigma^2} \right) \\ &= \exp \left(\frac{\hat{r}^2}{\sigma^2} \ln \left(1 + \frac{\epsilon}{\hat{r}} \right) - \frac{2\hat{r}\epsilon + \epsilon^2}{2\sigma^2} \right). \end{aligned} \quad (1.158)$$

By the Taylor series

$$\ln(1+x) = x - \frac{1}{2}x^2 + o(x^3), \quad (1.159)$$

the right hand side can be approximated as

$$\exp \left(\frac{\hat{r}^2}{\sigma^2} \left(\frac{\epsilon}{\hat{r}} - \frac{\epsilon^2}{2\hat{r}^2} \right) - \frac{2\hat{r}\epsilon + \epsilon^2}{2\sigma^2} \right) = \exp \left(-\frac{\epsilon^2}{\sigma^2} \right). \quad (1.160)$$

Therefore,

$$p(\hat{r} + \epsilon) \simeq p(\hat{r}) \exp \left(-\frac{\epsilon^2}{\sigma^2} \right). \quad (1.161)$$

Finally, let a vector of length \hat{r} be $\hat{\mathbf{r}}$. Then, by the definition of $p(\mathbf{x})$,

$$\frac{p(\mathbf{0})}{p(\hat{\mathbf{r}})} = \exp \left(\frac{\hat{r}^2}{2\sigma^2} \right). \quad (1.162)$$

Substituting the expression of \hat{r} above, the right hand side can be written as $\exp \left(\frac{D-1}{2} \right)$. Therefore,

$$\frac{p(\mathbf{0})}{p(\hat{\mathbf{r}})} = \exp \left(\frac{D-1}{2} \right). \quad (1.163)$$

1.21

If $0 \leq a \leq b$, then

$$0 \leq a(b-a). \quad (1.164)$$

Therefore,

$$a \leq (ab)^{\frac{1}{2}}. \quad (1.165)$$

For a two-class classification problem of \mathbf{x} , let the classes be \mathcal{C}_1 and \mathcal{C}_2 and let the decision regions be \mathcal{R}_1 and \mathcal{R}_2 . Let us choose the decision regions to minimise the probability of misclassification. Then,

$$p(\mathbf{x}, \mathcal{C}_1) > p(\mathbf{x}, \mathcal{C}_2) \Rightarrow \mathbf{x} \in \mathcal{C}_1, \quad (1.166)$$

and

$$p(\mathbf{x}, \mathcal{C}_2) > p(\mathbf{x}, \mathcal{C}_1) \Rightarrow \mathbf{x} \in \mathcal{C}_2. \quad (1.167)$$

Then, using the inequality above,

$$\int_{\mathcal{R}_1} p(\mathbf{x}, \mathcal{C}_2) d\mathbf{x} \leq \int_{\mathcal{R}_1} (p(\mathbf{x}, \mathcal{C}_1) p(\mathbf{x}, \mathcal{C}_2))^{\frac{1}{2}} d\mathbf{x}, \quad (1.168)$$

and

$$\int_{\mathcal{R}_2} p(\mathbf{x}, \mathcal{C}_1) d\mathbf{x} \leq \int_{\mathcal{R}_2} (p(\mathbf{x}, \mathcal{C}_1) p(\mathbf{x}, \mathcal{C}_2))^{\frac{1}{2}} d\mathbf{x}. \quad (1.169)$$

Therefore,

$$\int_{\mathcal{R}_1} p(\mathbf{x}, \mathcal{C}_2) d\mathbf{x} + \int_{\mathcal{R}_2} p(\mathbf{x}, \mathcal{C}_1) d\mathbf{x} \leq \int (p(\mathbf{x}, \mathcal{C}_1) p(\mathbf{x}, \mathcal{C}_2))^{\frac{1}{2}} d\mathbf{x}. \quad (1.170)$$

1.22

Let

$$E L = \sum_k \sum_j \int_{\mathcal{R}_j} L_{kj} p(\mathbf{x}, \mathcal{C}_k) d\mathbf{x}. \quad (1.171)$$

If

$$L_{kj} = 1 - \delta_{kj}, \quad (1.172)$$

then the right hand side can be written as

$$\sum_k \sum_j \int_{\mathcal{R}_j} (p(\mathbf{x}, \mathcal{C}_k) - p(\mathbf{x}, \mathcal{C}_j)) d\mathbf{x} = \sum_j \int_{\mathcal{R}_j} \left(\sum_k p(\mathbf{x}, \mathcal{C}_k) - p(\mathbf{x}, \mathcal{C}_j) \right) d\mathbf{x}. \quad (1.173)$$

The right hand side can be written as

$$\sum_j \int_{\mathcal{R}_j} (p(\mathbf{x}) - p(\mathbf{x}, \mathcal{C}_j)) d\mathbf{x} = 1 - \sum_j \int_{\mathcal{R}_j} p(\mathbf{x}, \mathcal{C}_j) d\mathbf{x}. \quad (1.174)$$

Therefore,

$$E L = 1 - \sum_j \int_{\mathcal{R}_j} p(\mathcal{C}_j | \mathbf{x}) p(\mathbf{x}) d\mathbf{x}. \quad (1.175)$$

Thus, minimising $E L$ reduces to choosing the criterion to maximise the posterior probability $p(\mathcal{C}_j | \mathbf{x})$.

1.23

Let

$$E L = \sum_k \sum_j \int_{\mathcal{R}_j} L_{kj} p(\mathbf{x}, \mathcal{C}_k) d\mathbf{x}. \quad (1.176)$$

The right hand side can be written as

$$\sum_j \int_{\mathcal{R}_j} \sum_k L_{kj} p(\mathbf{x}, \mathcal{C}_k) d\mathbf{x} = \sum_j \int_{\mathcal{R}_j} \left(\sum_k L_{kj} p(\mathcal{C}_k | \mathbf{x}) \right) p(\mathbf{x}) d\mathbf{x}. \quad (1.177)$$

Therefore,

$$E L = \sum_j \int_{\mathcal{R}_j} \left(\sum_k L_{kj} p(\mathcal{C}_k | \mathbf{x}) \right) p(\mathbf{x}) d\mathbf{x}. \quad (1.178)$$

Thus, minimising $E L$ reduces to choosing to minimise $\sum_k L_{kj} p(\mathcal{C}_k | \mathbf{x})$.

1.24 (Incomplete)

Let

$$E L = \sum_k \sum_j \int_{\mathcal{R}_j} L_{kj} p(\mathbf{x}, \mathcal{C}_k) d\mathbf{x} + \lambda \int_{\forall k p(\mathcal{C}_k | \mathbf{x}) < \theta} p(\mathbf{x}) d\mathbf{x}. \quad (1.179)$$

1.25

Let

$$E L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = \int \int \|\mathbf{y}(\mathbf{x}) - \mathbf{t}\|^2 p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t}. \quad (1.180)$$

Setting the derivative with respect to $\mathbf{y}(\mathbf{x})$ to zero gives

$$\mathbf{0} = 2 \int (\mathbf{y}(\mathbf{x}) - \mathbf{t}) p(\mathbf{x}, \mathbf{t}) d\mathbf{t}. \quad (1.181)$$

The integral of the right hand side can be written as

$$\mathbf{y}(\mathbf{x}) \int p(\mathbf{x}, \mathbf{t}) d\mathbf{t} - \int \mathbf{t} p(\mathbf{x}, \mathbf{t}) d\mathbf{t} = \mathbf{y}(\mathbf{x}) p(\mathbf{x}) - p(\mathbf{x}) \int \mathbf{t} p(\mathbf{t} | \mathbf{x}) d\mathbf{t}. \quad (1.182)$$

The integral in the second term of the right hand side can be written as $E_{\mathbf{t}}(\mathbf{t} | \mathbf{x})$. Therefore, the right hand side can be written as

$$\mathbf{0} = p(\mathbf{x}) (\mathbf{y}(\mathbf{x}) - E_{\mathbf{t}}(\mathbf{t} | \mathbf{x})). \quad (1.183)$$

Thus,

$$\operatorname{argmin}_{\mathbf{y}(\mathbf{x})} \mathbb{E} L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}). \quad (1.184)$$

Finally, for a single target variable t , it reduces to

$$\operatorname{argmin}_{\mathbf{y}(\mathbf{x})} \mathbb{E} L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = \mathbb{E}_t(t|\mathbf{x}). \quad (1.185)$$

1.26

Let

$$\mathbb{E} L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = \int \int \|\mathbf{y}(\mathbf{x}) - \mathbf{t}\|^2 p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t}. \quad (1.186)$$

The right hand side can be written as

$$\begin{aligned} & \int \int \|\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) + \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) - \mathbf{t}\|^2 p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t} \\ &= \int \int \|\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x})\|^2 p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t} \\ & \quad + 2 \int \int (\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}))^\top (\mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) - \mathbf{t}) p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t} \\ & \quad + \int \int \|\mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) - \mathbf{t}\|^2 p(\mathbf{x}, \mathbf{t}) d\mathbf{x} d\mathbf{t}. \end{aligned} \quad (1.187)$$

Let us look at each term of the right hand side. The first term can be written as

$$\int \|\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x})\|^2 \left(\int p(\mathbf{x}, \mathbf{t}) d\mathbf{t} \right) d\mathbf{x} = \int \|\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x})\|^2 p(\mathbf{x}) d\mathbf{x}. \quad (1.188)$$

The second term can be written as

$$2 \int (\mathbf{y}(\mathbf{x}) - \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}))^\top \left(\int (\mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) - \mathbf{t}) p(\mathbf{t}|\mathbf{x}) d\mathbf{t} \right) p(\mathbf{x}) d\mathbf{x}. \quad (1.189)$$

Since

$$\begin{aligned} \int \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) p(\mathbf{t}|\mathbf{x}) d\mathbf{t} &= \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) \frac{\int p(\mathbf{x}, \mathbf{t}) d\mathbf{t}}{p(\mathbf{x})}, \\ \int \mathbf{t} p(\mathbf{t}|\mathbf{x}) d\mathbf{t} &= \mathbb{E}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}), \end{aligned} \quad (1.190)$$

the second term is zero. The third term can be written as

$$\int \left(\int \|E_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) - \mathbf{t}\|^2 p(\mathbf{t}|\mathbf{x}) d\mathbf{t} \right) p(\mathbf{x}) d\mathbf{x} = \int \text{var}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) p(\mathbf{x}) d\mathbf{x}. \quad (1.191)$$

Therefore,

$$E L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = \int \|\mathbf{y}(\mathbf{x}) - E_{\mathbf{t}}(\mathbf{t}|\mathbf{x})\|^2 p(\mathbf{x}) d\mathbf{x} + \int \text{var}_{\mathbf{t}}(\mathbf{t}|\mathbf{x}) p(\mathbf{x}) d\mathbf{x}. \quad (1.192)$$

Thus,

$$\underset{\mathbf{y}(\mathbf{x})}{\text{argmin}} E L(\mathbf{t}, \mathbf{y}(\mathbf{x})) = E_{\mathbf{t}}(\mathbf{t}|\mathbf{x}). \quad (1.193)$$

1.27 (Incomplete)

Let

$$E L_q = \int \int |y(\mathbf{x}) - t|^q p(\mathbf{x}, t) d\mathbf{x} dt. \quad (1.194)$$

Setting the derivative with respect to $y(\mathbf{x})$ to zero gives

$$0 = q p(\mathbf{x}) \int |y(\mathbf{x}) - t|^{q-1} \text{sign}(y(\mathbf{x}) - t) p(t|\mathbf{x}) dt. \quad (1.195)$$

Therefore,

$$\underset{y(\mathbf{x})}{\text{argmin}} E L_q = \left\{ y(\mathbf{x}) \mid \int |y(\mathbf{x}) - t|^{q-1} \text{sign}(y(\mathbf{x}) - t) p(t|\mathbf{x}) dt = 0 \right\}. \quad (1.196)$$

Additionally, if $q = 1$, the integral can be written as

$$p(\mathbf{x}) \int \text{sign}(y(\mathbf{x}) - t) p(t|\mathbf{x}) dt = p(\mathbf{x}) \left(\int_{y(\mathbf{x})}^{\infty} p(t|\mathbf{x}) dt - \int_{-\infty}^{y(\mathbf{x})} p(t|\mathbf{x}) dt \right). \quad (1.197)$$

Therefore,

$$\underset{y(\mathbf{x})}{\text{argmin}} E L_1 = \text{median}(t|\mathbf{x}). \quad (1.198)$$

Finally,

$$\lim_{q \rightarrow 0} \left(\underset{y(\mathbf{x})}{\text{argmin}} E L_q \right) = \text{mode}(t|\mathbf{x})? \quad (1.199)$$

1.28

Let us assume that

$$p(x, y) = p(x)p(y) \Rightarrow h(x, y) = h(x) + h(y). \quad (1.200)$$

Let $h(p)$ be a function to relate h and p . Then

$$h(p^2) = 2h(p). \quad (1.201)$$

Let us assume that, for a positive integer n ,

$$h(p^n) = nh(p). \quad (1.202)$$

Then, by the first assumption,

$$h(p^{n+1}) = h(p^n) + h(p). \quad (1.203)$$

Therefore,

$$h(p^{n+1}) = (n+1)h(p). \quad (1.204)$$

Thus, the second assumption is proved by induction on n .

Additionally, for positive integers m and n ,

$$h(p^n) = h(p^{\frac{n}{m}m}). \quad (1.205)$$

By the second assumption, the left hand side can be written as $nh(p)$. By the first assumption, the right hand side can be written as $mh(p^{\frac{n}{m}})$. Therefore,

$$h(p^{\frac{n}{m}}) = \frac{n}{m}h(p). \quad (1.206)$$

Finally, by the continuity, for a positive real number a ,

$$h(p^a) = ah(p). \quad (1.207)$$

Taking the derivative with respect to a and substituting $a = 1$ gives

$$(p \ln p)h'(p) = h(p). \quad (1.208)$$

Therefore,

$$\int \frac{h'(p)}{h(p)} dp = \int \frac{1}{p \ln p} dp + \text{const}. \quad (1.209)$$

Ignoring the constants, the left hand side can be written as $\ln h(p)$ and the right hand side can be written as $\ln(\ln p)$. Thus,

$$h(p) \propto \ln p. \quad (1.210)$$

1.29

Let x be an M -state discrete random variable. Then, by the definition,

$$H(x) = - \sum_{i=1}^M p(x_i) \ln p(x_i), \quad (1.211)$$

where

$$\sum_{i=1}^M p(x_i) = 1. \quad (1.212)$$

By Jensen's inequality,

$$\sum_{i=1}^M p(x_i) \ln \frac{1}{p(x_i)} \leq \ln \left(\sum_{i=1}^M 1 \right). \quad (1.213)$$

Therefore,

$$H(x) \leq \ln M. \quad (1.214)$$

1.30

Let

$$\begin{aligned} p(x) &= \mathcal{N}(x|\mu, \sigma^2), \\ q(x) &= \mathcal{N}(x|m, s^2). \end{aligned} \quad (1.215)$$

By the definition,

$$\text{KL}(p||q) = - \int p(x) \ln \frac{q(x)}{p(x)} dx. \quad (1.216)$$

The right hand side can be written as

$$\begin{aligned} & - \int_{-\infty}^{\infty} p(x) \ln \frac{(2\pi s^2)^{-\frac{1}{2}} \exp\left(-\frac{(x-m)^2}{2s^2}\right)}{(2\pi \sigma^2)^{-\frac{1}{2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)} dx \\ &= - \int_{-\infty}^{\infty} p(x) \left(-\frac{1}{2} \ln \frac{s^2}{\sigma^2} - \frac{(x-m)^2}{2s^2} + \frac{(x-\mu)^2}{2\sigma^2} \right) dx. \end{aligned} \quad (1.217)$$

The right hand side can be written as

$$\ln \frac{s}{\sigma} \int_{-\infty}^{\infty} p(x) dx + \frac{1}{2s^2} \int_{-\infty}^{\infty} (x-m)^2 p(x) dx - \frac{1}{2\sigma^2} \int_{-\infty}^{\infty} (x-\mu)^2 p(x) dx. \quad (1.218)$$

The first term can be written as $\ln \frac{s}{\sigma}$. The second term can be written as

$$\frac{1}{2s^2} \int_{-\infty}^{\infty} (x - \mu + \mu - m)^2 p(x) dx = \frac{\sigma^2 + (\mu - m)^2}{2s^2}. \quad (1.219)$$

The third term can be written as $-\frac{1}{2}$. Therefore,

$$\text{KL}(p||q) = \ln \frac{s}{\sigma} + \frac{\sigma^2 + (\mu - m)^2}{2s^2} - \frac{1}{2}. \quad (1.220)$$

1.31

Let \mathbf{x} and \mathbf{y} be two variables. Then, by the definition,

$$\begin{aligned} H(\mathbf{x}) &= - \int p(\mathbf{x}) \ln p(\mathbf{x}) d\mathbf{x}, \\ H(\mathbf{y}) &= - \int p(\mathbf{y}) \ln p(\mathbf{y}) d\mathbf{y}, \\ H(\mathbf{x}, \mathbf{y}) &= - \int \int p(\mathbf{x}, \mathbf{y}) \ln p(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y}. \end{aligned} \quad (1.221)$$

Note that

$$\begin{aligned} H(\mathbf{x}) &= - \int \left(\int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} \right) \ln p(\mathbf{x}) d\mathbf{x}, \\ H(\mathbf{y}) &= - \int \left(\int p(\mathbf{x}, \mathbf{y}) d\mathbf{x} \right) \ln p(\mathbf{y}) d\mathbf{y}. \end{aligned} \quad (1.222)$$

Therefore,

$$H(\mathbf{x}) + H(\mathbf{y}) - H(\mathbf{x}, \mathbf{y}) = - \int \int p(\mathbf{x}, \mathbf{y}) \ln \frac{p(\mathbf{x})p(\mathbf{y})}{p(\mathbf{x}, \mathbf{y})} d\mathbf{x} d\mathbf{y}. \quad (1.223)$$

Since

$$\int \int p(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y} = 1, \quad (1.224)$$

Jensen's inequality can be used to write that

$$- \int \int p(\mathbf{x}, \mathbf{y}) \ln \frac{p(\mathbf{x})p(\mathbf{y})}{p(\mathbf{x}, \mathbf{y})} d\mathbf{x} d\mathbf{y} \geq - \ln \left(\int \int p(\mathbf{x})p(\mathbf{y}) d\mathbf{x} d\mathbf{y} \right). \quad (1.225)$$

The right hand side can be written as

$$- \ln \left(\int p(\mathbf{x}) d\mathbf{x} \int p(\mathbf{y}) d\mathbf{y} \right) = 0. \quad (1.226)$$

Thus,

$$H(\mathbf{x}, \mathbf{y}) \leq H(\mathbf{x}) + H(\mathbf{y}). \quad (1.227)$$

1.32

Let \mathbf{x} be a vector of continuous variables and

$$\mathbf{y} = \mathbf{A}\mathbf{x}, \quad (1.228)$$

where \mathbf{A} is a nonsingular matrix. By the definition,

$$H(\mathbf{y}) = - \int p_y(\mathbf{y}) \ln p_y(\mathbf{y}) d\mathbf{y}. \quad (1.229)$$

By the transformation

$$p_y(\mathbf{y}) = p_x(\mathbf{A}\mathbf{x}) |\det \mathbf{A}^{-1}|, \quad (1.230)$$

the right hand side can be written as

$$- \int p_x(\mathbf{A}\mathbf{x}) \ln p_x(\mathbf{A}\mathbf{x}) |\det \mathbf{A}| d\mathbf{x} - \ln |\det \mathbf{A}^{-1}| \int p_y(\mathbf{y}) d\mathbf{y}. \quad (1.231)$$

By the transformation

$$\mathbf{x}' = \mathbf{A}\mathbf{x}, \quad (1.232)$$

the first term can be written as

$$- \int p_x(\mathbf{x}') \ln p_x(\mathbf{x}') d\mathbf{x}' = H(\mathbf{x}), \quad (1.233)$$

and the second term can be written as

$$- \ln |\det \mathbf{A}^{-1}| = \ln |\det \mathbf{A}|. \quad (1.234)$$

Therefore,

$$H(\mathbf{y}) = H(\mathbf{x}) + \ln |\det \mathbf{A}|. \quad (1.235)$$

1.33

Let x and y be two discrete variables. By the definition,

$$H(y|x) = - \sum_i \sum_j p(x_i, y_j) \ln p(y_j|x_i). \quad (1.236)$$

If $H(y|x)$ is zero, then

$$0 = - \sum_i p(x_i) \sum_j p(y_j|x_i) \ln p(y_j|x_i). \quad (1.237)$$

Since

$$\begin{aligned} p(x_i) &\geq 0, \\ p(y_j|x_i) \ln p(y_j|x_i) &\leq 0. \end{aligned} \quad (1.238)$$

for all i and j , the equation reduces to

$$p(y_j|x_i) \ln p(y_j|x_i) = 0. \quad (1.239)$$

Therefore, $p(y_j|x_i)$ is zero or one. Thus, since

$$\sum_j p(y_j|x_i) = 1, \quad (1.240)$$

$p(y_j|x_i)$ is one for a unique x_i and zero for others.

1.34

Let

$$\begin{aligned} L(p(x)) = & - \int_{-\infty}^{\infty} p(x) \ln p(x) dx + \lambda_1 \left(\int_{-\infty}^{\infty} p(x) dx - 1 \right) \\ & + \lambda_2 \left(\int_{-\infty}^{\infty} xp(x) dx - \mu \right) + \lambda_3 \left(\int_{-\infty}^{\infty} (x - \mu)^2 p(x) dx - \sigma^2 \right). \end{aligned} \quad (1.241)$$

Setting the derivative with respect to $p(x)$ to zero gives

$$0 = -\ln p(x) - 1 + \lambda_1 + \lambda_2 x + \lambda_3 (x - \mu)^2. \quad (1.242)$$

Therefore,

$$p(x) = \exp \left(-1 + \lambda_1 + \lambda_2 x + \lambda_3 (x - \mu)^2 \right). \quad (1.243)$$

Therefore,

$$p(x) = \exp \left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3} + \lambda_3 \left(x - \left(\mu - \frac{\lambda_2}{2\lambda_3} \right) \right)^2 \right). \quad (1.244)$$

Substituting it to

$$\begin{aligned}\int_{-\infty}^{\infty} p(x) dx &= 1, \\ \int_{-\infty}^{\infty} xp(x) dx &= \mu, \\ \int_{-\infty}^{\infty} (x - \mu)^2 p(x) dx &= \sigma^2,\end{aligned}\tag{1.245}$$

gives

$$\begin{aligned}\exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} \exp\left(\lambda_3 \left(x - \left(\mu - \frac{\lambda_2}{2\lambda_3}\right)\right)^2\right) dx &= 1, \\ \exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} x \exp\left(\lambda_3 \left(x - \left(\mu - \frac{\lambda_2}{2\lambda_3}\right)\right)^2\right) dx &= \mu, \\ \exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} (x - \mu)^2 \exp\left(\lambda_3 \left(x - \left(\mu - \frac{\lambda_2}{2\lambda_3}\right)\right)^2\right) dx &= \sigma^2.\end{aligned}\tag{1.246}$$

By the transformation

$$y = \sqrt{-\lambda_3} \left(x - \left(\mu - \frac{\lambda_2}{2\lambda_3}\right)\right),\tag{1.247}$$

they can be written as

$$\begin{aligned}\exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} \exp(-y^2) (-\lambda_3)^{-\frac{1}{2}} dy &= 1, \\ \exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} \left((- \lambda_3)^{-\frac{1}{2}} y + \mu - \frac{\lambda_2}{2\lambda_3}\right) \exp(-y^2) (-\lambda_3)^{-\frac{1}{2}} dy &= \mu, \\ \exp\left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3}\right) \int_{-\infty}^{\infty} \left((- \lambda_3)^{-\frac{1}{2}} y - \frac{\lambda_2}{2\lambda_3}\right)^2 \exp(-y^2) (-\lambda_3)^{-\frac{1}{2}} dy &= \sigma^2.\end{aligned}\tag{1.248}$$

Since

$$\begin{aligned}\int_{-\infty}^{\infty} \exp(-y^2) dy &= \Gamma\left(\frac{1}{2}\right), \\ \int_{-\infty}^{\infty} y \exp(-y^2) dy &= 0, \\ \int_{-\infty}^{\infty} y^2 \exp(-y^2) dy &= \Gamma\left(\frac{3}{2}\right),\end{aligned}\tag{1.249}$$

they can be written as

$$\begin{aligned}
& \exp \left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3} \right) (-\lambda_3)^{-\frac{1}{2}} \Gamma \left(\frac{1}{2} \right) = 1, \\
& \exp \left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3} \right) \left(\mu - \frac{\lambda_2}{2\lambda_3} \right) (-\lambda_3)^{-\frac{1}{2}} \Gamma \left(\frac{1}{2} \right) = \mu, \\
& \exp \left(-1 + \lambda_1 - \frac{\lambda_2^2}{4\lambda_3} \right) \left((-\lambda_3)^{-\frac{3}{2}} \Gamma \left(\frac{3}{2} \right) + (-\lambda_3)^{-\frac{1}{2}} \frac{\lambda_2^2}{4\lambda_3^2} \Gamma \left(\frac{1}{2} \right) \right) = \sigma^2.
\end{aligned} \tag{1.250}$$

Therefore,

$$\begin{aligned}
\lambda_1 &= 1 - \frac{1}{2} \ln (2\pi\sigma^2), \\
\lambda_2 &= 0, \\
\lambda_3 &= -\frac{1}{2\sigma^2}.
\end{aligned} \tag{1.251}$$

Thus,

$$p(x) = (2\pi\sigma^2)^{-\frac{1}{2}} \exp \left(-\frac{1}{2\sigma^2} (x - \mu)^2 \right). \tag{1.252}$$

1.35

Let x be a variable such that

$$p(x) = \mathcal{N}(x|\mu, \sigma^2). \tag{1.253}$$

Then, by the definition,

$$H(x) = - \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) \ln \mathcal{N}(x|\mu, \sigma^2) dx. \tag{1.254}$$

The right hand side can be written as

$$\begin{aligned}
& - \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) \left(-\frac{1}{2} \ln (2\pi\sigma^2) - \frac{1}{2\sigma^2} (x - \mu)^2 \right) dx \\
&= \frac{1}{2} \ln (2\pi\sigma^2) \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) dx + \frac{1}{2\sigma^2} \int_{-\infty}^{\infty} (x - \mu)^2 \mathcal{N}(x|\mu, \sigma^2) dx.
\end{aligned} \tag{1.255}$$

Therefore,

$$H(x) = \frac{1}{2} (1 + \ln (2\pi\sigma^2)). \tag{1.256}$$

1.36 (Incomplete)

Let f be a strictly convex function. Then, by the definition,

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b), \quad (1.257)$$

where $a \leq b$ and $0 \leq \lambda \leq 1$. Let

$$x = \lambda a + (1 - \lambda)b. \quad (1.258)$$

Then, the inequality can be written as

$$f(x) \leq \frac{b-x}{b-a}f(a) + \frac{x-a}{b-a}f(b). \quad (1.259)$$

Let

$$g(x) = \frac{b-x}{b-a}f(a) + \frac{x-a}{b-a}f(b) - f(x). \quad (1.260)$$

Then,

$$g(x) \geq 0. \quad (1.261)$$

Additionally, for $x > a$,

$$g(x) = (x-a) \left(\frac{f(b)-f(a)}{b-a} - \frac{f(x)-f(a)}{x-a} \right). \quad (1.262)$$

By the mean value theorem, there exists c and y such that $a \leq c \leq b$, $a \leq y \leq x$ and

$$\begin{aligned} f'(c) &= \frac{f(b)-f(a)}{b-a}, \\ f'(y) &= \frac{f(x)-f(a)}{x-a}. \end{aligned} \quad (1.263)$$

Then, for $x > a$, the inequality reduces to

$$f'(y) \leq f'(c). \quad (1.264)$$

1.37

Let \mathbf{x} and \mathbf{y} be two variables. Then, by the definition,

$$H(\mathbf{x}, \mathbf{y}) = - \int \int p(\mathbf{x}, \mathbf{y}) \ln p(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y}. \quad (1.265)$$

The right hand side can be written as

$$\begin{aligned}
& - \int \int p(\mathbf{x}, \mathbf{y}) (\ln p(\mathbf{y}|\mathbf{x}) + \ln p(\mathbf{x})) d\mathbf{x}d\mathbf{y} \\
& = - \int \int p(\mathbf{x}, \mathbf{y}) \ln p(\mathbf{y}|\mathbf{x}) d\mathbf{x}d\mathbf{y} - \int \left(\int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} \right) \ln p(\mathbf{x}) d\mathbf{x}.
\end{aligned} \tag{1.266}$$

By the definition, the first term of the right hand side can be written as $H(\mathbf{y}|\mathbf{x})$ and the second term can be written as $H(\mathbf{x})$. Therefore,

$$H(\mathbf{x}, \mathbf{y}) = H(\mathbf{y}|\mathbf{x}) + H(\mathbf{x}). \tag{1.267}$$

1.38

Let f be a strictly convex function. Then, by the definition,

$$f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2), \tag{1.268}$$

where $0 \leq \lambda \leq 1$. Let us assume that

$$f\left(\sum_{i=1}^M \lambda_i x_i\right) \leq \sum_{i=1}^M \lambda_i f(x_i), \tag{1.269}$$

where $\lambda_i \geq 0$ and

$$\sum_{i=1}^M \lambda_i = 1. \tag{1.270}$$

Here, let $\lambda_i \geq 0$ and

$$\sum_{i=1}^{M+1} \lambda_i = 1. \tag{1.271}$$

Then, by the definition,

$$f\left(\sum_{i=1}^{M+1} \lambda_i x_i\right) \leq \lambda_{M+1} f(x_{M+1}) + (1 - \lambda_{M+1}) f\left(\sum_{i=1}^M \frac{\lambda_i}{1 - \lambda_{M+1}} x_i\right). \tag{1.272}$$

By the assumption,

$$f\left(\sum_{i=1}^M \frac{\lambda_i}{1 - \lambda_{M+1}} x_i\right) \leq \sum_{i=1}^M \frac{\lambda_i}{1 - \lambda_{M+1}} f(x_i). \tag{1.273}$$

Therefore,

$$f\left(\sum_{i=1}^{M+1} \lambda_i x_i\right) \leq \lambda_{M+1} f(x_{M+1}) + (1 - \lambda_{M+1}) \sum_{i=1}^M \frac{\lambda_i}{1 - \lambda_{M+1}} f(x_i). \quad (1.274)$$

Thus,

$$f\left(\sum_{i=1}^{M+1} \lambda_i x_i\right) \leq \sum_{i=1}^{M+1} \lambda_i f(x_i). \quad (1.275)$$

Hence, the assumption is proved by induction on M .

1.39

Let x and y be two binary variables where

$$\begin{aligned} p(x=0, y=0) &= \frac{1}{3}, \\ p(x=0, y=1) &= \frac{1}{3}, \\ p(x=1, y=0) &= 0, \\ p(x=1, y=1) &= \frac{1}{3}. \end{aligned} \quad (1.276)$$

(a)

By the definition,

$$H(x) = - \sum p(x) \ln p(x). \quad (1.277)$$

By the distribution,

$$\begin{aligned} p(x=0) &= \frac{2}{3}, \\ p(x=1) &= \frac{1}{3}. \end{aligned} \quad (1.278)$$

Therefore,

$$H(x) = \ln 3 - \frac{2}{3} \ln 2. \quad (1.279)$$

(b)

By the definition,

$$H(y) = - \sum p(y) \ln p(y). \quad (1.280)$$

By the distribution,

$$\begin{aligned} p(y = 0) &= \frac{1}{3}, \\ p(y = 1) &= \frac{2}{3}. \end{aligned} \quad (1.281)$$

Therefore,

$$H(y) = \ln 3 - \frac{2}{3} \ln 2. \quad (1.282)$$

(c)

By the definition,

$$H(y|x) = - \sum p(x, y) \ln p(y|x). \quad (1.283)$$

By the definition,

$$\begin{aligned} p(y = 0|x = 0) &= \frac{p(x = 0, y = 0)}{p(x = 0)}, \\ p(y = 0|x = 1) &= \frac{p(x = 1, y = 0)}{p(x = 1)}, \\ p(y = 1|x = 0) &= \frac{p(x = 0, y = 1)}{p(x = 0)}, \\ p(y = 1|x = 1) &= \frac{p(x = 1, y = 1)}{p(x = 1)}. \end{aligned} \quad (1.284)$$

Then, by the distribution,

$$\begin{aligned} p(y = 0|x = 0) &= \frac{1}{2}, \\ p(y = 0|x = 1) &= 0, \\ p(y = 1|x = 0) &= \frac{1}{2}, \\ p(y = 1|x = 1) &= 1. \end{aligned} \quad (1.285)$$

Therefore,

$$H(y|x) = \frac{2}{3} \ln 2. \quad (1.286)$$

(d)

By the definition,

$$H(x|y) = - \sum p(x, y) \ln p(x|y). \quad (1.287)$$

By the definition,

$$\begin{aligned} p(x = 0|y = 0) &= \frac{p(x = 0, y = 0)}{p(y = 0)}, \\ p(x = 0|y = 1) &= \frac{p(x = 0, y = 1)}{p(y = 1)}, \\ p(x = 1|y = 0) &= \frac{p(x = 1, y = 0)}{p(y = 0)}, \\ p(x = 1|y = 1) &= \frac{p(x = 1, y = 1)}{p(y = 1)}. \end{aligned} \quad (1.288)$$

Then, by the distribution,

$$\begin{aligned} p(x = 0|y = 0) &= 1, \\ p(x = 0|y = 1) &= \frac{1}{2}, \\ p(x = 1|y = 0) &= 0, \\ p(x = 1|y = 1) &= \frac{1}{2}. \end{aligned} \quad (1.289)$$

Therefore,

$$H(x|y) = \frac{2}{3} \ln 2. \quad (1.290)$$

(e)

By the definition,

$$H(x, y) = - \sum p(x, y) \ln p(x, y). \quad (1.291)$$

Therefore,

$$H(x, y) = \ln 3. \quad (1.292)$$

(f)

By the definition,

$$I(x, y) = - \sum p(x, y) \ln \frac{p(x)p(y)}{p(x, y)}. \quad (1.293)$$

By the distribution, the right hand side can be written as

$$H(x) + H(y) - H(x, y). \quad (1.294)$$

Therefore,

$$I(x, y) = \ln 3 - \frac{4}{3} \ln 2. \quad (1.295)$$

1.40

Let $\{x_i\}$ be a set of points where $x_i > 0$, and let $\{\lambda_i\}$ be a set of coefficients where $\lambda_i \geq 0$ and

$$\sum_{i=1}^M \lambda_i = 1. \quad (1.296)$$

By Jensen's inequality,

$$\sum_{i=1}^M \lambda_i \ln x_i \leq \ln \left(\sum_{i=1}^M \lambda_i x_i \right). \quad (1.297)$$

Therefore,

$$\prod_{i=1}^M x_i^{\lambda_i} \leq \sum_{i=1}^M \lambda_i x_i. \quad (1.298)$$

Substituting

$$\lambda_i = \frac{1}{M} \quad (1.299)$$

gives

$$\left(\prod_{i=1}^M x_i \right)^{\frac{1}{M}} \leq \frac{1}{M} \sum_{i=1}^M x_i. \quad (1.300)$$

1.41

Let \mathbf{x} and \mathbf{y} be continuous variables. Then, by the definition,

$$I(\mathbf{x}, \mathbf{y}) = - \int \int p(\mathbf{x}, \mathbf{y}) \ln \frac{p(\mathbf{x})p(\mathbf{y})}{p(\mathbf{x}, \mathbf{y})} d\mathbf{x}d\mathbf{y}. \quad (1.301)$$

The right hand side can be written as

$$\begin{aligned} & - \int \int p(\mathbf{x}, \mathbf{y}) \left(\ln p(\mathbf{x}) + \ln \frac{p(\mathbf{y})}{p(\mathbf{x}, \mathbf{y})} \right) d\mathbf{x}d\mathbf{y} \\ & = - \int \left(\int p(\mathbf{x}, \mathbf{y}) d\mathbf{y} \right) \ln p(\mathbf{x}) d\mathbf{x} + \int \int p(\mathbf{x}, \mathbf{y}) \ln p(\mathbf{x}|\mathbf{y}) d\mathbf{x}d\mathbf{y}. \end{aligned} \quad (1.302)$$

By the definition, the first term of the right hand side can be written as $H(\mathbf{x})$ and the second term can be written as $-H(\mathbf{x}|\mathbf{y})$. Therefore,

$$I(\mathbf{x}, \mathbf{y}) = H(\mathbf{x}) - H(\mathbf{x}|\mathbf{y}). \quad (1.303)$$

By the definition,

$$I(\mathbf{x}, \mathbf{y}) = I(\mathbf{y}, \mathbf{x}). \quad (1.304)$$

Thus,

$$I(\mathbf{x}, \mathbf{y}) = H(\mathbf{y}) - H(\mathbf{y}|\mathbf{x}). \quad (1.305)$$

2 Probability Distributions

2.1

Let x be a variable such that

$$p(x|\mu) = \mu^x(1 - \mu)^{1-x}, \quad (2.1)$$

where $x \in \{0, 1\}$. Then,

$$\sum_x p(x|\mu) = 1. \quad (2.2)$$

By the definition,

$$\begin{aligned} \mathbb{E} x &= \mu, \\ \mathbb{E} x^2 &= \mu, \end{aligned} \quad (2.3)$$

Since

$$\text{var } x = \mathbb{E} x^2 - (\mathbb{E} x)^2, \quad (2.4)$$

we have

$$\text{var } x = \mu(1 - \mu). \quad (2.5)$$

By the definition,

$$\mathbb{H}(x) = - \sum_x p(x|\mu) \ln p(x|\mu). \quad (2.6)$$

Therefore,

$$\mathbb{H}(x) = -\mu \ln \mu - (1 - \mu) \ln(1 - \mu). \quad (2.7)$$

2.2

Let x be a variable such that

$$p(x|\mu) = \left(\frac{1 - \mu}{2} \right)^{\frac{1-x}{2}} \left(\frac{1 + \mu}{2} \right)^{\frac{1+x}{2}}, \quad (2.8)$$

where $x \in \{-1, 1\}$. Then,

$$\sum_x p(x|\mu) = 1. \quad (2.9)$$

By the definition,

$$\begin{aligned} \mathbb{E} x &= \mu, \\ \mathbb{E} x^2 &= 1, \end{aligned} \quad (2.10)$$

Since

$$\text{var } x = \mathbb{E} x^2 - (\mathbb{E} x)^2, \quad (2.11)$$

we have

$$\text{var } x = 1 - \mu^2. \quad (2.12)$$

By the definition,

$$H(x) = - \sum_x p(x|\mu) \ln p(x|\mu). \quad (2.13)$$

Therefore,

$$H(x) = -\frac{1-\mu}{2} \ln \frac{1-\mu}{2} - \frac{1+\mu}{2} \ln \frac{1+\mu}{2}. \quad (2.14)$$

2.3

By the definition,

$$\begin{aligned} \binom{N}{m} &= \frac{N!}{m!(N-m)!}, \\ \binom{N}{m-1} &= \frac{N!}{(m-1)!(N-m+1)!} \end{aligned} \quad (2.15)$$

Therefore,

$$\binom{N}{m} + \binom{N}{m-1} = \frac{(N-m+1)N! + mN!}{m!(N-m+1)!}. \quad (2.16)$$

By the definition, the right hand side can be written as

$$\frac{(N+1)!}{m!(N+1-m)!} = \binom{N+1}{m}. \quad (2.17)$$

Thus,

$$\binom{N}{m} + \binom{N}{m-1} = \binom{N+1}{m}. \quad (2.18)$$

Note that

$$1+x = \sum_{m=0}^1 \binom{1}{m} x^m. \quad (2.19)$$

Let us assume that

$$(1+x)^N = \sum_{m=0}^N \binom{N}{m} x^m. \quad (2.20)$$

Then,

$$(1+x)^{N+1} = \sum_{m=0}^N \binom{N}{m} x^m + \sum_{m=0}^N \binom{N}{m} x^{m+1}. \quad (2.21)$$

By the result above, the right hand side can be written as

$$\sum_{m=0}^N \binom{N}{m} x^m + \sum_{m=1}^{N+1} \binom{N}{m-1} x^m = 1 + x^{N+1} + \sum_{m=1}^N \binom{N+1}{m} x^m. \quad (2.22)$$

Therefore,

$$(1+x)^{N+1} = \sum_{m=0}^{N+1} \binom{N+1}{m} x^m. \quad (2.23)$$

Thus, the assumption is proved by induction on N .

Finally, let m be a variable such that

$$p(m|\mu) = \binom{N}{m} \mu^m (1-\mu)^{N-m}. \quad (2.24)$$

Then

$$\sum_{m=0}^N p(m|\mu) = \sum_{m=0}^N \binom{N}{m} \mu^m (1-\mu)^{N-m}. \quad (2.25)$$

By the result above, the right hand side can be written as

$$(1-\mu)^N \sum_{m=0}^N \binom{N}{m} \left(\frac{\mu}{1-\mu} \right)^m = (1-\mu)^N \left(1 + \frac{\mu}{1-\mu} \right)^N. \quad (2.26)$$

Therefore,

$$\sum_{m=0}^N p(m|\mu) = 1. \quad (2.27)$$

2.4

Let m be a variable such that

$$p(m|\mu) = \binom{N}{m} \mu^m (1-\mu)^{N-m}. \quad (2.28)$$

Then

$$E m = \sum_{m=0}^N m \binom{N}{m} \mu^m (1-\mu)^{N-m}. \quad (2.29)$$

Taking the derivative of

$$\sum_{m=0}^N \binom{N}{m} \mu^m (1-\mu)^{N-m} = 1 \quad (2.30)$$

with respect to μ gives

$$\sum_{m=0}^N m \binom{N}{m} \mu^{m-1} (1-\mu)^{N-m} - \sum_{m=0}^N (N-m) \binom{N}{m} \mu^m (1-\mu)^{N-m-1} = 0. \quad (2.31)$$

The first term of the left hand side can be written as $\frac{1}{\mu} E m$. Since

$$(N-m) \binom{N}{m} = N \binom{N-1}{m}, \quad (2.32)$$

the second term of the left hand side can be written as

$$-N \sum_{m=0}^{N-1} \binom{N-1}{m} \mu^m (1-\mu)^{N-m-1} = -N. \quad (2.33)$$

Therefore,

$$E m = N\mu. \quad (2.34)$$

Taking the second derivative of

$$\sum_{m=0}^N \binom{N}{m} \mu^m (1-\mu)^{N-m} = 1 \quad (2.35)$$

with respect to μ gives

$$\begin{aligned} & \sum_{m=0}^N m(m-1) \binom{N}{m} \mu^{m-2} (1-\mu)^{N-m} \\ & - 2 \sum_{m=0}^N m(N-m) \binom{N}{m} \mu^{m-1} (1-\mu)^{N-m-1} \\ & + \sum_{m=0}^N (N-m)(N-m-1) \binom{N}{m} \mu^m (1-\mu)^{N-m-2} = 0. \end{aligned} \quad (2.36)$$

The first term of the left hand side can be written as $\frac{1}{\mu^2} E m(m-1)$. Since

$$\begin{aligned} m(N-m) \binom{N}{m} &= N(N-1) \binom{N-2}{m-1}, \\ (N-m)(N-m-1) \binom{N}{m} &= N(N-1) \binom{N-2}{m}, \end{aligned} \quad (2.37)$$

the second and third term of the left hand side can be written as

$$\begin{aligned} -2N(N-1) \sum_{m=1}^{N-1} \binom{N-2}{m-1} \mu^{m-1} (1-\mu)^{N-m-1} &= -2N(N-1), \\ N(N-1) \sum_{m=0}^N \binom{N-2}{m} \mu^m (1-\mu)^{N-m-2} &= N(N-1). \end{aligned} \quad (2.38)$$

Therefore,

$$E m(m-1) = N(N-1)\mu^2. \quad (2.39)$$

Thus, since

$$\text{var } m = E m(m-1) + E m - (E m)^2, \quad (2.40)$$

we have

$$\text{var } m = N\mu(1-\mu). \quad (2.41)$$

2.5

By the definition,

$$\Gamma(a)\Gamma(b) = \int_0^\infty x^{a-1} \exp(-x) dx \int_0^\infty y^{b-1} \exp(-y) dy. \quad (2.42)$$

By the transformation $t = x + y$, the right hand side can be written as

$$\begin{aligned} &\int_0^\infty x^{a-1} \left(\int_x^\infty (t-x)^{b-1} \exp(-t) dt \right) dx \\ &= \int_0^\infty \left(\int_0^t x^{a-1} (t-x)^{b-1} dx \right) \exp(-t) dt. \end{aligned} \quad (2.43)$$

By the transformation $x = t\mu$, the right hand side can be written as

$$\begin{aligned} &\int_0^\infty \left(\int_0^1 (t\mu)^{a-1} t^{b-1} (1-\mu)^{b-1} t d\mu \right) \exp(-t) dt \\ &= \int_0^1 \mu^{a-1} (1-\mu)^{b-1} d\mu \int_0^\infty t^{a+b-1} \exp(-t) dt. \end{aligned} \quad (2.44)$$

By the definition, the second integral of the right hand side can be written as $\Gamma(a+b)$. Therefore,

$$\int_0^1 \mu^{a-1}(1-\mu)^{b-1}d\mu = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}. \quad (2.45)$$

2.6

Let μ be a variable such that

$$p(\mu|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \mu^{a-1}(1-\mu)^{b-1}. \quad (2.46)$$

Then

$$\begin{aligned} E\mu &= \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \mu^a(1-\mu)^{b-1}d\mu, \\ E\mu^2 &= \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \int_0^1 \mu^{a+1}(1-\mu)^{b-1}d\mu. \end{aligned} \quad (2.47)$$

Since

$$\begin{aligned} \int_0^1 \mu^a(1-\mu)^{b-1}d\mu &= \frac{\Gamma(a+1)\Gamma(b)}{\Gamma(a+b+1)}, \\ \int_0^1 \mu^{a+1}(1-\mu)^{b-1}d\mu &= \frac{\Gamma(a+2)\Gamma(b)}{\Gamma(a+b+2)}, \end{aligned} \quad (2.48)$$

we have

$$\begin{aligned} E\mu &= \frac{a}{a+b}, \\ E\mu^2 &= \frac{a(a+1)}{(a+b)(a+b+1)}. \end{aligned} \quad (2.49)$$

Since

$$\text{var } \mu = E\mu^2 - (E\mu)^2, \quad (2.50)$$

we have

$$\text{var } \mu = \frac{ab}{(a+b)^2(a+b+1)}. \quad (2.51)$$

Since

$$\frac{\partial}{\partial \mu} p(\mu|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \mu^{a-1}(1-\mu)^{b-1} \left(\frac{a-1}{\mu} - \frac{b-1}{1-\mu} \right), \quad (2.52)$$

we have

$$\text{mode } \mu = \frac{a-1}{a+b-2}. \quad (2.53)$$

2.7

Let m and l be a variable such that

$$p(m, l|\mu) = \binom{m+l}{m} \mu^m (1-\mu)^l, \quad (2.54)$$

where

$$p(\mu|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \mu^{a-1} (1-\mu)^{b-1}. \quad (2.55)$$

By 2.6,

$$E(\mu|a, b) = \frac{a}{a+b}. \quad (2.56)$$

Note that

$$\mu_{\text{ML}} = \frac{m}{m+l}. \quad (2.57)$$

Since

$$p(\mu|m, l, a, b) \propto p(m, l|\mu)p(\mu|a, b), \quad (2.58)$$

we have

$$p(\mu|m, l, a, b) = \frac{\Gamma(m+l+a+b)}{\Gamma(m+a)\Gamma(l+b)} \mu^{m+a-1} (1-\mu)^{l+b-1}. \quad (2.59)$$

Therefore, by 2.6,

$$E(\mu|m, l, a, b) = \frac{m+a}{m+l+a+b}. \quad (2.60)$$

Thus,

$$E(\mu|m, l, a, b) = \lambda \mu_{\text{ML}} + (1-\lambda) E(\mu|a, b), \quad (2.61)$$

where

$$\lambda = \frac{m+l}{m+l+a+b}. \quad (2.62)$$

2.8

Let x and y be variables. Then, by the definition,

$$E x = \int x p(x) dx. \quad (2.63)$$

The right hand side can be written as

$$\int x \left(\int p(x, y) dy \right) dx = \int \left(\int x p(x|y) dx \right) p(y) dy. \quad (2.64)$$

Therefore,

$$\mathbb{E} x = \mathbb{E}_y (\mathbb{E}_x(x|y)) . \quad (2.65)$$

Additionally, by the definition,

$$\text{var } x = \mathbb{E} (x - \mathbb{E} x)^2 . \quad (2.66)$$

By the result above, the right hand side can be written as

$$\begin{aligned} & \mathbb{E}_y (\mathbb{E}_x ((x - \mathbb{E}_x(x|y) + \mathbb{E}_x(x|y) - \mathbb{E} x)^2 | y)) \\ &= \mathbb{E}_y (\mathbb{E}_x ((x - \mathbb{E}_x(x|y))^2 | y)) \\ & \quad + 2 \mathbb{E}_y (((\mathbb{E}_x(x|y) - \mathbb{E} x) \mathbb{E}_x(x - \mathbb{E}_x(x|y)) | y)) \\ & \quad + \mathbb{E}_y ((\mathbb{E}_x(x|y) - \mathbb{E} x)^2 | y) . \end{aligned} \quad (2.67)$$

Let us look at each term of the right hand side. By the definition, the first term can be written as $\mathbb{E}_y (\text{var}_x(x|y))$. The second term can be written as

$$2 \mathbb{E}_y ((\mathbb{E}_x(x|y) - \mathbb{E} x) (\mathbb{E}_x(x|y) - \mathbb{E}_x(x|y))) = 0. \quad (2.68)$$

By the result above, the third term can be written as

$$\mathbb{E}_y (\mathbb{E}_x(x|y) - \mathbb{E}_y(\mathbb{E}_x(x|y)))^2 = \text{var}_y (\mathbb{E}_x(x|y)) . \quad (2.69)$$

Therefore,

$$\text{var } x = \mathbb{E}_y (\text{var}_x(x|y)) + \text{var}_y (\mathbb{E}_x(x|y)) . \quad (2.70)$$

2.9

For a vector $\boldsymbol{\mu}$ in 2 dimensions, 2.5 gives

$$\int_{\substack{\mu_1 + \mu_2 = 1 \\ \mu_1 \geq 0, \mu_2 \geq 0}} \mu_1^{\alpha_1 - 1} \mu_2^{\alpha_2 - 1} d\boldsymbol{\mu} = \frac{\Gamma(\alpha_1) \Gamma(\alpha_2)}{\Gamma(\alpha_1 + \alpha_2)} .$$

For a vector $\boldsymbol{\mu}$ in M dimensions, let us assume that

$$\int_{\substack{\sum_{k=1}^M \mu_k = 1 \\ \mu_k \geq 0}} \prod_{k=1}^M \mu_k^{\alpha_k - 1} d\boldsymbol{\mu} = \frac{\prod_{k=1}^M \Gamma(\alpha_k)}{\Gamma(\sum_{k=1}^M \alpha_k)} .$$

Under the constraint

$$\sum_{k=1}^{M+1} \mu_k = 1, \quad (2.71)$$

we have

$$\begin{aligned}
& \int_0^{1-\sum_{k=1}^{M-1} \mu_k} \prod_{k=1}^{M+1} \mu_k^{\alpha_k-1} d\mu_{M+1} \\
&= \left(\prod_{k=1}^{M-1} \mu_k^{\alpha_k-1} \right) \int_0^{1-\sum_{k=1}^{M-1} \mu_k} \mu_{M+1}^{\alpha_{M+1}-1} \left(1 - \sum_{k=1}^{M-1} \mu_k - \mu_{M+1} \right)^{\alpha_M-1} d\mu_{M+1}.
\end{aligned} \tag{2.72}$$

By the transformation

$$\mu'_{M+1} = \frac{\mu_{M+1}}{1 - \sum_{k=1}^{M-1} \mu_k}, \tag{2.73}$$

the integral of the right hand side can be written as

$$\begin{aligned}
& \int_0^1 \left(\left(1 - \sum_{k=1}^{M-1} \mu_k \right) \mu'_{M+1} \right)^{\alpha_{M+1}-1} \left(\left(1 - \sum_{k=1}^{M-1} \mu_k \right) (1 - \mu'_{M+1}) \right)^{\alpha_M-1} \left(1 - \sum_{k=1}^{M-1} \mu_k \right) d\mu'_{M+1} \\
&= \left(1 - \sum_{k=1}^{M-1} \mu_k \right)^{\alpha_M + \alpha_{M+1} - 1} \int_0^1 \mu'_{M+1}^{\alpha_{M+1}-1} (1 - \mu'_{M+1})^{\alpha_M-1} d\mu'_{M+1}.
\end{aligned} \tag{2.74}$$

By 2.5, the integral of the right hand side can be written as

$$\frac{\Gamma(\alpha_M)\Gamma(\alpha_{M+1})}{\Gamma(\alpha_M + \alpha_{M+1})}. \tag{2.75}$$

Therefore,

$$\int_0^{1-\sum_{k=1}^{M-1} \mu_k} \prod_{k=1}^{M+1} \mu_k^{\alpha_k-1} d\mu_{M+1} = \left(\prod_{k=1}^{M-1} \mu_k^{\alpha_k-1} \right) \left(1 - \sum_{k=1}^{M-1} \mu_k \right)^{\alpha_M + \alpha_{M+1} - 1} \frac{\Gamma(\alpha_M)\Gamma(\alpha_{M+1})}{\Gamma(\alpha_M + \alpha_{M+1})}. \tag{2.76}$$

By the assumption,

$$\int_{\substack{\sum_{k=1}^M \mu_k = 1 \\ \mu_k \geq 0}} \left(\prod_{k=1}^{M-1} \mu_k^{\alpha_k-1} \right) \mu_M^{\alpha_M + \alpha_{M+1} - 1} d\boldsymbol{\mu} = \frac{(\prod_{k=1}^{M-1} \Gamma(\alpha_k))\Gamma(\alpha_M + \alpha_{M+1})}{\Gamma(\sum_{k=1}^{M+1} \alpha_k)}.$$

Thus, for a vector $\boldsymbol{\mu}$ in $M+1$ dimensions,

$$\int_{\substack{\sum_{k=1}^{M+1} \mu_k = 1 \\ \mu_k \geq 0}} \prod_{k=1}^{M+1} \mu_k^{\alpha_k-1} d\boldsymbol{\mu} = \frac{\Gamma(\alpha_M)\Gamma(\alpha_{M+1})}{\Gamma(\alpha_M + \alpha_{M+1})} \frac{(\prod_{k=1}^{M-1} \Gamma(\alpha_k))\Gamma(\alpha_M + \alpha_{M+1})}{\Gamma(\sum_{k=1}^{M+1} \alpha_k)}.$$

The right hand side can be written as

$$\frac{\prod_{k=1}^{M+1} \Gamma(\alpha_k)}{\Gamma(\sum_{k=1}^{M+1} \alpha_k)}. \quad (2.77)$$

Hence, the assumption is proved by induction on M .

2.10

Let $\boldsymbol{\mu}$ be a variable such that

$$p(\boldsymbol{\mu}|\boldsymbol{\alpha}) = \frac{\Gamma(\sum_{k=1}^K \alpha_k)}{\prod_{k=1}^K \Gamma(\alpha_k)} \prod_{k=1}^K \mu_k^{\alpha_k-1}. \quad (2.78)$$

Then

$$\begin{aligned} \mathbb{E} \mu_j &= \int \mu_j p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu}, \\ \mathbb{E} \mu_j^2 &= \int \mu_j^2 p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu}, \\ \mathbb{E} \mu_j \mu_l &= \int \mu_j \mu_l p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu}. \end{aligned} \quad (2.79)$$

If $j \neq l$, then the right hand sides can be written as

$$\begin{aligned} \frac{\Gamma(\sum_{k=1}^K \alpha_k)}{\prod_{k=1}^K \Gamma(\alpha_k)} \frac{\frac{\Gamma(\alpha_j+1)}{\Gamma(\alpha_j)} \prod_{k=1}^K \Gamma(\alpha_k)}{\Gamma(\sum_{k=1}^K \alpha_k + 1)} &= \frac{\alpha_j}{\sum_{k=1}^K \alpha_k}, \\ \frac{\Gamma(\sum_{k=1}^K \alpha_k)}{\prod_{k=1}^K \Gamma(\alpha_k)} \frac{\frac{\Gamma(\alpha_j+2)}{\Gamma(\alpha_j)} \prod_{k=1}^K \Gamma(\alpha_k)}{\Gamma(\sum_{k=1}^K \alpha_k + 2)} &= \frac{\alpha_j(\alpha_j + 1)}{\sum_{k=1}^K \alpha_k (\sum_{k=1}^K \alpha_k + 1)}, \\ \frac{\Gamma(\sum_{k=1}^K \alpha_k)}{\prod_{k=1}^K \Gamma(\alpha_k)} \frac{\frac{\Gamma(\alpha_j+1)\Gamma(\alpha_l+1)}{\Gamma(\alpha_j)\Gamma(\alpha_l)} \prod_{k=1}^K \Gamma(\alpha_k)}{\Gamma(\sum_{k=1}^K \alpha_k + 2)} &= \frac{\alpha_j \alpha_l}{\sum_{k=1}^K \alpha_k (\sum_{k=1}^K \alpha_k + 1)}. \end{aligned} \quad (2.80)$$

Therefore,

$$\begin{aligned} \mathbb{E} \mu_j &= \frac{\alpha_j}{\sum_{k=1}^K \alpha_k}, \\ \mathbb{E} \mu_j^2 &= \frac{\alpha_j(\alpha_j + 1)}{\sum_{k=1}^K \alpha_k (\sum_{k=1}^K \alpha_k + 1)}, \\ \mathbb{E} \mu_j \mu_l &= \frac{\alpha_j \alpha_l}{\sum_{k=1}^K \alpha_k (\sum_{k=1}^K \alpha_k + 1)}. \end{aligned} \quad (2.81)$$

Since

$$\begin{aligned}\text{var } \mu_j &= \text{E } \mu_j^2 - (\text{E } \mu_j)^2, \\ \text{cov } (\mu_j, \mu_l) &= \text{E } \mu_j \mu_l - \text{E } \mu_j \text{E } \mu_l,\end{aligned}\tag{2.82}$$

we have

$$\begin{aligned}\text{var } \mu_j &= \frac{\alpha_j (\sum_{k=1}^K \alpha_k - \alpha_j)}{\sum_{k=1}^K \alpha_k^2 (\sum_{k=1}^K \alpha_k + 1)}, \\ \text{cov } (\mu_j, \mu_l) &= -\frac{\alpha_j \alpha_l}{(\sum_{k=1}^K \alpha_k)^2 (\sum_{k=1}^K \alpha_k + 1)}.\end{aligned}\tag{2.83}$$

2.11

Let $\boldsymbol{\mu}$ be a variable such that

$$p(\boldsymbol{\mu}|\boldsymbol{\alpha}) = \frac{\Gamma\left(\sum_{k=1}^K \alpha_k\right)}{\prod_{k=1}^K \Gamma(\alpha_k)} \prod_{k=1}^K \mu_k^{\alpha_k-1}.\tag{2.84}$$

Then

$$\text{E } \ln \mu_j = \int (\ln \mu_j) p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu}.\tag{2.85}$$

Since

$$\frac{\partial}{\partial \alpha_j} p(\boldsymbol{\mu}|\boldsymbol{\alpha}) = \left(\frac{\Gamma'\left(\sum_{k=1}^K \alpha_k\right)}{\Gamma\left(\sum_{k=1}^K \alpha_k\right)} - \frac{\Gamma'(\alpha_j)}{\Gamma(\alpha_j)} + \ln \mu_j \right) p(\boldsymbol{\mu}|\boldsymbol{\alpha}),\tag{2.86}$$

we have

$$\text{E } \ln \mu_j = \frac{\partial}{\partial \alpha_j} \int p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu} + \left(\psi(\alpha_j) - \psi\left(\sum_{k=1}^K \alpha_k\right) \right) \int p(\boldsymbol{\mu}|\boldsymbol{\alpha}) d\boldsymbol{\mu},\tag{2.87}$$

where

$$\psi(a) = \frac{d}{da} \ln \Gamma(a).\tag{2.88}$$

Therefore,

$$\text{E } \ln \mu_j = \psi(\alpha_j) - \psi\left(\sum_{k=1}^K \alpha_k\right).\tag{2.89}$$

2.12

Let x be a variable such that

$$p(x|a, b) = \frac{1}{b-a}, \quad (2.90)$$

where $a < b$. Then

$$\int_a^b p(x|a, b) dx = 1. \quad (2.91)$$

Note that

$$\begin{aligned} \mathbb{E} x &= \int_a^b x p(x|a, b) dx, \\ \mathbb{E} x^2 &= \int_a^b x^2 p(x|a, b) dx. \end{aligned} \quad (2.92)$$

The right hand sides can be written as

$$\begin{aligned} \frac{1}{b-a} \int_a^b x dx &= \frac{1}{2}(a+b), \\ \frac{1}{b-a} \int_a^b x^2 dx &= \frac{1}{3}(a^2 + ab + b^2). \end{aligned} \quad (2.93)$$

Therefore,

$$\begin{aligned} \mathbb{E} x &= \frac{1}{2}(a+b), \\ \mathbb{E} x^2 &= \frac{1}{3}(a^2 + ab + b^2). \end{aligned} \quad (2.94)$$

Since

$$\text{var } x = \mathbb{E} x^2 - (\mathbb{E} x)^2, \quad (2.95)$$

we have

$$\text{var } x = \frac{1}{12}(b-a)^2. \quad (2.96)$$

2.13

Let \mathbf{x} be a variable in D dimensions and

$$\begin{aligned} p(\mathbf{x}) &= \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}), \\ q(\mathbf{x}) &= \mathcal{N}(\mathbf{x}|\mathbf{m}, \mathbf{L}). \end{aligned} \quad (2.97)$$

Then, by the definition,

$$\text{KL}(p||q) = - \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) \ln \frac{\mathcal{N}(\mathbf{x}|\mathbf{m}, \mathbf{L})}{\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma})} d\mathbf{x}. \quad (2.98)$$

Since

$$\ln \frac{\mathcal{N}(\mathbf{x}|\mathbf{m}, \mathbf{L})}{\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma})} = \ln \frac{(2\pi)^{-\frac{D}{2}} (|\det \mathbf{L}|)^{-\frac{1}{2}} \exp \left(-\frac{1}{2}(\mathbf{x} - \mathbf{m})^\top \mathbf{L}^{-1}(\mathbf{x} - \mathbf{m}) \right)}{(2\pi)^{-\frac{D}{2}} (|\det \boldsymbol{\Sigma}|)^{-\frac{1}{2}} \exp \left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \right)}, \quad (2.99)$$

The right hand side can be written as

$$\begin{aligned} & \frac{1}{2} \ln \left| \frac{\det \mathbf{L}}{\det \boldsymbol{\Sigma}} \right| \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ & + \frac{1}{2} \int (\mathbf{x} - \mathbf{m})^\top \mathbf{L}^{-1}(\mathbf{x} - \mathbf{m}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ & - \frac{1}{2} \int (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x}. \end{aligned} \quad (2.100)$$

Let us look at each term. Since

$$\int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} = 1, \quad (2.101)$$

the first term can be written as $\frac{1}{2} \ln \left| \frac{\det \mathbf{L}}{\det \boldsymbol{\Sigma}} \right|$. Since

$$(\mathbf{x} - \mathbf{m})^\top \mathbf{L}^{-1}(\mathbf{x} - \mathbf{m}) = (\mathbf{x} - \boldsymbol{\mu} + \boldsymbol{\mu} - \mathbf{m})^\top \mathbf{L}^{-1}(\mathbf{x} - \boldsymbol{\mu} + \boldsymbol{\mu} - \mathbf{m}), \quad (2.102)$$

the second term can be written as

$$\begin{aligned} & \frac{1}{2} \int (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{L}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ & + (\boldsymbol{\mu} - \mathbf{m})^\top \mathbf{L}^{-1} \int (\mathbf{x} - \boldsymbol{\mu}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ & + \frac{1}{2} (\boldsymbol{\mu} - \mathbf{m})^\top \mathbf{L}^{-1}(\boldsymbol{\mu} - \mathbf{m}) \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x}. \end{aligned} \quad (2.103)$$

Since

$$\begin{aligned} & \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} = 1, \\ & \int \mathbf{x} \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} = \boldsymbol{\mu}, \\ & \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} = \boldsymbol{\Sigma}, \end{aligned} \quad (2.104)$$

it can be written as

$$\frac{1}{2} \text{tr} (\mathbf{L}^{-1} \mathbf{\Sigma}) + \frac{1}{2} (\boldsymbol{\mu} - \mathbf{m})^\top \mathbf{L}^{-1} (\boldsymbol{\mu} - \mathbf{m}). \quad (2.105)$$

Since

$$\int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x} = \mathbf{\Sigma}, \quad (2.106)$$

the third term can be written as

$$-\frac{1}{2} \text{tr} (\mathbf{\Sigma}^{-1} \mathbf{\Sigma}) = -\frac{D}{2}. \quad (2.107)$$

Therefore,

$$\text{KL}(p||q) = \frac{1}{2} \left(\ln \left| \frac{\det \mathbf{L}}{\det \mathbf{\Sigma}} \right| + \text{tr} (\mathbf{L}^{-1} \mathbf{\Sigma}) + (\boldsymbol{\mu} - \mathbf{m})^\top \mathbf{L}^{-1} (\boldsymbol{\mu} - \mathbf{m}) - D \right). \quad (2.108)$$

2.14

Let \mathbf{x} be a variable in D dimensions and

$$\begin{aligned} L(p(\mathbf{x})) = & - \int p(\mathbf{x}) \ln p(\mathbf{x}) d\mathbf{x} + \lambda \left(\int p(\mathbf{x}) d\mathbf{x} - 1 \right) \\ & + \mathbf{l}^\top \left(\int \mathbf{x} p(\mathbf{x}) d\mathbf{x} - \boldsymbol{\mu} \right) + \mathbf{m}^\top \left(\int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top p(\mathbf{x}) d\mathbf{x} - \mathbf{\Sigma} \right) \mathbf{m}. \end{aligned} \quad (2.109)$$

Then

$$\frac{\delta L(p(\mathbf{x}))}{\delta p(\mathbf{x})} = -\ln p(\mathbf{x}) - 1 + \lambda + \mathbf{l}^\top \mathbf{x} + \mathbf{m}^\top (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{m}. \quad (2.110)$$

Setting the left hand side to zero gives

$$p(\mathbf{x}) = \exp (-1 + \lambda + \mathbf{l}^\top \mathbf{x} + \mathbf{m}^\top (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{m}), \quad (2.111)$$

so that

$$p(\mathbf{x}) = \exp \left(-1 + \lambda - \mathbf{l}^\top \mathbf{M} \mathbf{l} + (\mathbf{x} - \boldsymbol{\mu} - \mathbf{M} \mathbf{l})^\top \mathbf{M}^{-1} (\mathbf{x} - \boldsymbol{\mu} - \mathbf{M} \mathbf{l}) \right), \quad (2.112)$$

where

$$\mathbf{M} = (\mathbf{m} \mathbf{m}^\top)^{-1}. \quad (2.113)$$

Substituting it to

$$\begin{aligned}\int p(\mathbf{x})d\mathbf{x} &= 1, \\ \int \mathbf{x}p(\mathbf{x})d\mathbf{x} &= \boldsymbol{\mu}, \\ \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top p(\mathbf{x})d\mathbf{x} &= \boldsymbol{\Sigma},\end{aligned}\tag{2.114}$$

and the transformation

$$\mathbf{y} = \mathbf{x} - \boldsymbol{\mu} - \mathbf{M}\mathbf{l}\tag{2.115}$$

gives

$$\begin{aligned}\exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) \int \exp(-\mathbf{y}^\top \mathbf{M}^{-1} \mathbf{y}) d\mathbf{y} &= 1, \\ \exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) \int (\mathbf{y} + \boldsymbol{\mu} + \mathbf{M}\mathbf{l}) \exp(-\mathbf{y}^\top \mathbf{M}^{-1} \mathbf{y}) d\mathbf{y} &= \boldsymbol{\mu}, \\ \exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) \int (\mathbf{y} + \mathbf{M}\mathbf{l}) (\mathbf{y} + \mathbf{M}\mathbf{l})^\top \exp(-\mathbf{y}^\top \mathbf{M}^{-1} \mathbf{y}) d\mathbf{y} &= \boldsymbol{\Sigma}.\end{aligned}\tag{2.116}$$

Since

$$\begin{aligned}\int \exp(-\mathbf{y}^\top \mathbf{y}) d\mathbf{y} &= \left(\Gamma\left(\frac{1}{2}\right) \right)^D, \\ \int \mathbf{y} \exp(-\mathbf{y}^\top \mathbf{y}) d\mathbf{y} &= \mathbf{0}, \\ \int \mathbf{y}\mathbf{y}^\top \exp(-\mathbf{y}^\top \mathbf{y}) d\mathbf{y} &= \Gamma\left(\frac{3}{2}\right) \left(\Gamma\left(\frac{1}{2}\right) \right)^{D-1} \mathbf{I},\end{aligned}\tag{2.117}$$

they can be written as

$$\begin{aligned}\exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) \left(\Gamma\left(\frac{1}{2}\right) \right)^D (\det \mathbf{M})^{\frac{1}{2}} &= 1, \\ \exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) (\boldsymbol{\mu} + \mathbf{M}\mathbf{l}) \left(\Gamma\left(\frac{1}{2}\right) \right)^D (\det \mathbf{M})^{\frac{1}{2}} &= \boldsymbol{\mu}, \\ \exp(-1 + \lambda - \mathbf{l}^\top \mathbf{M}\mathbf{l}) \left(\Gamma\left(\frac{3}{2}\right) \left(\Gamma\left(\frac{1}{2}\right) \right)^{D-1} \mathbf{M} + \mathbf{M}\mathbf{l}(\mathbf{M}\mathbf{l})^\top \left(\Gamma\left(\frac{1}{2}\right) \right)^D \right) (\det \mathbf{M})^{\frac{1}{2}} &= \boldsymbol{\Sigma}.\end{aligned}\tag{2.118}$$

Therefore,

$$\begin{aligned}\lambda &= 1 - \frac{D}{2} \ln \pi - \frac{1}{2} \ln(\det \mathbf{M}), \\ \mathbf{l} &= \mathbf{0}, \\ \mathbf{M} &= 2\mathbf{\Sigma}.\end{aligned}\tag{2.119}$$

Thus,

$$p(\mathbf{x}) = (2\pi)^{-\frac{D}{2}} (\det \mathbf{\Sigma})^{-\frac{1}{2}} \exp \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right).\tag{2.120}$$

2.15

Let \mathbf{x} be a variable in D dimensions such that

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}).\tag{2.121}$$

Then, by the definition,

$$\mathbf{H}(\mathbf{x}) = - \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) \ln \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x}.\tag{2.122}$$

The right hand side can be written as

$$\begin{aligned}& - \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) \left(-\frac{D}{2} \ln(2\pi) - \frac{1}{2} \ln |\det \mathbf{\Sigma}| - \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right) d\mathbf{x} \\ &= \left(\frac{D}{2} \ln(2\pi) + \frac{1}{2} \ln |\det \mathbf{\Sigma}| \right) \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x} \\ &+ \frac{1}{2} \int (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x}.\end{aligned}\tag{2.123}$$

Since

$$\begin{aligned}\int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x} &= 1, \\ \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}) d\mathbf{x} &= \mathbf{\Sigma},\end{aligned}\tag{2.124}$$

the first and second term of the right hand side can be written as

$$\frac{D}{2} \ln(2\pi) + \frac{1}{2} \ln |\det \mathbf{\Sigma}|\tag{2.125}$$

and

$$\frac{1}{2} \text{tr} (\mathbf{\Sigma}^{-1} \mathbf{\Sigma}) = \frac{D}{2}. \quad (2.126)$$

Therefore,

$$H(\mathbf{x}) = \frac{D}{2} (1 + \ln(2\pi)) + \frac{1}{2} \ln |\det \mathbf{\Sigma}|. \quad (2.127)$$

2.16

Let x be a variable such that

$$x = x_1 + x_2, \quad (2.128)$$

where

$$\begin{aligned} p(x_1) &= \mathcal{N}(x_1 | \mu_1, \tau_1^{-1}), \\ p(x_2) &= \mathcal{N}(x_2 | \mu_2, \tau_2^{-1}). \end{aligned} \quad (2.129)$$

By marginalisation,

$$p(x) = \int_{-\infty}^{\infty} p(x|x_2)p(x_2)dx_2. \quad (2.130)$$

The right hand side can be written as

$$\begin{aligned} & \int_{-\infty}^{\infty} \mathcal{N}(x | \mu_1 + x_2, \tau_1^{-1}) \mathcal{N}(x_2 | \mu_2, \tau_2^{-1}) dx_2 \\ &= \int_{-\infty}^{\infty} \left(\frac{\tau_1}{2\pi}\right)^{\frac{1}{2}} \exp\left(-\frac{\tau_1}{2}(x - \mu_1 - x_2)^2\right) \left(\frac{\tau_2}{2\pi}\right)^{\frac{1}{2}} \exp\left(-\frac{\tau_2}{2}(x_2 - \mu_2)^2\right) dx_2. \end{aligned} \quad (2.131)$$

The logarithm of the integrand except the terms independent of x and z is given by

$$\begin{aligned} & -\frac{\tau_1 + \tau_2}{2} \left(x_2 - \frac{\tau_1(x - \mu_1) + \tau_2\mu_2}{\tau_1 + \tau_2}\right)^2 - \frac{\tau_1}{2}(x - \mu_1)^2 - \frac{\tau_2}{2}\mu_2^2 \\ & + \frac{\tau_1 + \tau_2}{2} \left(\frac{\tau_1(x - \mu_1) + \tau_2\mu_2}{\tau_1 + \tau_2}\right)^2 \\ & = -\frac{\tau_1 + \tau_2}{2} \left(x_2 - \frac{\tau_1(x - \mu_1) + \tau_2\mu_2}{\tau_1 + \tau_2}\right)^2 - \frac{\tau_1\tau_2}{2(\tau_1 + \tau_2)}(x - \mu_1 - \mu_2)^2. \end{aligned} \quad (2.132)$$

Therefore,

$$p(x) = \mathcal{N}(x | \mu_1 + \mu_2, \tau_1^{-1} + \tau_2^{-1}). \quad (2.133)$$

Thus, by 1.35,

$$H(x) = \frac{1}{2} (1 + \ln(2\pi) + \ln(\tau_1^{-1} + \tau_2^{-1})). \quad (2.134)$$

2.17

Let Σ be a matrix and

$$\begin{aligned} \mathbf{S} &= \frac{1}{2} (\Sigma^{-1} + (\Sigma^{-1})^\top), \\ \mathbf{A} &= \frac{1}{2} (\Sigma^{-1} - (\Sigma^{-1})^\top). \end{aligned} \quad (2.135)$$

Then

$$\Sigma^{-1} = \mathbf{S} + \mathbf{A}. \quad (2.136)$$

Therefore,

$$(\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) = (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{S} (\mathbf{x} - \boldsymbol{\mu}) + (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A} (\mathbf{x} - \boldsymbol{\mu}). \quad (2.137)$$

The second term of the right hand side can be written as

$$\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) - \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top (\Sigma^{-1})^\top (\mathbf{x} - \boldsymbol{\mu}). \quad (2.138)$$

The second term of the right hand side can be written as

$$-\frac{1}{2} (\Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}))^\top (\mathbf{x} - \boldsymbol{\mu}) = -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}). \quad (2.139)$$

Thus,

$$(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A} (\mathbf{x} - \boldsymbol{\mu}) = 0. \quad (2.140)$$

Hence

$$(\mathbf{x} - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) = (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{S} (\mathbf{x} - \boldsymbol{\mu}). \quad (2.141)$$

2.18

Let Σ be a $D \times D$ real symmetric matrix such that

$$\Sigma \mathbf{u}_i = \lambda_i \mathbf{u}_i, \quad (2.142)$$

where $i = 1, \dots, D$ and \mathbf{u}_i are unit vectors. Taking the inner product with $\bar{\mathbf{u}}_i$ on both sides gives

$$\bar{\mathbf{u}}_i^\top \Sigma \mathbf{u}_i = \lambda_i. \quad (2.143)$$

Since Σ is real and symmetric, the left hand side can be written as

$$\bar{\mathbf{u}}_i^\top \Sigma^\top \mathbf{u}_i = (\bar{\Sigma} \bar{\mathbf{u}}_i)^\top \mathbf{u}_i. \quad (2.144)$$

The right hand side can be written as

$$\bar{\lambda}_i \bar{\mathbf{u}}_i^\top \mathbf{u}_i = \bar{\lambda}_i. \quad (2.145)$$

Therefore,

$$\lambda_i = \bar{\lambda}_i. \quad (2.146)$$

Additionally, for $i \neq j$, taking the inner product with \mathbf{u}_j on both sides of the original equation gives

$$\mathbf{u}_j^\top \Sigma \mathbf{u}_i = \lambda_i \mathbf{u}_j^\top \mathbf{u}_i. \quad (2.147)$$

Since Σ is symmetric, the left hand side can be written as

$$\mathbf{u}_j^\top \Sigma^\top \mathbf{u}_i = (\Sigma \mathbf{u}_j)^\top \mathbf{u}_i. \quad (2.148)$$

The right hand side can be written as $\lambda_j \mathbf{u}_j^\top \mathbf{u}_i$. Therefore,

$$\lambda_i \mathbf{u}_j^\top \mathbf{u}_i = \lambda_j \mathbf{u}_j^\top \mathbf{u}_i. \quad (2.149)$$

Thus, if $\lambda_i \neq \lambda_j$, then

$$\mathbf{u}_j^\top \mathbf{u}_i = 0. \quad (2.150)$$

2.19

Let Σ be a $D \times D$ real symmetric matrix such that

$$\Sigma \mathbf{u}_i = \lambda_i \mathbf{u}_i, \quad (2.151)$$

where $i = 1, \dots, D$ and \mathbf{u}_i are unit vectors. Let

$$\begin{aligned}\mathbf{\Lambda} &= \text{diag}(\lambda_1, \dots, \lambda_D), \\ \mathbf{U} &= [\mathbf{u}_1 \cdots \mathbf{u}_D].\end{aligned}\tag{2.152}$$

Then

$$\mathbf{\Sigma U} = \mathbf{U \Lambda}.\tag{2.153}$$

By 2.18,

$$\mathbf{U}^\top \mathbf{U} = \mathbf{I}.\tag{2.154}$$

Therefore,

$$\begin{aligned}\mathbf{\Sigma} &= \mathbf{U \Lambda U}^\top, \\ \mathbf{\Sigma}^{-1} &= \mathbf{U \Lambda}^{-1} \mathbf{U}^\top,\end{aligned}\tag{2.155}$$

Thus,

$$\begin{aligned}\mathbf{\Sigma} &= \sum_{i=1}^D \lambda_i \mathbf{u}_i \mathbf{u}_i^\top, \\ \mathbf{\Sigma}^{-1} &= \sum_{i=1}^D \frac{1}{\lambda_i} \mathbf{u}_i \mathbf{u}_i^\top.\end{aligned}\tag{2.156}$$

2.20

Let $\mathbf{\Sigma}$ be a $D \times D$ real symmetric matrix such that

$$\mathbf{\Sigma u}_i = \lambda_i \mathbf{u}_i,\tag{2.157}$$

where $i = 1, \dots, D$ and \mathbf{u}_i are unit vectors. Let

$$\begin{aligned}\mathbf{\Lambda} &= \text{diag}(\lambda_1, \dots, \lambda_D), \\ \mathbf{U} &= [\mathbf{u}_1 \cdots \mathbf{u}_D].\end{aligned}\tag{2.158}$$

By 2.19,

$$\mathbf{a}^\top \mathbf{\Sigma a} = \mathbf{b}^\top \mathbf{\Lambda b},\tag{2.159}$$

where

$$\mathbf{b} = \mathbf{U}^\top \mathbf{a}.\tag{2.160}$$

The right hand side can be written as $\sum_{i=1}^D \lambda_i b_i^2$. Therefore, the necessary and sufficient condition for

$$\mathbf{a}^\top \mathbf{\Sigma a} > 0\tag{2.161}$$

for any real vector \mathbf{a} is

$$\lambda_i > 0.\tag{2.162}$$

2.21

Let Σ be a $D \times D$ real symmetric matrix. Then the number of independent parameters is $\frac{D(D+1)}{2}$.

2.22

Let Σ be a $D \times D$ symmetric matrix and

$$\Sigma \Lambda = \mathbf{I}. \quad (2.163)$$

Taking the transpose of the both sides gives

$$\Lambda^\top \Sigma = \mathbf{I}. \quad (2.164)$$

Therefore,

$$\Lambda^\top = \Lambda. \quad (2.165)$$

2.23

Let Σ be a $D \times D$ real symmetric matrix such that

$$\Sigma \mathbf{u}_i = \lambda_i \mathbf{u}_i, \quad (2.166)$$

where $i = 1, \dots, D$ and \mathbf{u}_i are unit vectors. Let

$$\begin{aligned} \Lambda' &= \text{diag} \left(\lambda_1^{-\frac{1}{2}}, \dots, \lambda_D^{-\frac{1}{2}} \right), \\ \mathbf{U} &= [\mathbf{u}_1 \cdots \mathbf{u}_D]. \end{aligned} \quad (2.167)$$

By 2.19,

$$\int_{(\mathbf{x}-\boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}-\boldsymbol{\mu}) = \Delta} d\mathbf{x} = \int_{(\mathbf{x}-\boldsymbol{\mu})^\top \mathbf{U} \Lambda' \Lambda'^\top \mathbf{U}^\top (\mathbf{x}-\boldsymbol{\mu}) = \Delta} d\mathbf{x}. \quad (2.168)$$

By the transformation

$$\mathbf{y} = \Lambda'^\top \mathbf{U}^\top (\mathbf{x} - \boldsymbol{\mu}) \quad (2.169)$$

and the property

$$\mathbf{U}^\top \mathbf{U} = \mathbf{I}, \quad (2.170)$$

the right hand side can be written as

$$\int_{\|\mathbf{y}\|^2=\Delta} \left| \det \left(\mathbf{U} \mathbf{\Lambda}'^{-1} \right) \right| d\mathbf{y} = |\det \mathbf{\Sigma}|^{\frac{1}{2}} \int_{\|\mathbf{y}\|^2=\Delta} d\mathbf{y}. \quad (2.171)$$

Therefore,

$$\int_{(\mathbf{x}-\boldsymbol{\mu})^\top \mathbf{\Sigma}^{-1} (\mathbf{x}-\boldsymbol{\mu})=\Delta} d\mathbf{x} = |\det \mathbf{\Sigma}|^{\frac{1}{2}} \Delta^D V_D, \quad (2.172)$$

where

$$V_D = \int_{\|\mathbf{x}\|=1} d\mathbf{x}. \quad (2.173)$$

2.24

Let

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}$$

be a partitioned matrix where \mathbf{A} is a square matrix and \mathbf{D} is an invertible matrix. By an LDU decomposition, we have

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix} = \begin{bmatrix} \mathbf{I} & \mathbf{B}\mathbf{D}^{-1} \\ \mathbf{O} & \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C} & \mathbf{O} \\ \mathbf{O} & \mathbf{D} \end{bmatrix} \begin{bmatrix} \mathbf{I} & \mathbf{O} \\ \mathbf{D}^{-1}\mathbf{C} & \mathbf{I} \end{bmatrix}.$$

Therefore,

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}^{-1} = \begin{bmatrix} \mathbf{I} & \mathbf{O} \\ -\mathbf{D}^{-1}\mathbf{C} & \mathbf{I} \end{bmatrix} \begin{bmatrix} (\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1} & \mathbf{O} \\ \mathbf{O} & \mathbf{D}^{-1} \end{bmatrix} \begin{bmatrix} \mathbf{I} & -\mathbf{B}\mathbf{D}^{-1} \\ \mathbf{O} & \mathbf{I} \end{bmatrix}.$$

Thus,

$$\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}^{-1} = \begin{bmatrix} (\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1} & -\mathbf{B}\mathbf{D}^{-1}(\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1} \\ -\mathbf{D}^{-1}\mathbf{C}(\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1} & \mathbf{D}^{-1} + \mathbf{D}^{-1}\mathbf{C}(\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1}\mathbf{B}\mathbf{D}^{-1} \end{bmatrix}.$$

2.25

Let \mathbf{x} be a variable in D dimensions such that

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Sigma}), \quad (2.174)$$

where

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_a \\ \mathbf{x}_b \\ \mathbf{x}_c \end{bmatrix}, \boldsymbol{\mu} = \begin{bmatrix} \boldsymbol{\mu}_a \\ \boldsymbol{\mu}_b \\ \boldsymbol{\mu}_c \end{bmatrix}, \boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\Sigma}_{aa} & \boldsymbol{\Sigma}_{ab} & \boldsymbol{\Sigma}_{ac} \\ \boldsymbol{\Sigma}_{ba} & \boldsymbol{\Sigma}_{bb} & \boldsymbol{\Sigma}_{bc} \\ \boldsymbol{\Sigma}_{ca} & \boldsymbol{\Sigma}_{cb} & \boldsymbol{\Sigma}_{cc} \end{bmatrix}.$$

Let

$$\boldsymbol{\Lambda} = \boldsymbol{\Sigma}^{-1}, \quad (2.175)$$

where

$$\boldsymbol{\Lambda} = \begin{bmatrix} \boldsymbol{\Lambda}_{aa} & \boldsymbol{\Lambda}_{ab} & \boldsymbol{\Lambda}_{ac} \\ \boldsymbol{\Lambda}_{ba} & \boldsymbol{\Lambda}_{bb} & \boldsymbol{\Lambda}_{bc} \\ \boldsymbol{\Lambda}_{ca} & \boldsymbol{\Lambda}_{cb} & \boldsymbol{\Lambda}_{cc} \end{bmatrix}.$$

Then

$$\begin{aligned} & -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \\ = & -\frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_a)^\top \boldsymbol{\Lambda}_{aa}(\mathbf{x}_a - \boldsymbol{\mu}_a) - \frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_a)^\top \boldsymbol{\Lambda}_{ab}(\mathbf{x}_b - \boldsymbol{\mu}_b) - \frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_a)^\top \boldsymbol{\Lambda}_{ac}(\mathbf{x}_c - \boldsymbol{\mu}_c) \\ & - \frac{1}{2}(\mathbf{x}_b - \boldsymbol{\mu}_b)^\top \boldsymbol{\Lambda}_{ba}(\mathbf{x}_a - \boldsymbol{\mu}_a) - \frac{1}{2}(\mathbf{x}_b - \boldsymbol{\mu}_b)^\top \boldsymbol{\Lambda}_{bb}(\mathbf{x}_b - \boldsymbol{\mu}_b) - \frac{1}{2}(\mathbf{x}_b - \boldsymbol{\mu}_b)^\top \boldsymbol{\Lambda}_{bc}(\mathbf{x}_c - \boldsymbol{\mu}_c) \\ & - \frac{1}{2}(\mathbf{x}_c - \boldsymbol{\mu}_c)^\top \boldsymbol{\Lambda}_{ca}(\mathbf{x}_a - \boldsymbol{\mu}_a) - \frac{1}{2}(\mathbf{x}_c - \boldsymbol{\mu}_c)^\top \boldsymbol{\Lambda}_{cb}(\mathbf{x}_b - \boldsymbol{\mu}_b) - \frac{1}{2}(\mathbf{x}_c - \boldsymbol{\mu}_c)^\top \boldsymbol{\Lambda}_{cc}(\mathbf{x}_c - \boldsymbol{\mu}_c). \end{aligned} \quad (2.176)$$

Excluding the terms independent of \mathbf{x}_a , the right hand side can be written as

$$-\frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_{a|b,c})^\top \boldsymbol{\Sigma}_{a|b,c}^{-1}(\mathbf{x}_a - \boldsymbol{\mu}_{a|b,c}), \quad (2.177)$$

where

$$\begin{aligned} \boldsymbol{\mu}_{a|b,c} &= \boldsymbol{\mu}_a - \boldsymbol{\Lambda}_{aa}^{-1} \boldsymbol{\Lambda}_{ab}(\mathbf{x}_b - \boldsymbol{\mu}_b) - \boldsymbol{\Lambda}_{aa}^{-1} \boldsymbol{\Lambda}_{ac}(\mathbf{x}_c - \boldsymbol{\mu}_c), \\ \boldsymbol{\Sigma}_{a|b,c} &= \boldsymbol{\Lambda}_{aa}^{-1}. \end{aligned} \quad (2.178)$$

Therefore,

$$p(\mathbf{x}_a | \mathbf{x}_b, \mathbf{x}_c) = \mathcal{N}(\mathbf{x}_a | \boldsymbol{\mu}_{a|b,c}, \boldsymbol{\Sigma}_{a|b,c}). \quad (2.179)$$

Multiplying both sides by $p(\mathbf{x}_c)$ and integrating both sides with respect to \mathbf{x}_c gives

$$p(\mathbf{x}_a | \mathbf{x}_b) = \int \mathcal{N}(\mathbf{x}_a | \boldsymbol{\mu}_{a|b,c}, \boldsymbol{\Sigma}_{a|b,c}) p(\mathbf{x}_c) d\mathbf{x}_c. \quad (2.180)$$

Thus,

$$p(\mathbf{x}_a | \mathbf{x}_b) = \mathcal{N}(\mathbf{x}_a | \boldsymbol{\mu}_{a|b}, \boldsymbol{\Sigma}_{a|b}), \quad (2.181)$$

where

$$\begin{aligned}\boldsymbol{\mu}_{a|b} &= \boldsymbol{\mu}_a - \boldsymbol{\Lambda}_{aa}^{-1} \boldsymbol{\Lambda}_{ab} (\mathbf{x}_b - \boldsymbol{\mu}_b) + \boldsymbol{\Lambda}_{aa}^{-1} \boldsymbol{\Lambda}_{ac} \boldsymbol{\mu}_c, \\ \boldsymbol{\Sigma}_{a|b} &= \boldsymbol{\Lambda}_{aa}^{-1}.\end{aligned}\tag{2.182}$$

2.26

We have

$$\begin{aligned}& \left(\mathbf{A}^{-1} - \mathbf{A}^{-1} \mathbf{B} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} \mathbf{D} \mathbf{A}^{-1} \right) (\mathbf{A} + \mathbf{B} \mathbf{C} \mathbf{D}) \\ &= \mathbf{I} - \mathbf{A}^{-1} \mathbf{B} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} \mathbf{D} + \mathbf{A}^{-1} \mathbf{B} \mathbf{C} \mathbf{D} \\ & \quad - \mathbf{A}^{-1} \mathbf{B} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} \mathbf{D} \mathbf{A}^{-1} \mathbf{B} \mathbf{C} \mathbf{D}.\end{aligned}\tag{2.183}$$

The right hand side except the first term can be written as

$$\begin{aligned}& \mathbf{A}^{-1} \mathbf{B} \left(\mathbf{C} - (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} (\mathbf{I} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B} \mathbf{C}) \right) \mathbf{D} \\ &= \mathbf{A}^{-1} \mathbf{B} \left(\mathbf{C} - (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B}) \mathbf{C} \right) \mathbf{D}.\end{aligned}\tag{2.184}$$

The right hand side can be written as

$$\mathbf{A}^{-1} \mathbf{B} (\mathbf{C} - \mathbf{C}) \mathbf{D} = \mathbf{O}.\tag{2.185}$$

Therefore,

$$\left(\mathbf{A}^{-1} - \mathbf{A}^{-1} \mathbf{B} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} \mathbf{D} \mathbf{A}^{-1} \right) (\mathbf{A} + \mathbf{B} \mathbf{C} \mathbf{D}) = \mathbf{I}.\tag{2.186}$$

Thus,

$$(\mathbf{A} + \mathbf{B} \mathbf{C} \mathbf{D})^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1} \mathbf{B} (\mathbf{C}^{-1} + \mathbf{D} \mathbf{A}^{-1} \mathbf{B})^{-1} \mathbf{D} \mathbf{A}^{-1}.\tag{2.187}$$

2.27

Let \mathbf{x} and \mathbf{z} be two variables. Then

$$\mathbf{E}(\mathbf{x} + \mathbf{z}) = \int \int (\mathbf{x} + \mathbf{z}) p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z}.\tag{2.188}$$

The right hand side can be written as

$$\int \mathbf{x} \left(\int p(\mathbf{x}, \mathbf{z}) d\mathbf{z} \right) d\mathbf{x} + \int \mathbf{z} \left(\int p(\mathbf{x}, \mathbf{z}) d\mathbf{x} \right) d\mathbf{z} = \int \mathbf{x} p(\mathbf{x}) d\mathbf{x} + \int \mathbf{z} p(\mathbf{z}) d\mathbf{z}.\tag{2.189}$$

The right hand side can be written as $E \mathbf{x} + E \mathbf{z}$. Therefore,

$$E(\mathbf{x} + \mathbf{z}) = E \mathbf{x} + E \mathbf{z}. \quad (2.190)$$

Additionally,

$$\text{cov}(\mathbf{x} + \mathbf{z}) = \int \int (\mathbf{x} + \mathbf{z} - E(\mathbf{x} + \mathbf{z})) (\mathbf{x} + \mathbf{z} - E(\mathbf{x} + \mathbf{z}))^\top p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z}. \quad (2.191)$$

The right hand side can be written as

$$\begin{aligned} & \int \int (\mathbf{x} - E \mathbf{x}) (\mathbf{x} - E \mathbf{x})^\top p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z} + \int \int (\mathbf{x} - E \mathbf{x}) (\mathbf{z} - E \mathbf{z})^\top p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z} \\ & + \int \int (\mathbf{z} - E \mathbf{z}) (\mathbf{x} - E \mathbf{x})^\top p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z} + \int \int (\mathbf{z} - E \mathbf{z}) (\mathbf{z} - E \mathbf{z})^\top p(\mathbf{x}, \mathbf{z}) d\mathbf{x} d\mathbf{z}. \end{aligned} \quad (2.192)$$

The first and fourth terms can be written as $\text{cov } \mathbf{x}$ and $\text{cov } \mathbf{z}$. If \mathbf{x} and \mathbf{z} are independent, the second and third terms can be written as

$$\begin{aligned} & \int (\mathbf{x} - E \mathbf{x}) p(\mathbf{x}) d\mathbf{x} \int (\mathbf{z} - E \mathbf{z})^\top p(\mathbf{z}) d\mathbf{z} = \mathbf{O}, \\ & \int (\mathbf{z} - E \mathbf{z}) p(\mathbf{z}) d\mathbf{z} \int (\mathbf{x} - E \mathbf{x})^\top p(\mathbf{x}) d\mathbf{x} = \mathbf{O}. \end{aligned} \quad (2.193)$$

Therefore,

$$\text{cov}(\mathbf{x} + \mathbf{z}) = \text{cov } \mathbf{x} + \text{cov } \mathbf{z}. \quad (2.194)$$

2.28

Let \mathbf{x} and \mathbf{y} be Gaussian variables and

$$\mathbf{z} = \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix},$$

where

$$E \mathbf{z} = \begin{bmatrix} \boldsymbol{\mu} \\ \mathbf{A}\boldsymbol{\mu} + \mathbf{b} \end{bmatrix}$$

and

$$\text{cov } \mathbf{z} = \begin{bmatrix} \boldsymbol{\Lambda}^{-1} & \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top \\ \mathbf{A} \boldsymbol{\Lambda}^{-1} & \mathbf{L}^{-1} + \mathbf{A} \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top \end{bmatrix}.$$

Then, by 2.29,

$$(\text{cov } \mathbf{z})^{-1} = \begin{bmatrix} \mathbf{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A} & -\mathbf{A}^\top \mathbf{L} \\ -\mathbf{L} \mathbf{A} & \mathbf{L} \end{bmatrix}.$$

Then, $\ln p(\mathbf{z})$ except the terms independent of \mathbf{x} and \mathbf{y} is given by

$$\begin{aligned} & -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top (\mathbf{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A}) (\mathbf{x} - \boldsymbol{\mu}) + \frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A}^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b}) \\ & + \frac{1}{2}(\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b})^\top \mathbf{L} \mathbf{A} (\mathbf{x} - \boldsymbol{\mu}) - \frac{1}{2}(\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b}) \\ = & -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top (\mathbf{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A}) (\mathbf{x} - \boldsymbol{\mu}) + \\ & -\frac{1}{2}(\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b} - \mathbf{A}(\mathbf{x} - \boldsymbol{\mu}))^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b} - \mathbf{A}(\mathbf{x} - \boldsymbol{\mu})) \\ & -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A}^\top \mathbf{L} \mathbf{A} (\mathbf{x} - \boldsymbol{\mu}). \end{aligned} \tag{2.195}$$

The right hand side can be written as

$$-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Lambda} (\mathbf{x} - \boldsymbol{\mu}) - \frac{1}{2}(\mathbf{y} - \mathbf{A}\mathbf{x} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\mathbf{x} - \mathbf{b}). \tag{2.196}$$

Therefore,

$$\begin{aligned} p(\mathbf{x}) &= \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \mathbf{\Lambda}^{-1}), \\ p(\mathbf{y} | \mathbf{x}) &= \mathcal{N}(\mathbf{y} | \mathbf{A}\mathbf{x} + \mathbf{b}, \mathbf{L}^{-1}). \end{aligned} \tag{2.197}$$

2.29

Let

$$\mathbf{R} = \begin{bmatrix} \mathbf{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A} & -\mathbf{A}^\top \mathbf{L} \\ -\mathbf{L} \mathbf{A} & \mathbf{L} \end{bmatrix}.$$

Then, by 2.24,

$$\mathbf{R}^{-1} = \begin{bmatrix} \mathbf{\Lambda}^{-1} & \mathbf{\Lambda}^{-1} \mathbf{A}^\top \\ \mathbf{A} \mathbf{\Lambda}^{-1} & \mathbf{L}^{-1} + \mathbf{A} \mathbf{\Lambda}^{-1} \mathbf{A}^\top \end{bmatrix}.$$

2.30

Let

$$\mathbf{R}^{-1} = \begin{bmatrix} \mathbf{\Lambda}^{-1} & \mathbf{\Lambda}^{-1} \mathbf{A}^\top \\ \mathbf{A} \mathbf{\Lambda}^{-1} & \mathbf{L}^{-1} + \mathbf{A} \mathbf{\Lambda}^{-1} \mathbf{A}^\top \end{bmatrix}.$$

Then

$$\mathbf{R}^{-1} \begin{bmatrix} \mathbf{A}\boldsymbol{\mu} - \mathbf{A}^\top \mathbf{L}\mathbf{b} \\ \mathbf{L}\mathbf{b} \end{bmatrix} = \begin{bmatrix} \boldsymbol{\mu} \\ \mathbf{A}\boldsymbol{\mu} + \mathbf{b} \end{bmatrix}.$$

2.31

Let \mathbf{y} be a variable such that

$$\mathbf{y} = \mathbf{x} + \mathbf{z}, \quad (2.198)$$

where

$$\begin{aligned} p(\mathbf{x}) &= \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_{\mathbf{x}}, \boldsymbol{\Sigma}_{\mathbf{x}}), \\ p(\mathbf{z}) &= \mathcal{N}(\mathbf{z} | \boldsymbol{\mu}_{\mathbf{z}}, \boldsymbol{\Sigma}_{\mathbf{z}}). \end{aligned} \quad (2.199)$$

By marginalisation,

$$p(\mathbf{y}) = \int p(\mathbf{y} | \mathbf{x}) p(\mathbf{x}) d\mathbf{x}. \quad (2.200)$$

The right hand side can be written as

$$\int \mathcal{N}(\mathbf{y} | \mathbf{x} + \boldsymbol{\mu}_{\mathbf{z}}, \boldsymbol{\Sigma}_{\mathbf{z}}) \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_{\mathbf{x}}, \boldsymbol{\Sigma}_{\mathbf{x}}) d\mathbf{x}. \quad (2.201)$$

The logarithm of the integrand except the terms independent of \mathbf{x} and \mathbf{y} is given by

$$-\frac{1}{2}(\mathbf{y} - \mathbf{x} - \boldsymbol{\mu}_{\mathbf{z}})^\top \boldsymbol{\Sigma}_{\mathbf{z}}^{-1}(\mathbf{y} - \mathbf{x} - \boldsymbol{\mu}_{\mathbf{z}}) - \frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{x}})^\top \boldsymbol{\Sigma}_{\mathbf{x}}^{-1}(\mathbf{x} - \boldsymbol{\mu}_{\mathbf{x}}). \quad (2.202)$$

The first and second order terms can be written as

$$-\mathbf{x}^\top (\boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \boldsymbol{\mu}_{\mathbf{z}} - \boldsymbol{\Sigma}_{\mathbf{x}}^{-1} \boldsymbol{\mu}_{\mathbf{x}}) + \mathbf{y}^\top \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \boldsymbol{\mu}_{\mathbf{z}} = \mathbf{u}^\top \mathbf{v} \quad (2.203)$$

and

$$-\frac{1}{2} \mathbf{x}^\top (\boldsymbol{\Sigma}_{\mathbf{x}}^{-1} + \boldsymbol{\Sigma}_{\mathbf{z}}^{-1}) \mathbf{x} + \frac{1}{2} \mathbf{x}^\top \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \mathbf{y} + \frac{1}{2} \mathbf{y}^\top \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \mathbf{x} - \frac{1}{2} \mathbf{y}^\top \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \mathbf{y} = -\frac{1}{2} \mathbf{u}^\top \mathbf{R} \mathbf{u}, \quad (2.204)$$

respectively, where

$$\mathbf{u} = \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix}, \mathbf{v} = \begin{bmatrix} \boldsymbol{\Sigma}_{\mathbf{x}}^{-1} \boldsymbol{\mu}_{\mathbf{x}} - \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \boldsymbol{\mu}_{\mathbf{z}} \\ \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \boldsymbol{\mu}_{\mathbf{z}} \end{bmatrix}, \mathbf{R} = \begin{bmatrix} \boldsymbol{\Sigma}_{\mathbf{x}}^{-1} & -\boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \\ -\boldsymbol{\Sigma}_{\mathbf{z}}^{-1} & \boldsymbol{\Sigma}_{\mathbf{z}}^{-1} \end{bmatrix}.$$

Therefore, the logarithm of the integrand except the terms independent of \mathbf{u} can be written as

$$-\frac{1}{2} (\mathbf{u} - \mathbf{R}^{-1}\mathbf{v})^\top \mathbf{R} (\mathbf{u} - \mathbf{R}^{-1}\mathbf{v}), \quad (2.205)$$

where

$$\mathbf{R}^{-1} = \begin{bmatrix} \Sigma_{\mathbf{x}} & \Sigma_{\mathbf{x}} \\ \Sigma_{\mathbf{x}} & \Sigma_{\mathbf{x}} + \Sigma_{\mathbf{z}} \end{bmatrix}, \mathbf{R}^{-1}\mathbf{v} = \begin{bmatrix} \mu_{\mathbf{x}} \\ \mu_{\mathbf{x}} + \mu_{\mathbf{z}} \end{bmatrix}.$$

by 2.29 and 2.30. Thus,

$$p(\mathbf{y}) = \mathcal{N}(\mathbf{y} | \mu_{\mathbf{x}} + \mu_{\mathbf{z}}, \Sigma_{\mathbf{x}} + \Sigma_{\mathbf{z}}). \quad (2.206)$$

2.32

Let \mathbf{x} and \mathbf{y} be variables such that

$$\begin{aligned} p(\mathbf{x}) &= \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Lambda}^{-1}), \\ p(\mathbf{y} | \mathbf{x}) &= \mathcal{N}(\mathbf{y} | \mathbf{A}\mathbf{x} + \mathbf{b}, \mathbf{L}^{-1}). \end{aligned} \quad (2.207)$$

By the Bayes' theorem,

$$p(\mathbf{y} | \mathbf{x})p(\mathbf{x}) = p(\mathbf{x} | \mathbf{y})p(\mathbf{y}). \quad (2.208)$$

The logarithm of the left hand side except the terms independent of \mathbf{x} and \mathbf{y} is given by

$$-\frac{1}{2} (\mathbf{y} - \mathbf{A}\mathbf{x} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\mathbf{x} - \mathbf{b}) - \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda} (\mathbf{x} - \boldsymbol{\mu}). \quad (2.209)$$

Since the first term can be written as

$$\begin{aligned} & -\frac{1}{2} (\mathbf{y} - \mathbf{A}(\mathbf{x} - \boldsymbol{\mu}) - \mathbf{A}\boldsymbol{\mu} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A}(\mathbf{x} - \boldsymbol{\mu}) - \mathbf{A}\boldsymbol{\mu} - \mathbf{b}) \\ &= -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A}^\top \mathbf{L} \mathbf{A} (\mathbf{x} - \boldsymbol{\mu}) + (\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{A}^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b}) \\ & \quad -\frac{1}{2} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A}\boldsymbol{\mu} - \mathbf{b}), \end{aligned} \quad (2.210)$$

the logarithm except the terms independent of \mathbf{x} and \mathbf{y} can be written as

$$\begin{aligned}
& -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu} - \mathbf{z})^\top (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A}) (\mathbf{x} - \boldsymbol{\mu} - \mathbf{z}) + \frac{1}{2} \mathbf{z}^\top (\mathbf{A}^\top \mathbf{L} \mathbf{A} + \boldsymbol{\Lambda}) \mathbf{z} \\
& -\frac{1}{2}(\mathbf{y} - \mathbf{A} \boldsymbol{\mu} - \mathbf{b})^\top \mathbf{L} (\mathbf{y} - \mathbf{A} \boldsymbol{\mu} - \mathbf{b}) \\
& = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu} - \mathbf{z})^\top (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A}) (\mathbf{x} - \boldsymbol{\mu} - \mathbf{z}) \\
& -\frac{1}{2}(\mathbf{y} - \mathbf{A} \boldsymbol{\mu} - \mathbf{b})^\top \mathbf{M} (\mathbf{y} - \mathbf{A} \boldsymbol{\mu} - \mathbf{b}),
\end{aligned} \tag{2.211}$$

where

$$\begin{aligned}
\mathbf{z} &= (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{L} (\mathbf{y} - \mathbf{A} \boldsymbol{\mu} - \mathbf{b}), \\
\mathbf{M} &= \mathbf{L} - \mathbf{L} \mathbf{A} (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{L}.
\end{aligned} \tag{2.212}$$

we have

$$\boldsymbol{\mu} + \mathbf{z} = (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1} (\mathbf{A}^\top \mathbf{L} (\mathbf{y} - \mathbf{b}) + \boldsymbol{\Lambda} \boldsymbol{\mu}). \tag{2.213}$$

By 2.26,

$$(\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1} = \boldsymbol{\Lambda}^{-1} - \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top (\mathbf{L}^{-1} + \mathbf{A} \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top)^{-1} \mathbf{A} \boldsymbol{\Lambda}^{-1}. \tag{2.214}$$

Therefore,

$$\mathbf{M} = (\mathbf{L}^{-1} + \mathbf{A} \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top)^{-1}. \tag{2.215}$$

Thus,

$$\begin{aligned}
p(\mathbf{x}|\mathbf{y}) &= \mathcal{N}(\mathbf{x} | (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1} (\mathbf{A}^\top \mathbf{L} (\mathbf{y} - \mathbf{b}) + \boldsymbol{\Lambda} \boldsymbol{\mu}), (\boldsymbol{\Lambda} + \mathbf{A}^\top \mathbf{L} \mathbf{A})^{-1}), \\
p(\mathbf{y}) &= \mathcal{N}(\mathbf{y} | \mathbf{A} \boldsymbol{\mu} + \mathbf{b}, \mathbf{L}^{-1} + \mathbf{A} \boldsymbol{\Lambda}^{-1} \mathbf{A}^\top).
\end{aligned} \tag{2.216}$$

2.33

Refer to 2.32.

2.34

Let \mathbf{X} be a set of N variables such that

$$\ln p(\mathbf{X} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = -\frac{ND}{2} \ln(2\pi) - \frac{N}{2} \ln(\det \boldsymbol{\Sigma}) - \frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}). \tag{2.217}$$

To maximise it with respect to $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$, setting the partial derivatives to zero gives

$$\begin{aligned}\mathbf{0} &= \sum_{n=1}^N (\boldsymbol{\Sigma}^{-1} + (\boldsymbol{\Sigma}^{-1})^\top) (\mathbf{x}_n - \boldsymbol{\mu}), \\ \mathbf{O} &= -\frac{N}{2} (\boldsymbol{\Sigma}^{-1})^\top + \frac{1}{2} (\boldsymbol{\Sigma}^{-1})^2 \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})(\mathbf{x}_n - \boldsymbol{\mu})^\top.\end{aligned}\tag{2.218}$$

Therefore,

$$\begin{aligned}\boldsymbol{\mu}_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n, \\ \boldsymbol{\Sigma}_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})(\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top.\end{aligned}\tag{2.219}$$

2.35

Let \mathbf{x} be a variable such that

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}).\tag{2.220}$$

Then

$$\mathbb{E} \mathbf{x} \mathbf{x}^\top = \int \mathbf{x} \mathbf{x}^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x}.\tag{2.221}$$

The right hand side can be written as

$$\begin{aligned}& \int (\mathbf{x} - \boldsymbol{\mu} + \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu} + \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ &= \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} + \boldsymbol{\mu} \int (\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \\ &+ \left(\int (\mathbf{x} - \boldsymbol{\mu}) \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} \right) \boldsymbol{\mu}^\top + \boldsymbol{\mu} \boldsymbol{\mu}^\top \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x}.\end{aligned}\tag{2.222}$$

Since

$$\begin{aligned}\int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} &= 1, \\ \int \mathbf{x} \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} &= \boldsymbol{\mu}, \\ \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) d\mathbf{x} &= \boldsymbol{\Sigma},\end{aligned}\tag{2.223}$$

the right hand side can be written as $\Sigma + \boldsymbol{\mu}\boldsymbol{\mu}^\top$. Therefore,

$$\mathbb{E} \mathbf{x}\mathbf{x}^\top = \Sigma + \boldsymbol{\mu}\boldsymbol{\mu}^\top. \quad (2.224)$$

Additionally, let \mathbf{x}_n and \mathbf{x}_m be variables such that

$$\begin{aligned} p(\mathbf{x}_n) &= \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}, \Sigma), \\ p(\mathbf{x}_m) &= \mathcal{N}(\mathbf{x}_m | \boldsymbol{\mu}, \Sigma). \end{aligned} \quad (2.225)$$

If $n \neq m$, then

$$\mathbb{E} \mathbf{x}_n \mathbf{x}_m^\top = \mathbb{E} \mathbf{x}_n \mathbb{E} \mathbf{x}_m^\top. \quad (2.226)$$

The right hand side can be written as $\boldsymbol{\mu}\boldsymbol{\mu}^\top$. Therefore,

$$\mathbb{E} \mathbf{x}_n \mathbf{x}_m^\top = \delta_{nm} \Sigma + \boldsymbol{\mu}\boldsymbol{\mu}^\top. \quad (2.227)$$

Finally, let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables such that

$$p(\mathbf{x}_n) = \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}, \Sigma). \quad (2.228)$$

By 2.34,

$$\begin{aligned} \boldsymbol{\mu}_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n, \\ \Sigma_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})(\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top. \end{aligned} \quad (2.229)$$

Then

$$\mathbb{E} \Sigma_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N \mathbb{E} (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})(\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top. \quad (2.230)$$

The right hand side can be written as

$$\begin{aligned} & \frac{1}{N} \sum_{n=1}^N \mathbb{E} \mathbf{x}_n \mathbf{x}_n^\top - \frac{1}{N^2} \sum_{n=1}^N \mathbb{E} \left(\sum_{n=1}^N \mathbf{x}_n \right) \mathbf{x}_n^\top - \frac{1}{N^2} \sum_{n=1}^N \mathbb{E} \mathbf{x}_n \left(\sum_{n=1}^N \mathbf{x}_n \right)^\top \\ & + \frac{1}{N^3} \sum_{n=1}^N \mathbb{E} \left(\sum_{n=1}^N \mathbf{x}_n \right) \left(\sum_{n=1}^N \mathbf{x}_n \right)^\top. \end{aligned} \quad (2.231)$$

The first term can be written as $\Sigma + \boldsymbol{\mu}\boldsymbol{\mu}^\top$. The second and third terms can be written as

$$-\frac{1}{N} ((\Sigma + \boldsymbol{\mu}\boldsymbol{\mu}^\top) + (N-1)\boldsymbol{\mu}\boldsymbol{\mu}^\top) = -\frac{1}{N} \Sigma - \boldsymbol{\mu}\boldsymbol{\mu}^\top. \quad (2.232)$$

The fourth term can be written as

$$\frac{1}{N^2} (N (\mathbf{\Sigma} + \boldsymbol{\mu}\boldsymbol{\mu}^\top) + N(N-1)\boldsymbol{\mu}\boldsymbol{\mu}^\top) = \frac{1}{N}\mathbf{\Sigma} + \boldsymbol{\mu}\boldsymbol{\mu}^\top. \quad (2.233)$$

Therefore,

$$\mathbb{E} \mathbf{\Sigma}_{\text{ML}} = \frac{N-1}{N}\mathbf{\Sigma}. \quad (2.234)$$

2.36

Let x_1, \dots, x_N be variables such that

$$p(x_n) = \mathcal{N}(x_n | \mu, \sigma^2). \quad (2.235)$$

Let us assume that μ is known. Then, by 2.34,

$$\sigma_{\text{ML}}^{2(N)} = \frac{1}{N} \sum_{n=1}^N (x_n - \mu)^2. \quad (2.236)$$

The right hand side can be written as

$$\frac{1}{N}(x_N - \mu)^2 + \frac{1}{N} \sum_{n=1}^{N-1} (x_n - \mu)^2 = \frac{1}{N}(x_N - \mu)^2 + \frac{N-1}{N} \sigma_{\text{ML}}^{2(N-1)}. \quad (2.237)$$

Therefore,

$$\sigma_{\text{ML}}^{2(N)} = \sigma_{\text{ML}}^{2(N-1)} + \frac{1}{N} \left((x_N - \mu)^2 - \sigma_{\text{ML}}^{2(N-1)} \right). \quad (2.238)$$

Since

$$\frac{\partial}{\partial \sigma^2} (-\ln p(x_n | \sigma^2)) = \frac{1}{2\sigma^2} - \frac{1}{2(\sigma^2)^2} (x_n - \mu)^2, \quad (2.239)$$

we have

$$\sigma_{\text{ML}}^{2(N)} = \sigma_{\text{ML}}^{2(N-1)} - \frac{\sigma_{\text{ML}}^{2(N-1)}}{N} \frac{\partial}{\partial \sigma_{\text{ML}}^{2(N-1)}} \left(-\ln p(x_N | \sigma_{\text{ML}}^{2(N-1)}) \right). \quad (2.240)$$

2.37

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables such that

$$p(\mathbf{x}_n) = \mathcal{N}(\mathbf{x}_n | \boldsymbol{\mu}, \boldsymbol{\Sigma}). \quad (2.241)$$

Let us assume that $\boldsymbol{\mu}$ is known. Then, by 2.34,

$$\boldsymbol{\Sigma}_{\text{ML}}^{(N)} = \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})(\mathbf{x}_n - \boldsymbol{\mu})^\top. \quad (2.242)$$

The right hand side can be written as

$$\begin{aligned} & \frac{1}{N} (\mathbf{x}_N - \boldsymbol{\mu})(\mathbf{x}_N - \boldsymbol{\mu})^\top + \frac{1}{N} \sum_{n=1}^{N-1} (\mathbf{x}_n - \boldsymbol{\mu})(\mathbf{x}_n - \boldsymbol{\mu})^\top \\ &= \frac{1}{N} (\mathbf{x}_N - \boldsymbol{\mu})(\mathbf{x}_N - \boldsymbol{\mu})^\top + \frac{N-1}{N} \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)}. \end{aligned} \quad (2.243)$$

Therefore,

$$\boldsymbol{\Sigma}_{\text{ML}}^{(N)} = \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)} + \frac{1}{N} \left((\mathbf{x}_N - \boldsymbol{\mu})(\mathbf{x}_N - \boldsymbol{\mu})^\top - \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)} \right). \quad (2.244)$$

Since

$$\frac{\partial}{\partial \boldsymbol{\Sigma}} (-\ln p(\mathbf{x}_n | \boldsymbol{\Sigma})) = -\frac{1}{2} (\boldsymbol{\Sigma}^{-1})^\top + \frac{1}{2} (\boldsymbol{\Sigma}^{-1})^2 (\mathbf{x}_n - \boldsymbol{\mu})(\mathbf{x}_n - \boldsymbol{\mu})^\top, \quad (2.245)$$

we have

$$\boldsymbol{\Sigma}_{\text{ML}}^{(N)} = \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)} - \frac{\boldsymbol{\Sigma}_{\text{ML}}^{(N-1)}}{N} \frac{\partial}{\partial \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)}} \left(-\ln p(\mathbf{x}_N | \boldsymbol{\Sigma}_{\text{ML}}^{(N-1)}) \right). \quad (2.246)$$

2.38

Let x_1, \dots, x_N be variables such that

$$\begin{aligned} p(x_n | \mu) &= \mathcal{N}(x_n | \mu, \sigma^2), \\ p(\mu) &= \mathcal{N}(\mu | \mu_0, \sigma_0^2). \end{aligned} \quad (2.247)$$

By the Bayes' theorem,

$$p(\mu | \mathbf{x}) p(\mathbf{x}) = p(\mathbf{x} | \mu) p(\mu). \quad (2.248)$$

The logarithm of the right hand side except the terms independent of \mathbf{x} and μ can be written as

$$-\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu)^2 - \frac{1}{2\sigma_0^2} (\mu - \mu_0)^2. \quad (2.249)$$

The first term can be written as

$$-\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu_{\text{ML}} + \mu_{\text{ML}} - \mu)^2 = -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2 - \frac{N}{2\sigma^2} (\mu_{\text{ML}} - \mu)^2. \quad (2.250)$$

where

$$\mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N x_n, \quad (2.251)$$

as derived in 2.34. Therefore, the logarithm except the terms independent of \mathbf{x} and μ can be written as

$$\begin{aligned} & -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2 - \frac{N}{2\sigma^2} (\mu_{\text{ML}} - \mu)^2 - \frac{1}{2\sigma_0^2} (\mu - \mu_0)^2 \\ & = -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2 - \frac{1}{2\sigma_N^2} (\mu - \mu_N)^2 + \frac{\mu_N^2}{2\sigma_N^2}, \end{aligned} \quad (2.252)$$

where

$$\begin{aligned} \mu_N &= \frac{N\sigma_0^2}{N\sigma_0^2 + \sigma^2} \mu_{\text{ML}} + \frac{\sigma^2}{N\sigma_0^2 + \sigma^2} \mu_0, \\ \sigma_N^2 &= \frac{\sigma^2 \sigma_0^2}{N\sigma_0^2 + \sigma^2}. \end{aligned} \quad (2.253)$$

Therefore,

$$p(\mu|\mathbf{x}) = \mathcal{N} \left(\mu \mid \frac{N\sigma_0^2}{N\sigma_0^2 + \sigma^2} \mu_{\text{ML}} + \frac{\sigma^2}{N\sigma_0^2 + \sigma^2} \mu_0, \frac{\sigma^2 \sigma_0^2}{N\sigma_0^2 + \sigma^2} \right). \quad (2.254)$$

2.39 (Incomplete)

Let x_1, \dots, x_N be variables such that

$$\begin{aligned} p(x_n|\mu) &= \mathcal{N}(x_n|\mu, \sigma^2), \\ p(\mu) &= \mathcal{N}(\mu|\mu_0, \sigma_0^2). \end{aligned} \quad (2.255)$$

Then, by 2.38,

$$p(\mu|\mathbf{x}) = \mathcal{N}(\mu|\mu_N, \sigma_N^2), \quad (2.256)$$

where

$$\begin{aligned} \mu_N &= \frac{\sigma_0^2}{N\sigma_0^2 + \sigma^2} \sum_{n=1}^N x_n + \frac{\sigma^2}{N\sigma_0^2 + \sigma^2} \mu_0, \\ \sigma_N^2 &= \frac{\sigma^2 \sigma_0^2}{N\sigma_0^2 + \sigma^2}. \end{aligned} \quad (2.257)$$

Then

$$\begin{aligned} \mu_N &= \frac{(N-1)\sigma_0^2 + \sigma^2}{N\sigma_0^2 + \sigma^2} \mu_{N-1} + \frac{\sigma_0^2}{N\sigma_0^2 + \sigma^2} x_N + \frac{\sigma^2 - \sigma_0^2}{N\sigma_0^2 + \sigma^2} \mu_0, \\ \sigma_N^2 &= \frac{(N-1)\sigma_0^2 + \sigma^2}{N\sigma_0^2 + \sigma^2} \sigma_{N-1}^2. \end{aligned} \quad (2.258)$$

Additionally, we have

$$\begin{aligned} p(\mu|x_1, \dots, x_{N-1}) &= \mathcal{N}(\mu|\mu_{N-1}, \sigma_{N-1}^2), \\ p(x_N|\mu) &= \mathcal{N}(x_N|\mu, \sigma^2). \end{aligned} \quad (2.259)$$

Then, $\ln p(\mu|x_1, \dots, x_{N-1}) + \ln p(x_N|\mu)$ except the terms independent of μ or x_N can be written as

$$\begin{aligned} & -\frac{1}{2\sigma_{N-1}^2}(\mu - \mu_{N-1})^2 - \frac{1}{2\sigma^2}(x_N - \mu)^2 \\ &= -\frac{1}{2\frac{\sigma_{N-1}^2 \sigma^2}{\sigma_{N-1}^2 + \sigma^2}} \left(\mu - \frac{\sigma^2}{\sigma_{N-1}^2 + \sigma^2} \mu_{N-1} - \frac{\sigma_{N-1}^2}{\sigma_{N-1}^2 + \sigma^2} x_N \right)^2 \\ &+ \frac{1}{2\frac{\sigma_{N-1}^2 \sigma^2}{\sigma_{N-1}^2 + \sigma^2}} \left(\frac{\sigma^2}{\sigma_{N-1}^2 + \sigma^2} \mu_{N-1} + \frac{\sigma_{N-1}^2}{\sigma_{N-1}^2 + \sigma^2} x_N \right) - \frac{\mu_{N-1}^2}{2\sigma_{N-1}^2} - \frac{x_N^2}{2\sigma^2}. \end{aligned} \quad (2.260)$$

Therefoere,

$$\begin{aligned} \mu_N &= \frac{\sigma^2}{\sigma_{N-1}^2 + \sigma^2} \mu_{N-1} + \frac{\sigma_{N-1}^2}{\sigma_{N-1}^2 + \sigma^2} x_N, \\ \sigma_N^2 &= \frac{\sigma_{N-1}^2 \sigma^2}{\sigma_{N-1}^2 + \sigma^2}. \end{aligned} \quad (2.261)$$

2.40

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables such that

$$\begin{aligned} p(\mathbf{x}_n|\boldsymbol{\mu}) &= \mathcal{N}(\mathbf{x}_n|\boldsymbol{\mu}, \boldsymbol{\Sigma}), \\ p(\boldsymbol{\mu}) &= \mathcal{N}(\boldsymbol{\mu}|\boldsymbol{\mu}_0, \boldsymbol{\Sigma}). \end{aligned} \quad (2.262)$$

By the Bayes' theorem,

$$p(\boldsymbol{\mu}|\mathbf{X})p(\mathbf{X}) = p(\mathbf{X}|\boldsymbol{\mu})p(\boldsymbol{\mu}). \quad (2.263)$$

The logarithm of the right hand side except the terms independent of \mathbf{X} and $\boldsymbol{\mu}$ can be written as

$$-\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) - \frac{1}{2} (\boldsymbol{\mu} - \boldsymbol{\mu}_0)^\top \boldsymbol{\Sigma}_0^{-1} (\boldsymbol{\mu} - \boldsymbol{\mu}_0). \quad (2.264)$$

The first term can be written as

$$\begin{aligned} & -\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}} + \boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}} + \boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu}) \\ &= -\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}}) - \frac{N}{2} (\boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu}). \end{aligned} \quad (2.265)$$

where

$$\boldsymbol{\mu}_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n, \quad (2.266)$$

as derived in 2.34. Therefore, the logarithm except the terms independent of \mathbf{X} and $\boldsymbol{\mu}$ can be written as

$$\begin{aligned} & -\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}}) - \frac{N}{2} (\boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_{\text{ML}} - \boldsymbol{\mu}) \\ & -\frac{1}{2} (\boldsymbol{\mu} - \boldsymbol{\mu}_0)^\top \boldsymbol{\Sigma}_0^{-1} (\boldsymbol{\mu} - \boldsymbol{\mu}_0) \\ &= -\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}}) - \frac{1}{2} (\boldsymbol{\mu} - \boldsymbol{\mu}_N)^\top \boldsymbol{\Sigma}_N^{-1} (\boldsymbol{\mu} - \boldsymbol{\mu}_N) \\ & + \frac{1}{2} \boldsymbol{\mu}_N^\top \boldsymbol{\Sigma}_N^{-1} \boldsymbol{\mu}_N, \end{aligned} \quad (2.267)$$

where

$$\begin{aligned}\boldsymbol{\mu}_N &= (N\boldsymbol{\Sigma}_0^{-1} + \boldsymbol{\Sigma}^{-1})^{-1} (N\boldsymbol{\Sigma}_0^{-1}\boldsymbol{\mu}_{\text{ML}} + \boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}_0), \\ \boldsymbol{\Sigma}_N &= (N\boldsymbol{\Sigma}_0^{-1} + \boldsymbol{\Sigma}^{-1})^{-1}.\end{aligned}\tag{2.268}$$

Therefore,

$$p(\boldsymbol{\mu}|\mathbf{X}) = \mathcal{N}\left(\boldsymbol{\mu} \mid (N\boldsymbol{\Sigma}_0^{-1} + \boldsymbol{\Sigma}^{-1})^{-1} (N\boldsymbol{\Sigma}_0^{-1}\boldsymbol{\mu}_{\text{ML}} + \boldsymbol{\Sigma}^{-1}\boldsymbol{\mu}_0), (N\boldsymbol{\Sigma}_0^{-1} + \boldsymbol{\Sigma}^{-1})^{-1}\right).\tag{2.269}$$

2.41

By the definition,

$$\text{Gam}(\lambda|a, b) = \frac{b^a}{\Gamma(a)} \lambda^{a-1} \exp(-b\lambda).\tag{2.270}$$

Then

$$\int_0^\infty \text{Gam}(\lambda|a, b) d\lambda = \frac{b^a}{\Gamma(a)} \int_0^\infty \lambda^{a-1} \exp(-b\lambda) d\lambda.\tag{2.271}$$

By the transformation

$$\lambda' = b\lambda,\tag{2.272}$$

the right hand side can be written as

$$\frac{b^a}{\Gamma(a)} \int_0^\infty \left(\frac{\lambda'}{b}\right)^{a-1} \exp(-\lambda') \frac{1}{b} d\lambda' = \frac{1}{\Gamma(a)} \int_0^\infty \lambda'^{a-1} \exp(-\lambda') d\lambda'.\tag{2.273}$$

The right hand side can be written as

$$\frac{1}{\Gamma(a)} \Gamma(a) = 1.\tag{2.274}$$

Therefore,

$$\int_0^\infty \text{Gam}(\lambda|a, b) d\lambda = 1.\tag{2.275}$$

2.42

Let λ be a variable such that

$$p(\lambda) = \text{Gam}(\lambda|a, b).\tag{2.276}$$

By the definition,

$$\text{Gam}(\lambda|a, b) = \frac{b^a}{\Gamma(a)} \lambda^{a-1} \exp(-b\lambda). \quad (2.277)$$

Then

$$\text{E } \lambda = \frac{b^a}{\Gamma(a)} \int_0^\infty \lambda^a \exp\left(-\frac{\lambda}{b}\right) d\lambda. \quad (2.278)$$

By the transformation

$$\lambda' = b\lambda, \quad (2.279)$$

the right hand side can be written as

$$\frac{b^a}{\Gamma(a)} \int_0^\infty \left(\frac{\lambda'}{b}\right)^a \exp(-\lambda') \frac{1}{b} d\lambda' = \frac{1}{b\Gamma(a)} \int_0^\infty \lambda'^a \exp(-\lambda') d\lambda'. \quad (2.280)$$

The right hand side can be written as

$$\frac{1}{b\Gamma(a)} \Gamma(a+1) = \frac{a}{b}. \quad (2.281)$$

Therefore,

$$\text{E } \lambda = \frac{a}{b}. \quad (2.282)$$

Additionally,

$$\text{E } \lambda^2 = \frac{b^a}{\Gamma(a)} \int_0^\infty \lambda^{a+1} \exp\left(-\frac{\lambda}{b}\right) d\lambda. \quad (2.283)$$

By the transformation

$$\lambda' = b\lambda, \quad (2.284)$$

the right hand side can be written as

$$\frac{b^a}{\Gamma(a)} \int_0^\infty \left(\frac{\lambda'}{b}\right)^{a+1} \exp(-\lambda') \frac{1}{b} d\lambda' = \frac{1}{b^2\Gamma(a)} \int_0^\infty \lambda'^{a+1} \exp(-\lambda') d\lambda'. \quad (2.285)$$

The right hand side can be written as

$$\frac{1}{b^2\Gamma(a)} \Gamma(a+2) = \frac{a(a+1)}{b^2}. \quad (2.286)$$

Therefore,

$$E \lambda^2 = \frac{a(a+1)}{b^2}. \quad (2.287)$$

By the definition,

$$\text{var } \lambda = E \lambda^2 - (E \lambda)^2. \quad (2.288)$$

Therefore,

$$\text{var } \lambda = \frac{a}{b^2}. \quad (2.289)$$

Finally, setting the derivative of $\text{Gam}(\lambda|a, b)$ with respect to λ to zero gives

$$0 = \frac{b^a}{\Gamma(a)} \left(\frac{a-1}{\lambda} - b \right) \lambda^{a-1} \exp \left(-\frac{\lambda}{b} \right). \quad (2.290)$$

Therefore,

$$\text{mode } \lambda = \frac{a-1}{b}. \quad (2.291)$$

2.43

Let

$$p(x|\sigma^2, q) = \frac{q}{2\Gamma(\frac{1}{q})} (2\sigma^2)^{-\frac{1}{q}} \exp \left(-\frac{|x|^q}{2\sigma^2} \right). \quad (2.292)$$

Then

$$\int_{-\infty}^{\infty} p(x|\sigma^2, q) dx = \frac{q}{\Gamma(\frac{1}{q})} (2\sigma^2)^{-\frac{1}{q}} \int_0^{\infty} \exp \left(-\frac{x^q}{2\sigma^2} \right) dx. \quad (2.293)$$

By the transformation

$$x' = \frac{x^q}{2\sigma^2}, \quad (2.294)$$

the right hand side can be written as

$$\begin{aligned} & \frac{q}{\Gamma(\frac{1}{q})} (2\sigma^2)^{-\frac{1}{q}} \int_0^{\infty} \exp(-x') (2\sigma^2)^{\frac{1}{q}} \frac{1}{q} x'^{\frac{1}{q}-1} dx' \\ &= \frac{1}{\Gamma(\frac{1}{q})} \int_0^{\infty} x'^{\frac{1}{q}-1} \exp(-x') dx'. \end{aligned} \quad (2.295)$$

The right hand side can be written as

$$\frac{1}{\Gamma(\frac{1}{q})} \Gamma \left(\frac{1}{q} \right) = 1. \quad (2.296)$$

Therefore,

$$\int_{-\infty}^{\infty} p(x|\sigma^2, q) dx = 1. \quad (2.297)$$

Additionally,

$$p(x|\sigma^2, 2) = \frac{1}{\Gamma(\frac{1}{2})} (2\sigma^2)^{-\frac{1}{2}} \exp\left(-\frac{x^2}{2\sigma^2}\right). \quad (2.298)$$

Therefore,

$$p(x|\sigma^2, 2) = \mathcal{N}(x|0, \sigma^2). \quad (2.299)$$

Finally, let $\mathbf{t} = (t_1, \dots, t_N)^\top$ and $\mathbf{X} = \{x_1, \dots, x_N\}$ such that

$$t_n = y(\mathbf{x}_n, \mathbf{w}) + \epsilon_n, \quad (2.300)$$

where

$$p(\epsilon_n) = p(\epsilon_n|\sigma^2, q). \quad (2.301)$$

Therefore, the logarithm of $p(\epsilon_n)$ except the terms independent of \mathbf{w} and σ^2 can be written as

$$-\frac{|\epsilon_n|^q}{2\sigma^2} - \frac{1}{q} \ln(2\sigma^2). \quad (2.302)$$

Thus, the logarithm of $p(\mathbf{t}|\mathbf{X}, \mathbf{w}, \sigma^2)$ except the terms independent of \mathbf{w} and σ^2 can be written as

$$-\frac{1}{2\sigma^2} \sum_{n=1}^N |y(\mathbf{x}_n, \mathbf{w}) - t_n|^q - \frac{N}{q} \ln(2\sigma^2). \quad (2.303)$$

2.44

Let x_1, \dots, x_N be variables such that

$$\begin{aligned} p(x_n|\mu, \tau) &= \mathcal{N}(x_n|\mu, \tau^{-1}), \\ p(\mu, \tau) &= \mathcal{N}(\mu|\mu_0, (\beta\tau)^{-1}) \text{Gam}(\tau|a, b). \end{aligned} \quad (2.304)$$

By the Bayes' theorem,

$$p(\mu, \tau|\mathbf{x})p(\mathbf{x}) = p(\mathbf{x}|\mu, \tau)p(\mu, \tau). \quad (2.305)$$

The logarithm of the right hand side except the terms independent of \mathbf{x} , μ and τ can be written as

$$\begin{aligned} & \frac{N}{2} \ln \tau - \frac{\tau}{2} \sum_{n=1}^N (x_n - \mu)^2 + \frac{1}{2} \ln \tau - \frac{\beta \tau}{2} (\mu - \mu_0)^2 + (a - 1) \ln \tau - b\tau \\ &= \left(a + \frac{N-1}{2} \right) \ln \tau - \frac{N\tau}{2} (\bar{x} - \mu)^2 - \frac{\beta \tau}{2} (\mu - \mu_0)^2 - b\tau - \frac{\tau}{2} \sum_{n=1}^N (x_n - \bar{x})^2, \end{aligned} \quad (2.306)$$

where

$$\bar{x} = \frac{1}{N} \sum_{n=1}^N x_n. \quad (2.307)$$

Since

$$-\frac{N\tau}{2} (\bar{x} - \mu)^2 - \frac{\beta \tau}{2} (\mu - \mu_0)^2 = -\frac{(N + \beta)\tau}{2} \left(\mu - \frac{N\bar{x} + \beta\mu_0}{N + \beta} \right)^2 - \frac{N\beta\tau(\bar{x} - \mu_0)^2}{2(N + \beta)}, \quad (2.308)$$

the right hand side can be written as

$$\begin{aligned} & -\frac{(N + \beta)\tau}{2} \left(\mu - \frac{N\bar{x} + \beta\mu_0}{N + \beta} \right)^2 \\ & + \left(a + \frac{N-1}{2} \right) \ln \tau - \left(b + \frac{N\beta(\bar{x} - \mu_0)^2}{2(N + \beta)} + \frac{1}{2} \sum_{n=1}^N (x_n - \bar{x})^2 \right) \tau. \end{aligned} \quad (2.309)$$

Therefore,

$$\begin{aligned} p(\mu, \tau | \mathbf{x}) &= \mathcal{N} \left(\mu \mid \frac{N\bar{x} + \beta\mu_0}{N + \beta}, ((N + \beta)\tau)^{-1} \right) \\ & \quad \text{Gam} \left(\tau \mid a + \frac{N+1}{2}, b + \frac{N\beta(\bar{x} - \mu_0)^2}{2(N + \beta)} + \frac{1}{2} \sum_{n=1}^N (x_n - \bar{x})^2 \right). \end{aligned} \quad (2.310)$$

2.45

Let \mathbf{x} be a variable in D dimensions such that

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} | \boldsymbol{\mu}, \boldsymbol{\Lambda}^{-1}). \quad (2.311)$$

Then

$$p(\mathbf{X}|\mathbf{\Lambda}) = \prod_{n=1}^N \mathcal{N}(\mathbf{x}_n|\boldsymbol{\mu}, \mathbf{\Lambda}^{-1}). \quad (2.312)$$

The right hand side except the terms independent of $\mathbf{\Lambda}$ can be written as

$$(\det \mathbf{\Lambda})^{\frac{N}{2}} \exp\left(-\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \mathbf{\Lambda} (\mathbf{x}_n - \boldsymbol{\mu})\right) = (\det \mathbf{\Lambda})^{\frac{N}{2}} \exp\left(-\frac{1}{2} \text{tr}(\mathbf{S}\mathbf{\Lambda})\right), \quad (2.313)$$

where

$$\mathbf{S} = \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})(\mathbf{x}_n - \boldsymbol{\mu})^\top. \quad (2.314)$$

Therefore,

$$p(\mathbf{X}|\mathbf{\Lambda}) \propto (\det \mathbf{\Lambda})^{\frac{N}{2}} \exp\left(-\frac{1}{2} \text{tr}(\mathbf{S}\mathbf{\Lambda})\right). \quad (2.315)$$

Let us assume that a prior distribution of $\mathbf{\Lambda}$ is given by

$$\mathcal{W}(\mathbf{\Lambda}|\mathbf{W}, \nu) = B(\mathbf{W}, \nu) (\det \mathbf{\Lambda})^{\frac{\nu-D-1}{2}} \exp\left(-\frac{1}{2} \text{tr}(\mathbf{W}^{-1}\mathbf{\Lambda})\right). \quad (2.316)$$

Then, by the definition,

$$p(\mathbf{\Lambda}|\mathbf{X}, \mathbf{W}, \nu) \propto p(\mathbf{X}|\mathbf{\Lambda}) \mathcal{W}(\mathbf{\Lambda}|\mathbf{W}, \nu), \quad (2.317)$$

where the right hand side except the terms independent of $\mathbf{\Lambda}$ can be written as

$$(\det \mathbf{\Lambda})^{\frac{\nu+N-D-1}{2}} \exp\left(-\frac{1}{2} \text{tr}((\mathbf{W}^{-1} + \mathbf{S})\mathbf{\Lambda})\right). \quad (2.318)$$

Therefore,

$$p(\mathbf{\Lambda}|\mathbf{X}, \mathbf{W}, \nu) = \mathcal{W}\left(\mathbf{\Lambda} \mid (\mathbf{W}^{-1} + \mathbf{S})^{-1}, \nu + N\right). \quad (2.319)$$

Thus, \mathcal{W} is a conjugate prior distribution of $\mathbf{\Lambda}$.

2.46

Let x be a variable such that

$$p(x|\mu, \tau, a, b) = \mathcal{N}(x|\mu, \tau^{-1}) \text{Gam}(\tau|a, b). \quad (2.320)$$

Then

$$p(x|\mu, a, b) = \int_0^\infty \mathcal{N}(x|\mu, \tau^{-1}) \text{Gam}(\tau|a, b) d\tau. \quad (2.321)$$

The right hand side can be written as

$$\begin{aligned} & \int_0^\infty \left(\frac{\tau}{2\pi}\right)^{\frac{1}{2}} \exp\left(-\frac{\tau}{2}(x-\mu)^2\right) \frac{b^a}{\Gamma(a)} \tau^{a-1} \exp(-b\tau) d\tau \\ &= (2\pi)^{-\frac{1}{2}} \frac{b^a}{\Gamma(a)} \int_0^\infty \tau^{a-\frac{1}{2}} \exp\left(-\left(b + \frac{(x-\mu)^2}{2}\right)\tau\right) d\tau. \end{aligned} \quad (2.322)$$

By the transformation

$$\tau' = \left(b + \frac{(x-\mu)^2}{2}\right) \tau, \quad (2.323)$$

the integral of the right hand side can be written as

$$\int_0^\infty \left(\frac{\tau'}{b + \frac{(x-\mu)^2}{2}}\right)^{a-\frac{1}{2}} \exp(-\tau') \frac{d\tau'}{b + \frac{(x-\mu)^2}{2}} = \Gamma\left(a + \frac{1}{2}\right) \left(b + \frac{(x-\mu)^2}{2}\right)^{-a-\frac{1}{2}}. \quad (2.324)$$

Therefore,

$$p(x|\mu, \tau, a, b) = (2\pi)^{-\frac{1}{2}} \frac{\Gamma(a + \frac{1}{2})}{\Gamma(a)} b^a \left(b + \frac{(x-\mu)^2}{2}\right)^{-a-\frac{1}{2}}. \quad (2.325)$$

Let

$$\begin{aligned} \nu &= 2a, \\ \lambda &= \frac{a}{b}. \end{aligned} \quad (2.326)$$

Then

$$p(x|\mu, \lambda, \nu) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})} \left(\frac{\lambda}{\pi\nu}\right)^{\frac{1}{2}} \left(1 + \frac{\lambda(x-\mu)^2}{\nu}\right)^{-\frac{\nu+1}{2}}. \quad (2.327)$$

2.47

By the definition,

$$\text{St}(x|\mu, \lambda, \nu) = \frac{\Gamma(\frac{\nu+1}{2})}{\Gamma(\frac{\nu}{2})} \left(\frac{\lambda}{\pi\nu}\right)^{\frac{1}{2}} \left(1 + \frac{\lambda(x-\mu)^2}{\nu}\right)^{-\frac{\nu+1}{2}}. \quad (2.328)$$

By the transformation

$$y = \frac{\lambda(x - \mu)^2}{\nu}, \quad (2.329)$$

the right hand side except the terms independent of x can be written as

$$(1 + y)^{-\frac{\lambda(x - \mu)^2}{2y} - \frac{1}{2}}. \quad (2.330)$$

In the limit $y \rightarrow \infty$, it becomes

$$\exp\left(-\frac{\lambda}{2}(x - \mu)^2\right). \quad (2.331)$$

Therefore, in the limit $\nu \rightarrow \infty$, $\text{St}(x|\mu, \lambda, \nu)$ becomes $\mathcal{N}(x|\mu, \lambda^{-1})$.

2.48

Let \mathbf{x} be a variable in D dimensions such that

$$p(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \eta, \nu) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right). \quad (2.332)$$

Then

$$p(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) = \int_0^\infty \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta. \quad (2.333)$$

The right hand side can be written as

$$\begin{aligned} & \int_0^\infty (2\pi)^{-\frac{D}{2}} (\det(\eta\boldsymbol{\Lambda}))^{\frac{1}{2}} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \eta\boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu})\right) \frac{(\frac{\nu}{2})^{\frac{\nu}{2}}}{\Gamma(\frac{\nu}{2})} \eta^{\frac{\nu}{2}-1} \exp\left(-\frac{\nu}{2}\eta\right) d\eta \\ &= (2\pi)^{-\frac{D}{2}} \frac{(\frac{\nu}{2})^{\frac{\nu}{2}}}{\Gamma(\frac{\nu}{2})} (\det \boldsymbol{\Lambda})^{\frac{1}{2}} \int_0^\infty \eta^{\frac{D+\nu}{2}-1} \exp\left(-\frac{1}{2}(\nu + (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu}))\eta\right) d\eta. \end{aligned} \quad (2.334)$$

By the transformation

$$\eta' = \frac{1}{2}(\nu + (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu}))\eta, \quad (2.335)$$

the integral of the right hand side can be written as

$$\begin{aligned} & \int_0^\infty \left(\frac{2\eta'}{\nu + (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu})}\right)^{\frac{D+\nu}{2}-1} \exp(-\eta') \frac{2}{\nu + (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu})} d\eta' \\ &= \left(\frac{2}{\nu + (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu})}\right)^{\frac{D+\nu}{2}} \Gamma\left(\frac{D+\nu}{2}\right). \end{aligned} \quad (2.336)$$

Therefore,

$$p(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) = \frac{\Gamma(\frac{D+\nu}{2})}{\Gamma(\frac{\nu}{2})} \frac{(\det \boldsymbol{\Lambda})^{\frac{1}{2}}}{(\pi\nu)^{\frac{D}{2}}} \left(1 + \frac{(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda} (\mathbf{x} - \boldsymbol{\mu})}{\nu}\right)^{-\frac{D+\nu}{2}}. \quad (2.337)$$

2.49

Let \mathbf{x} be a variable such that

$$p(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) = \text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu). \quad (2.338)$$

By the definition,

$$\text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) = \int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta. \quad (2.339)$$

First,

$$\mathbb{E} \mathbf{x} = \int \mathbf{x} \text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) d\mathbf{x}. \quad (2.340)$$

The right hand side can be written as

$$\begin{aligned} & \int \mathbf{x} \left(\int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta \right) d\mathbf{x} \\ &= \int \left(\int \mathbf{x} \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) d\mathbf{x} \right) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta. \end{aligned} \quad (2.341)$$

The right hand side can be written as

$$\boldsymbol{\mu} \int \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta = \boldsymbol{\mu}. \quad (2.342)$$

Therefore,

$$\mathbb{E} \mathbf{x} = \boldsymbol{\mu}. \quad (2.343)$$

Additionally,

$$\text{cov} \mathbf{x} = \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) d\mathbf{x}. \quad (2.344)$$

The right hand side can be written as

$$\begin{aligned} & \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \left(\int \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta \right) d\mathbf{x} \\ &= \int \left(\int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^\top \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) d\mathbf{x} \right) \text{Gam}\left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2}\right) d\eta. \end{aligned} \quad (2.345)$$

The right hand side can be written as

$$\int (\eta \mathbf{\Lambda})^{-1} \text{Gam} \left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2} \right) d\eta = \mathbf{\Lambda}^{-1} \frac{(\frac{\nu}{2})^{\frac{\nu}{2}}}{\Gamma(\frac{\nu}{2})} \int \eta^{\frac{\nu}{2}-2} \exp \left(-\frac{\nu}{2} \eta \right) d\eta. \quad (2.346)$$

By the transformation

$$\eta' = \frac{\nu}{2} \eta, \quad (2.347)$$

the integral of the right hand side can be written as

$$\int \left(\frac{2}{\nu} \eta' \right)^{-\frac{\nu}{2}-2} \exp(-\eta') \frac{2}{\nu} d\eta' = \left(\frac{2}{\nu} \right)^{\frac{\nu}{2}-1} \Gamma \left(\frac{\nu}{2} - 1 \right). \quad (2.348)$$

Therefore, the right hand side can be written as

$$\mathbf{\Lambda}^{-1} \frac{(\frac{\nu}{2})^{\frac{\nu}{2}}}{\Gamma(\frac{\nu}{2})} \left(\frac{2}{\nu} \right)^{\frac{\nu}{2}-1} \Gamma \left(\frac{\nu}{2} - 1 \right) = \frac{\frac{\nu}{2}}{\frac{\nu}{2} - 1} \mathbf{\Lambda}^{-1}. \quad (2.349)$$

Thus,

$$\text{cov } \mathbf{x} = \frac{\nu}{\nu - 2} \mathbf{\Lambda}^{-1}. \quad (2.350)$$

Finally, setting the derivative of $\text{St}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Lambda}, \nu)$ with respect to \mathbf{x} to zero gives

$$\mathbf{0} = -\frac{1}{2} (\mathbf{\Lambda} + \mathbf{\Lambda}^\top) (\mathbf{x} - \boldsymbol{\mu}) \int \eta \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta \mathbf{\Lambda})^{-1}) \text{Gam} \left(\eta \mid \frac{\nu}{2}, \frac{\nu}{2} \right) d\eta. \quad (2.351)$$

Therefore,

$$\text{mode } \mathbf{x} = \boldsymbol{\mu}. \quad (2.352)$$

2.50

By the definition,

$$\text{St}(\mathbf{x}|\boldsymbol{\mu}, \mathbf{\Lambda}, \nu) = \frac{\Gamma(\frac{D+\nu}{2})}{\Gamma(\frac{\nu}{2})} \frac{(\det \mathbf{\Lambda})^{\frac{1}{2}}}{(\pi\nu)^{\frac{D}{2}}} \left(1 + \frac{(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Lambda} (\mathbf{x} - \boldsymbol{\mu})}{\nu} \right)^{-\frac{D+\nu}{2}}. \quad (2.353)$$

By the transformation

$$y = \frac{(\mathbf{x} - \boldsymbol{\mu})^\top \mathbf{\Lambda} (\mathbf{x} - \boldsymbol{\mu})}{\nu}, \quad (2.354)$$

the right hand side except the terms independent of \mathbf{x} can be written as

$$(1 + y)^{-\frac{(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x}-\boldsymbol{\mu})}{2y} - \frac{D}{2}}. \quad (2.355)$$

In the limit $y \rightarrow \infty$, it becomes

$$\exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Lambda}(\mathbf{x} - \boldsymbol{\mu})\right). \quad (2.356)$$

Therefore, in the limit $\nu \rightarrow \infty$, $\text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu)$ becomes $\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}^{-1})$.

2.51

We have

$$\exp(iA) \exp(-iA) = 1. \quad (2.357)$$

The left hand side can be written as

$$(\cos A + i \sin A)(\cos A - i \sin A) = \cos^2 A + \sin^2 A. \quad (2.358)$$

Therefore,

$$\cos^2 A + \sin^2 A = 1. \quad (2.359)$$

Additionally,

$$\cos(A - B) = \text{Re}(\exp(i(A - B))). \quad (2.360)$$

The right hand side can be written as

$$\text{Re}(\exp(iA) \exp(-iB)) = \text{Re}((\cos A + i \sin A)(\cos B - i \sin B)). \quad (2.361)$$

The right hand side can be written as $\cos A \cos B + \sin A \sin B$. Therefore,

$$\cos(A - B) = \cos A \cos B + \sin A \sin B. \quad (2.362)$$

Finally,

$$\sin(A - B) = \text{Im}(\exp(i(A - B))). \quad (2.363)$$

The right hand side can be written as

$$\text{Im}(\exp(iA) \exp(-iB)) = \text{Im}((\cos A + i \sin A)(\cos B - i \sin B)). \quad (2.364)$$

The right hand side can be written as $\sin A \cos B - \cos A \sin B$. Therefore,

$$\sin(A - B) = \sin A \cos B - \cos A \sin B. \quad (2.365)$$

2.52 (Incomplete)

Let θ be a variable such that

$$p(\theta|\theta_0, m) = \frac{1}{2\pi I_0(m)} \exp(m \cos(\theta - \theta_0)), \quad (2.366)$$

where

$$I_0(m) = \frac{1}{2\pi} \int_0^{2\pi} \exp(m \cos \theta) d\theta. \quad (2.367)$$

By the Taylor series

$$\cos \alpha = 1 - \frac{1}{2}\alpha^2 + O(\alpha^4) \quad (2.368)$$

and the transformation

$$\xi = m^{\frac{1}{2}}(\theta - \theta_0), \quad (2.369)$$

we have

$$\exp(m \cos(\theta - \theta_0)) = \exp\left(m \left(1 - \frac{1}{2}(\theta - \theta_0)^2 + O((\theta - \theta_0)^4)\right)\right). \quad (2.370)$$

2.53

Let θ_0 be a parameter such that

$$\sum_{n=1}^N \sin(\theta_n - \theta_0) = 0. \quad (2.371)$$

The left hand side can be written as

$$\sum_{n=1}^N (\sin \theta_n \cos \theta_0 - \cos \theta_n \sin \theta_0) = \cos \theta_0 \sum_{n=1}^N \sin \theta_n - \sin \theta_0 \sum_{n=1}^N \cos \theta_n. \quad (2.372)$$

Therefore,

$$\theta_0 = \arctan \left(\frac{\sum_{n=1}^N \sin \theta_n}{\sum_{n=1}^N \cos \theta_n} \right). \quad (2.373)$$

2.54

Let θ be a variable such that

$$p(\theta|\theta_0, m) = \frac{1}{2\pi I_0(m)} \exp(m \cos(\theta - \theta_0)), \quad (2.374)$$

where

$$I_0(m) = \frac{1}{2\pi} \int_0^{2\pi} \exp(m \cos \theta) d\theta. \quad (2.375)$$

Setting the first and second derivatives with respect to θ to zero gives

$$\begin{aligned} 0 &= -m \sin(\theta - \theta_0) p(\theta|\theta_0, m), \\ 0 &= (m^2 \sin^2(\theta - \theta_0) - m \cos(\theta - \theta_0)) p(\theta|\theta_0, m). \end{aligned} \quad (2.376)$$

Therefore,

$$\begin{aligned} \operatorname{argmax}_{\theta} p(\theta|\theta_0, m) &= \theta_0, \\ \operatorname{argmin}_{\theta} p(\theta|\theta_0, m) &= \theta_0 - \pi \operatorname{sgn}(\theta_0 - \pi). \end{aligned} \quad (2.377)$$

2.55

Let

$$\theta_0^{\text{ML}} = \arctan \left(\frac{\sum_{n=1}^N \sin \theta_n}{\sum_{n=1}^N \cos \theta_n} \right). \quad (2.378)$$

Let

$$\begin{aligned} \bar{r} \cos \bar{\theta} &= \frac{1}{N} \sum_{n=1}^N \cos \theta_n, \\ \bar{r} \sin \bar{\theta} &= \frac{1}{N} \sum_{n=1}^N \sin \theta_n. \end{aligned} \quad (2.379)$$

Then

$$\theta_0^{\text{ML}} = \bar{\theta}. \quad (2.380)$$

Here,

$$\frac{1}{N} \sum_{n=1}^N \cos(\theta_n - \theta_0^{\text{ML}}) = \left(\frac{1}{N} \sum_{n=1}^N \cos \theta_n \right) \cos \theta_0^{\text{ML}} + \left(\frac{1}{N} \sum_{n=1}^N \sin \theta_n \right) \sin \theta_0^{\text{ML}}. \quad (2.381)$$

By the result above, the right hand side can be written as

$$\bar{r} \cos^2 \bar{\theta} + \bar{r} \sin^2 \bar{\theta} = \bar{r}. \quad (2.382)$$

Therefore,

$$\frac{1}{N} \sum_{n=1}^N \cos(\theta_n - \theta_0^{\text{ML}}) = \bar{r}. \quad (2.383)$$

2.56

By the definition,

$$\text{Beta}(\mu|a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \mu^{a-1} (1-\mu)^{b-1}. \quad (2.384)$$

The right hand side can be written as

$$\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} \exp((a-1) \ln \mu + (b-1) \ln(1-\mu)) \quad (2.385)$$

Therefore, the natural parameters are given by

$$\boldsymbol{\eta} = \begin{bmatrix} a-1 \\ b-1 \end{bmatrix}.$$

Additionally, by the definition,

$$\text{Gam}(\lambda|a, b) = \frac{b^a}{\Gamma(a)} \lambda^{a-1} \exp(-b\lambda). \quad (2.386)$$

The right hand side can be written as

$$\frac{b^a}{\Gamma(a)} \exp((a-1) \ln \lambda - b\lambda). \quad (2.387)$$

Therefore, the natural parameters are given by

$$\boldsymbol{\eta} = \begin{bmatrix} a-1 \\ -b \end{bmatrix}.$$

Finally, for

$$p(\theta|\theta_0, m) = \frac{1}{2\pi I_0(m)} \exp(m \cos(\theta - \theta_0)), \quad (2.388)$$

where

$$I_0(m) = \frac{1}{2\pi} \int_0^{2\pi} \exp(m \cos \theta) d\theta, \quad (2.389)$$

the right hand side can be written as

$$\frac{1}{2\pi I_0(m)} \exp(m \cos \theta_0 \cos \theta + m \sin \theta_0 \sin \theta). \quad (2.390)$$

Therefore, the natural parameters are given by

$$\boldsymbol{\eta} = \begin{bmatrix} m \cos \theta_0 \\ m \sin \theta_0 \end{bmatrix}.$$

2.57

By the definition,

$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-\frac{D}{2}} (\det \boldsymbol{\Sigma})^{-\frac{1}{2}} \exp \left(-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right). \quad (2.391)$$

Therefore,

$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = h(\mathbf{x}) g(\boldsymbol{\eta}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})), \quad (2.392)$$

where

$$\begin{aligned} h(\mathbf{x}) &= (2\pi)^{-\frac{D}{2}}, \\ g(\boldsymbol{\eta}) &= (\det(-2\boldsymbol{\eta}_2))^{-\frac{1}{2}} \exp \left(\frac{1}{4} \boldsymbol{\eta}_1^\top \boldsymbol{\eta}_2^{-1} \boldsymbol{\eta}_1 \right), \\ \boldsymbol{\eta} &= \begin{bmatrix} \boldsymbol{\Sigma}^{-1} \boldsymbol{\mu} \\ -\frac{1}{2} \boldsymbol{\Sigma}^{-1} \end{bmatrix}, \\ \mathbf{u}(\mathbf{x}) &= \begin{bmatrix} \mathbf{x} \\ \mathbf{x} \mathbf{x}^\top \end{bmatrix}. \end{aligned}$$

2.58

Let \mathbf{x} be a variable such that

$$p(\mathbf{x}|\boldsymbol{\eta}) = h(\mathbf{x}) g(\boldsymbol{\eta}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})). \quad (2.393)$$

Then, taking the first derivative of

$$\int p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} = 1 \quad (2.394)$$

with respect to $\boldsymbol{\eta}$ gives

$$\nabla g(\boldsymbol{\eta}) \int h(\mathbf{x}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})) d\mathbf{x} + g(\boldsymbol{\eta}) \int \mathbf{u}(\mathbf{x}) h(\mathbf{x}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})) d\mathbf{x} = \mathbf{0}. \quad (2.395)$$

The left hand side can be written as

$$\frac{\nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} \int p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} + \int \mathbf{u}(\mathbf{x}) p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} = \frac{\nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} + \mathbb{E} \mathbf{u}(\mathbf{x}). \quad (2.396)$$

Therefore,

$$\mathbb{E} \mathbf{u}(\mathbf{x}) = -\frac{\nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})}. \quad (2.397)$$

Thus,

$$\mathbb{E} \mathbf{u}(\mathbf{x}) = -\nabla \ln g(\boldsymbol{\eta}). \quad (2.398)$$

Taking the second derivative with respect to $\boldsymbol{\eta}$ gives

$$\begin{aligned} \nabla \nabla g(\boldsymbol{\eta}) \int h(\mathbf{x}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})) d\mathbf{x} + 2\nabla g(\boldsymbol{\eta}) \int \mathbf{u}(\mathbf{x})^\top h(\mathbf{x}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})) d\mathbf{x} \\ + g(\boldsymbol{\eta}) \int \mathbf{u}(\mathbf{x}) \mathbf{u}(\mathbf{x})^\top h(\mathbf{x}) \exp(\boldsymbol{\eta}^\top \mathbf{u}(\mathbf{x})) d\mathbf{x} = \mathbf{0}. \end{aligned} \quad (2.399)$$

The left hand side can be written as

$$\begin{aligned} \frac{\nabla \nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} \int p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} + \frac{2\nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} \int \mathbf{u}(\mathbf{x})^\top p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} + \int \mathbf{u}(\mathbf{x}) \mathbf{u}(\mathbf{x})^\top p(\mathbf{x}|\boldsymbol{\eta}) d\mathbf{x} \\ = \frac{\nabla \nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} - 2 \mathbb{E} \mathbf{u}(\mathbf{x}) \mathbb{E} \mathbf{u}(\mathbf{x})^\top + \mathbb{E} (\mathbf{u}(\mathbf{x}) \mathbf{u}(\mathbf{x})^\top). \end{aligned} \quad (2.400)$$

Therefore,

$$\mathbb{E} (\mathbf{u}(\mathbf{x}) \mathbf{u}(\mathbf{x})^\top) = -\frac{\nabla \nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} + \frac{2\nabla g(\boldsymbol{\eta})(\nabla g(\boldsymbol{\eta}))^\top}{g^2(\boldsymbol{\eta})}. \quad (2.401)$$

By the definition,

$$\text{cov} \mathbf{u}(\mathbf{x}) = \mathbb{E} (\mathbf{u}(\mathbf{x}) \mathbf{u}(\mathbf{x})^\top) - \mathbb{E} \mathbf{u}(\mathbf{x}) \mathbb{E} \mathbf{u}(\mathbf{x})^\top. \quad (2.402)$$

Thus,

$$\text{cov} \mathbf{u}(\mathbf{x}) = -\frac{\nabla \nabla g(\boldsymbol{\eta})}{g(\boldsymbol{\eta})} + \frac{\nabla g(\boldsymbol{\eta})(\nabla g(\boldsymbol{\eta}))^\top}{g^2(\boldsymbol{\eta})}. \quad (2.403)$$

Hence,

$$\text{cov} \mathbf{u}(\mathbf{x}) = -\nabla \nabla \ln g(\boldsymbol{\eta}). \quad (2.404)$$

2.59

Let

$$p(x|\sigma) = \frac{1}{\sigma} f\left(\frac{x}{\sigma}\right). \quad (2.405)$$

Then

$$\int p(x|\sigma) dx = \frac{1}{\sigma} \int f\left(\frac{x}{\sigma}\right) dx. \quad (2.406)$$

By the transformation

$$x' = \frac{x}{\sigma}, \quad (2.407)$$

the right hand side can be written as

$$\frac{1}{\sigma} \int f(x') \sigma dx' = \int f(x') dx'. \quad (2.408)$$

Therefore, $p(x|\sigma)$ will be normalised if $f(x)$ is normalised.

2.60

Let \mathbf{x} be a variable such that

$$\mathbf{x} \in \mathcal{R}_i \Rightarrow p(\mathbf{x}) = h_i, \quad (2.409)$$

where

$$\int_{\mathcal{R}_i} d\mathbf{x} = \Delta_i. \quad (2.410)$$

Since

$$\int p(\mathbf{x}) d\mathbf{x} = 1, \quad (2.411)$$

we have

$$\sum_i h_i \Delta_i = 1. \quad (2.412)$$

Let N be the total number of observations and n_i be the number of observations which fall in \mathcal{R}_i . Then, the logarithm of the likelihood is given by

$$\ln \left(\prod_i h_i^{n_i} \right) = \sum_i n_i \ln h_i, \quad (2.413)$$

where

$$\sum_i n_i = N. \quad (2.414)$$

Setting the derivatives of

$$\sum_i n_i \ln h_i + \lambda \left(\sum_i h_i \Delta_i - 1 \right) \quad (2.415)$$

with respect to h_i and λ to zero gives

$$\begin{aligned} \frac{n_i}{h_i} + \lambda \Delta_i &= 0, \\ \sum_i h_i \Delta_i - 1 &= 0. \end{aligned} \quad (2.416)$$

Then,

$$\begin{aligned} \lambda &= -N, \\ h_i &= \frac{n_i}{N \Delta_i}. \end{aligned} \quad (2.417)$$

Therefore, the maximum likelihood estimator for the $\{h_i\}$ is $\frac{n_i}{N \Delta_i}$.

2.61 (Incomplete)

Let \mathbf{x} be a variable and $\mathbf{x}_1, \dots, \mathbf{x}_N$ be observations. Let

$$p(\mathbf{x}) = \frac{K}{NV(\mathbf{x})}, \quad (2.418)$$

where

$$V(\mathbf{x}) = \int_{\|\mathbf{x}' - \mathbf{x}\| \leq \|\mathbf{x}_{(K)} - \mathbf{x}\|} d\mathbf{x}', \quad (2.419)$$

K is a constant and $\mathbf{x}_{(K)}$ is the K th nearest observation from the point \mathbf{x} .

3 Linear Models for Regression

3.1

By the definition,

$$\tanh a = \frac{\exp(a) - \exp(-a)}{\exp(a) + \exp(-a)}. \quad (3.1)$$

The right hand side can be written as

$$\frac{1 - \exp(-2a)}{1 + \exp(-2a)} = \frac{2}{1 + \exp(-2a)} - 1. \quad (3.2)$$

Therefore,

$$\tanh a = 2\sigma(2a) - 1, \quad (3.3)$$

where

$$\sigma(a) = \frac{1}{1 + \exp(-a)}. \quad (3.4)$$

Let

$$y(x_n, \mathbf{w}) = w_0 + \sum_{m=1}^M w_m \sigma\left(\frac{x - \mu_j}{s}\right). \quad (3.5)$$

By the result above, the right hand side can be written as

$$w_0 + \sum_{m=1}^M w_m \frac{1 + \tanh\left(\frac{x - \mu_m}{2s}\right)}{2} = w_0 + \frac{1}{2} \sum_{m=1}^M w_m + \frac{1}{2} \sum_{m=1}^M w_m \tanh\left(\frac{x - \mu_m}{2s}\right). \quad (3.6)$$

Therefore, $y(x_n, \mathbf{w})$ is equivalent to

$$y(x_n, \mathbf{u}) = u_0 + \sum_{m=1}^M u_m \tanh\left(\frac{x - \mu_m}{2s}\right), \quad (3.7)$$

where

$$\begin{aligned} u_0 &= w_0 + \frac{1}{2} \sum_{m=1}^M w_m, \\ u_m &= \frac{1}{2} w_m. \end{aligned} \quad (3.8)$$

3.2 (Incomplete)

Let Φ be an $N \times M$ matrix. Then, for any vector \mathbf{v} in N dimensions,

$$\Phi (\Phi^\top \Phi)^{-1} \Phi^\top \mathbf{v} \quad (3.9)$$

is a projection of \mathbf{v} onto the space spanned by the columns of Φ ?

Additionally, for a vector \mathbf{t} in N dimensions,

$$(\Phi^\top \Phi)^{-1} \Phi^\top \mathbf{t} \quad (3.10)$$

is an orthogonal projection of \mathbf{t} onto the space spanned by the columns of Φ ?

3.3

Let

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N r_n (t_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2. \quad (3.11)$$

The right hand side can be written as

$$\frac{1}{2} \|\mathbf{t}' - \Phi' \mathbf{w}\|^2, \quad (3.12)$$

where

$$\mathbf{t}' = \begin{bmatrix} \sqrt{r_1} t_1 \\ \vdots \\ \sqrt{r_N} t_N \end{bmatrix}, \Phi' = \begin{bmatrix} \sqrt{r_1} \phi(\mathbf{x}_1)^\top \\ \vdots \\ \sqrt{r_N} \phi(\mathbf{x}_N)^\top \end{bmatrix}.$$

Setting the derivative with respect to \mathbf{w} to zero gives

$$\mathbf{0} = -\Phi'^\top (\mathbf{t}' - \Phi' \mathbf{w}). \quad (3.13)$$

Therefore,

$$\underset{\mathbf{w}}{\operatorname{argmin}} E(\mathbf{w}) = (\Phi'^\top \Phi')^{-1} \Phi'^\top \mathbf{t}'. \quad (3.14)$$

3.4 (Incomplete)

Let

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N (y(\mathbf{x}_n, \mathbf{w}) - t_n)^2, \quad (3.15)$$

where

$$\begin{aligned} y(\mathbf{x}, \mathbf{w}) &= w_0 + \sum_{m=1}^M w_m (x_m + \epsilon_m), \\ p(\epsilon_m) &= \mathcal{N}(\epsilon_m | 0, \sigma^2). \end{aligned} \quad (3.16)$$

Setting the derivative with respect to \mathbf{w} to zero gives

$$\mathbf{0} = \sum_{n=1}^N \begin{bmatrix} 1 \\ \mathbf{x}_n + \boldsymbol{\epsilon}_n \end{bmatrix} (y(\mathbf{x}_n, \mathbf{w}) - t_n).$$

The right hand side can be written as

$$\sum_{n=1}^N \begin{bmatrix} 1 \\ \mathbf{x}_n \end{bmatrix} (y(\mathbf{x}_n, \mathbf{w}) - t_n) + \sum_{n=1}^N \begin{bmatrix} 0 \\ \boldsymbol{\epsilon}_n \end{bmatrix} (y(\mathbf{x}_n, \mathbf{w}) - t_n).$$

3.5

Let

$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n))^2. \quad (3.17)$$

Then, the minimisation of $E(\mathbf{w})$ under the constraint

$$\sum_{m=1}^M |w_m|^q \leq \eta \quad (3.18)$$

reduces to the minimisation of

$$E(\mathbf{w}) + \lambda \left(\sum_{m=1}^M |w_m|^q - \eta \right) \quad (3.19)$$

with respect to \mathbf{w} and λ . Then,

$$\eta = \sum_{m=1}^M |w_m^*(\lambda)|^q, \quad (3.20)$$

where

$$\mathbf{w}^*(\lambda) = \underset{\mathbf{w}}{\operatorname{argmin}} \left(E(\mathbf{w}) + \lambda \left(\sum_{m=1}^M |w_m|^q - \eta \right) \right). \quad (3.21)$$

3.6

Let $\mathbf{t}_1, \dots, \mathbf{t}_N$ be variables in D dimensions such that

$$p(\mathbf{t}_n | \mathbf{W}, \Sigma) = \mathcal{N}(\mathbf{t}_n | \mathbf{y}(\mathbf{x}_n, \mathbf{W}), \Sigma), \quad (3.22)$$

where

$$\mathbf{y}(\mathbf{x}_n, \mathbf{W}) = \mathbf{W}^\top \phi(\mathbf{x}_n). \quad (3.23)$$

Then,

$$\begin{aligned} & \ln \left(\prod_{n=1}^N p(\mathbf{t}_n | \mathbf{W}, \Sigma) \right) \\ &= -\frac{ND}{2} \ln(2\pi) - \frac{N}{2} \ln(\det \Sigma) - \frac{1}{2} \sum_{n=1}^N (\mathbf{t}_n - \mathbf{W}^\top \phi(\mathbf{x}_n))^\top \Sigma^{-1} (\mathbf{t}_n - \mathbf{W}^\top \phi(\mathbf{x}_n)). \end{aligned} \quad (3.24)$$

By 3.21(a), setting the derivatives with respect to \mathbf{W} and Σ to zero gives

$$\begin{aligned} \mathbf{O} &= -\frac{1}{2} (\Sigma^{-1} + (\Sigma^{-1})^\top) \sum_{n=1}^N (\mathbf{t}_n - \mathbf{W}^\top \phi(\mathbf{x}_n)) (\phi(\mathbf{x}_n))^\top, \\ \mathbf{O} &= -\frac{N}{2} (\Sigma^{-1})^\top + \frac{1}{2} (\Sigma^{-1})^2 \sum_{n=1}^N (\mathbf{t}_n - \mathbf{W}^\top \phi(\mathbf{x}_n)) (\mathbf{t}_n - \mathbf{W}^\top \phi(\mathbf{x}_n))^\top. \end{aligned} \quad (3.25)$$

Therefore,

$$\begin{aligned} \mathbf{W}_{\text{ML}} &= (\Phi^\top \Phi)^{-1} \Phi^\top \mathbf{t}, \\ \Sigma_{\text{ML}} &= \frac{1}{N} \sum_{n=1}^N (\mathbf{t}_n - \mathbf{W}_{\text{ML}}^\top \phi(\mathbf{x}_n)) (\mathbf{t}_n - \mathbf{W}_{\text{ML}}^\top \phi(\mathbf{x}_n))^\top, \end{aligned} \quad (3.26)$$

where

$$\Phi = \begin{bmatrix} \phi(\mathbf{x}_1)^\top \\ \vdots \\ \phi(\mathbf{x}_N)^\top \end{bmatrix}.$$

3.7

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{m}_0, \mathbf{S}_0). \end{aligned} \quad (3.27)$$

By the Bayes' theorem,

$$p(\mathbf{w}|\mathbf{t})p(\mathbf{t}) = p(\mathbf{t}|\mathbf{w})p(\mathbf{w}). \quad (3.28)$$

The logarithm of the right hand side except the terms independent of \mathbf{t} and \mathbf{w} can be written as

$$\begin{aligned} & -\frac{\beta}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n))^2 - \frac{1}{2}(\mathbf{w} - \mathbf{m}_0)^\top \mathbf{S}_0^{-1}(\mathbf{w} - \mathbf{m}_0) \\ &= -\frac{\beta}{2}(\mathbf{t} - \boldsymbol{\Phi}\mathbf{w})^\top (\mathbf{t} - \boldsymbol{\Phi}\mathbf{w}) - \frac{1}{2}(\mathbf{w} - \mathbf{m}_0)^\top \mathbf{S}_0^{-1}(\mathbf{w} - \mathbf{m}_0), \end{aligned} \quad (3.29)$$

where

$$\boldsymbol{\Phi} = \begin{bmatrix} \boldsymbol{\phi}(\mathbf{x}_1)^\top \\ \vdots \\ \boldsymbol{\phi}(\mathbf{x}_N)^\top \end{bmatrix}.$$

The right hand side can be written as

$$-\frac{1}{2}(\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1}(\mathbf{w} - \mathbf{m}_N) + \frac{1}{2}\mathbf{m}_N^\top \mathbf{S}_N^{-1}\mathbf{m}_N - \frac{\beta}{2}\mathbf{t}^\top \mathbf{t} - \frac{1}{2}\mathbf{m}_0^\top \mathbf{S}_0^{-1}\mathbf{m}_0, \quad (3.30)$$

where

$$\begin{aligned} \mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1}\mathbf{m}_0 + \beta\boldsymbol{\Phi}^\top \mathbf{t}), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \beta\boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.31)$$

Therefore,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N). \quad (3.32)$$

3.8

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{m}_0, \mathbf{S}_0). \end{aligned} \quad (3.33)$$

Then, by 3.7,

$$p(\mathbf{w}|\mathbf{t}_N) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.34)$$

where

$$\begin{aligned} \mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \beta \Phi_N^\top \mathbf{t}_N), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \beta \Phi_N^\top \Phi_N. \end{aligned} \quad (3.35)$$

By the Bayes' theorem,

$$p(\mathbf{w}|\mathbf{t}_{N+1})p(\mathbf{t}_{N+1}) = p(\mathbf{t}_{N+1}|\mathbf{w})p(\mathbf{w}). \quad (3.36)$$

The right hand side can be written as

$$p(t_{N+1}|\mathbf{w})p(\mathbf{t}_N|\mathbf{w})p(\mathbf{w}) = p(t_{N+1}|\mathbf{w})p(\mathbf{w}|\mathbf{t}_N)p(\mathbf{t}_N). \quad (3.37)$$

Therefore,

$$p(\mathbf{w}|\mathbf{t}_{N+1})p(t_{N+1}) = p(t_{N+1}|\mathbf{w})p(\mathbf{w}|\mathbf{t}_N). \quad (3.38)$$

The logarithm of the right hand side except the terms independent of \mathbf{w} can be written as

$$\begin{aligned} & -\frac{\beta}{2} (t_{N+1} - \mathbf{w}^\top \phi(\mathbf{x}_{N+1}))^2 - \frac{1}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N) \\ &= -\frac{1}{2} (\mathbf{w} - \boldsymbol{\mu}_{N+1})^\top \boldsymbol{\Lambda}_{N+1} (\mathbf{w} - \boldsymbol{\mu}_{N+1}) + \frac{1}{2} \boldsymbol{\mu}_{N+1}^\top \boldsymbol{\Lambda}_{N+1} \boldsymbol{\mu}_{N+1} \\ & \quad - \frac{1}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N - \frac{\beta}{2} t_{N+1}^2, \end{aligned} \quad (3.39)$$

where

$$\begin{aligned} \boldsymbol{\mu}_{N+1} &= \boldsymbol{\Lambda}_{N+1}^{-1} (\mathbf{S}_N^{-1} \mathbf{m}_N + \beta t_{N+1} \phi(\mathbf{x}_{N+1})), \\ \boldsymbol{\Lambda}_{N+1} &= \mathbf{S}_N^{-1} + \beta \phi(\mathbf{x}_{N+1}) \phi(\mathbf{x}_{N+1})^\top. \end{aligned} \quad (3.40)$$

Therefore,

$$\begin{aligned} \boldsymbol{\mu}_{N+1} &= \mathbf{m}_{N+1}, \\ \boldsymbol{\Lambda}_{N+1} &= \mathbf{S}_{N+1}^{-1}. \end{aligned} \quad (3.41)$$

Thus,

$$p(\mathbf{w}|\mathbf{t}_{N+1}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_{N+1}, \mathbf{S}_{N+1}). \quad (3.42)$$

3.9 (Incomplete)

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{m}_0, \mathbf{S}_0). \end{aligned} \quad (3.43)$$

Then, by 3.7,

$$p(\mathbf{w}|\mathbf{t}_N) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.44)$$

where

$$\begin{aligned} \mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \beta \boldsymbol{\Phi}_N^\top \mathbf{t}_N), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \beta \boldsymbol{\Phi}_N^\top \boldsymbol{\Phi}_N. \end{aligned} \quad (3.45)$$

By the Bayes' theorem,

$$p(\mathbf{w}|\mathbf{t}_{N+1})p(\mathbf{t}_{N+1}) = p(\mathbf{t}_{N+1}|\mathbf{w})p(\mathbf{w}). \quad (3.46)$$

The right hand side can be written as

$$p(t_{N+1}|\mathbf{w})p(\mathbf{t}_N|\mathbf{w})p(\mathbf{w}) = p(t_{N+1}|\mathbf{w})p(\mathbf{w}|\mathbf{t}_N)p(\mathbf{t}_N). \quad (3.47)$$

Therefore,

$$p(\mathbf{w}|\mathbf{t}_{N+1})p(t_{N+1}) = p(t_{N+1}|\mathbf{w})p(\mathbf{w}|\mathbf{t}_N). \quad (3.48)$$

The logarithm of the right hand side except the terms independent of \mathbf{w} can be written as

$$-\frac{\beta}{2} (t_{N+1} - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_{N+1}))^2 - \frac{1}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N). \quad (3.49)$$

3.10

Let t be a variable such that

$$\begin{aligned} p(t|\mathbf{w}) &= \mathcal{N}(t|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{m}_0, \mathbf{S}_0). \end{aligned} \quad (3.50)$$

Then, by 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.51)$$

where

$$\begin{aligned} \mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \beta \boldsymbol{\Phi}^\top \mathbf{t}), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.52)$$

By marginalisation,

$$p(t|\mathbf{t}) = \int p(t|\mathbf{w})p(\mathbf{w}|\mathbf{t})d\mathbf{w}. \quad (3.53)$$

The logarithm of the integrand of the right hand side except the terms independent of t and \mathbf{w} can be written as

$$-\frac{\beta}{2}(t - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}))^2 - \frac{1}{2}(\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1}(\mathbf{w} - \mathbf{m}_N). \quad (3.54)$$

It can be written as

$$-\frac{1}{2} \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix}^\top \begin{bmatrix} \mathbf{S}_N^{-1} + \beta \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\beta \boldsymbol{\phi}(\mathbf{x}) \\ -\beta \boldsymbol{\phi}(\mathbf{x})^\top & \beta \end{bmatrix} \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix} + \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix}^\top \begin{bmatrix} \mathbf{S}_N^{-1} \mathbf{m}_N \\ 0 \end{bmatrix} - \frac{1}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N.$$

By 2.24,

$$\begin{bmatrix} \mathbf{S}_N^{-1} + \beta \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\beta \boldsymbol{\phi}(\mathbf{x}) \\ -\beta \boldsymbol{\phi}(\mathbf{x})^\top & \beta \end{bmatrix}^{-1} = \begin{bmatrix} \mathbf{S}_N & \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}) \\ \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N & \frac{1}{\beta} + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}) \end{bmatrix}.$$

Therefore,

$$\begin{bmatrix} \mathbf{S}_N^{-1} + \beta \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\beta \boldsymbol{\phi}(\mathbf{x}) \\ -\beta \boldsymbol{\phi}(\mathbf{x})^\top & \beta \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{S}_N^{-1} \mathbf{m}_N \\ 0 \end{bmatrix} = \begin{bmatrix} \mathbf{m}_N \\ \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}) \end{bmatrix}.$$

Thus,

$$p(t|\mathbf{t}) = \mathcal{N}(t|\mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}), \sigma_N^2(\mathbf{x})), \quad (3.55)$$

where

$$\sigma_N^2(\mathbf{x}) = \frac{1}{\beta} + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}). \quad (3.56)$$

3.11

Let t be a variable such that

$$\begin{aligned} p(t|\mathbf{w}) &= \mathcal{N}(t|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{m}_0, \mathbf{S}_0). \end{aligned} \quad (3.57)$$

Then, by 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.58)$$

where

$$\begin{aligned}\mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \beta \Phi_N^\top \mathbf{t}_N), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \beta \Phi_N^\top \Phi_N.\end{aligned}\tag{3.59}$$

Then, by 3.10,

$$p(t|\mathbf{t}) = \mathcal{N}(t \mid \mathbf{m}_N^\top \phi(\mathbf{x}), \sigma_N^2(\mathbf{x})),\tag{3.60}$$

where

$$\sigma_N^2(\mathbf{x}) = \frac{1}{\beta} + \phi(\mathbf{x})^\top \mathbf{S}_N \phi(\mathbf{x}).\tag{3.61}$$

Then,

$$\sigma_N^2(\mathbf{x}) - \sigma_{N+1}^2(\mathbf{x}) = \phi(\mathbf{x})^\top (\mathbf{S}_N - \mathbf{S}_{N+1}) \phi(\mathbf{x}).\tag{3.62}$$

By the expression of \mathbf{S}_N above,

$$\mathbf{S}_{N+1} = (\mathbf{S}_N^{-1} + \beta \phi(\mathbf{x}_{N+1}) \phi(\mathbf{x}_{N+1})^\top)^{-1}.\tag{3.63}$$

By the identity

$$(\mathbf{M} + \mathbf{v} \mathbf{v}^\top)^{-1} = \mathbf{M}^{-1} - \frac{(\mathbf{M}^{-1} \mathbf{v}) (\mathbf{v}^\top \mathbf{M}^{-1})}{1 + \mathbf{v}^\top \mathbf{M}^{-1} \mathbf{v}},\tag{3.64}$$

the right hand side can be written as

$$\mathbf{S}_N - \frac{\beta (\mathbf{S}_N \phi(\mathbf{x}_{N+1})) (\phi(\mathbf{x}_{N+1})^\top \mathbf{S}_N)}{1 + \beta \phi(\mathbf{x}_{N+1})^\top \mathbf{S}_N \phi(\mathbf{x}_{N+1})}.\tag{3.65}$$

Therefore,

$$\phi(\mathbf{x})^\top (\mathbf{S}_N - \mathbf{S}_{N+1}) \phi(\mathbf{x}) = \frac{\beta (\phi(\mathbf{x})^\top \mathbf{S}_N \phi(\mathbf{x}_{N+1}))^2}{1 + \beta \phi(\mathbf{x}_{N+1})^\top \mathbf{S}_N \phi(\mathbf{x}_{N+1})}.\tag{3.66}$$

Thus,

$$\sigma_{N+1}^2(\mathbf{x}) \leq \sigma_N^2(\mathbf{x}).\tag{3.67}$$

3.12

Let t_1, \dots, t_N be variables such that

$$\begin{aligned}p(t_n | \mathbf{w}, \beta) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}, \beta) &= \mathcal{N}(\mathbf{w} | \mathbf{m}_0, \beta^{-1} \mathbf{S}_0) \text{Gam}(\beta | a_0, b_0),\end{aligned}\tag{3.68}$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By the Bayes' theorem,

$$p(\mathbf{w}, \beta | \mathbf{t}) p(\mathbf{t}) = p(\mathbf{t} | \mathbf{w}, \beta) p(\mathbf{w}, \beta). \quad (3.69)$$

The logarithm of the right hand side except the terms independent of \mathbf{t} , \mathbf{w} and β can be written as

$$\begin{aligned} & -\frac{N}{2} \ln \beta^{-1} - \frac{\beta}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n))^2 - \frac{M}{2} \ln \beta^{-1} - \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_0)^\top \mathbf{S}_0^{-1} (\mathbf{w} - \mathbf{m}_0) \\ & + (a_0 - 1) \ln \beta - b_0 \beta \\ = & -\frac{M}{2} \ln \beta - \frac{\beta}{2} \mathbf{w}^\top (\mathbf{S}_0^{-1} + \boldsymbol{\Phi}^\top \boldsymbol{\Phi}) \mathbf{w} + \beta \mathbf{w}^\top (\mathbf{S}_0^{-1} \mathbf{m}_0 + \boldsymbol{\Phi}^\top \mathbf{t}) - \frac{\beta}{2} \|\mathbf{t}\|^2 - \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0 \\ & + \left(a_0 + \frac{N}{2} - 1 \right) \ln \beta - b_0 \beta. \end{aligned} \quad (3.70)$$

The right hand side can be written as

$$-\frac{M}{2} \ln \beta - \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N) + (a_N - 1) \ln \beta - b_N \beta, \quad (3.71)$$

where

$$\begin{aligned} \mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \boldsymbol{\Phi}^\top \mathbf{t}), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \boldsymbol{\Phi}^\top \boldsymbol{\Phi}, \\ a_N &= a_0 + \frac{N}{2}, \\ b_N &= b_0 + \frac{1}{2} \|\mathbf{t}\|^2 + \frac{1}{2} \mathbf{m}_0^\top \mathbf{S}_0 \mathbf{m}_0 - \frac{1}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N. \end{aligned} \quad (3.72)$$

Therefore,

$$p(\mathbf{w}, \beta | \mathbf{t}) = \mathcal{N}(\mathbf{w} | \mathbf{m}_N, \beta^{-1} \mathbf{S}_N) \text{Gam}(\beta | a_N, b_N). \quad (3.73)$$

Substituting it to the result of the Bayes' theorem above, we have

$$p(\mathbf{t}) = \frac{\mathcal{N}(\mathbf{t} | \boldsymbol{\Phi} \mathbf{w}, \beta^{-1} \mathbf{I}) \mathcal{N}(\mathbf{w} | \mathbf{m}_0, \beta^{-1} \mathbf{S}_0) \text{Gam}(\beta | a_0, b_0)}{\mathcal{N}(\mathbf{w} | \mathbf{m}_N, \beta^{-1} \mathbf{S}_N) \text{Gam}(\beta | a_N, b_N)}. \quad (3.74)$$

The logarithm of the right hand side can be written as

$$\begin{aligned}
& -\frac{N}{2} \ln(2\pi) - \frac{N}{2} \ln \beta^{-1} - \frac{\beta}{2} (\mathbf{t} - \Phi \mathbf{w})^\top (\mathbf{t} - \Phi \mathbf{w}) \\
& -\frac{M}{2} \ln(2\pi) - \frac{M}{2} \ln \beta^{-1} - \frac{1}{2} \det \mathbf{S}_0 - \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_0)^\top \mathbf{S}_0^{-1} (\mathbf{w} - \mathbf{m}_0) \\
& + a_0 \ln b_0 - \ln \Gamma(a_0) + (a_0 - 1) \ln \beta - b_0 \beta \\
& + \frac{M}{2} \ln(2\pi) + \frac{M}{2} \ln \beta^{-1} + \frac{1}{2} \det \mathbf{S}_N + \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N) \\
& - a_N \ln b_N + \ln \Gamma(a_N) - (a_N - 1) \ln \beta + b_N \beta \\
& = -\frac{N}{2} \ln(2\pi) - \frac{1}{2} \det \mathbf{S}_0 + a_0 \ln b_0 - \ln \Gamma(a_0) + \frac{1}{2} \det \mathbf{S}_N - a_N \ln b_N + \ln \Gamma(a_N).
\end{aligned} \tag{3.75}$$

Therefore,

$$p(\mathbf{t}) = (2\pi)^{-\frac{N}{2}} \left(\frac{\det \mathbf{S}_N}{\det \mathbf{S}_0} \right)^{\frac{1}{2}} \frac{\Gamma(a_N)}{\Gamma(a_0)} \frac{b_0^{a_0}}{b_N^{a_N}}. \tag{3.76}$$

3.13

Let t_1, \dots, t_N be variables such that

$$\begin{aligned}
p(t_n | \mathbf{w}, \beta) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\
p(\mathbf{w}, \beta) &= \mathcal{N}(\mathbf{w} | \mathbf{m}_0, \beta^{-1} \mathbf{S}_0) \text{Gam}(\beta | a_0, b_0),
\end{aligned} \tag{3.77}$$

where \mathbf{w} and ϕ are vectors in M dimensions. Then, by 3.12,

$$p(\mathbf{w}, \beta | \mathbf{t}) = \mathcal{N}(\mathbf{w} | \mathbf{m}_N, \beta^{-1} \mathbf{S}_N) \text{Gam}(\beta | a_N, b_N), \tag{3.78}$$

where

$$\begin{aligned}
\mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \Phi^\top \mathbf{t}), \\
\mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \Phi^\top \Phi, \\
a_N &= a_0 + \frac{N}{2}, \\
b_N &= b_0 + \frac{1}{2} \|\mathbf{t}\|^2 + \frac{1}{2} \mathbf{m}_0^\top \mathbf{S}_0 \mathbf{m}_0 - \frac{1}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N.
\end{aligned} \tag{3.79}$$

By marginalisation,

$$p(t | \mathbf{t}) = \int \int p(t | \mathbf{w}, \beta) p(\mathbf{w}, \beta | \mathbf{t}) d\mathbf{w} d\beta. \tag{3.80}$$

The right hand side can be written as

$$\int \left(\int \mathcal{N}(t|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}), \beta^{-1}) \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \beta^{-1}\mathbf{S}_N) d\mathbf{w} \right) \text{Gam}(\beta|a_N, b_N) d\beta. \quad (3.81)$$

The logarithm of the integrand with respect to \mathbf{w} except the terms independent of \mathbf{w} can be written as

$$-\frac{\beta}{2} (t - \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}))^2 - \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N). \quad (3.82)$$

It can be written as

$$-\frac{\beta}{2} \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix}^\top \begin{bmatrix} \mathbf{S}_N^{-1} + \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\boldsymbol{\phi}(\mathbf{x}) \\ -\boldsymbol{\phi}(\mathbf{x})^\top & 1 \end{bmatrix} \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix} + \beta \begin{bmatrix} \mathbf{w} \\ t \end{bmatrix}^\top \begin{bmatrix} \mathbf{S}_N^{-1} \mathbf{m}_N \\ 0 \end{bmatrix} - \frac{\beta}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N.$$

By 2.24,

$$\begin{bmatrix} \mathbf{S}_N^{-1} + \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\boldsymbol{\phi}(\mathbf{x}) \\ -\boldsymbol{\phi}(\mathbf{x})^\top & 1 \end{bmatrix}^{-1} = \begin{bmatrix} \mathbf{S}_N & \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}) \\ \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N & 1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}) \end{bmatrix}.$$

Then,

$$\begin{bmatrix} \mathbf{S}_N^{-1} + \boldsymbol{\phi}(\mathbf{x})\boldsymbol{\phi}(\mathbf{x})^\top & -\boldsymbol{\phi}(\mathbf{x}) \\ -\boldsymbol{\phi}(\mathbf{x})^\top & 1 \end{bmatrix}^{-1} \begin{bmatrix} \mathbf{S}_N^{-1} \mathbf{m}_N \\ 0 \end{bmatrix} = \begin{bmatrix} \mathbf{m}_N \\ \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}) \end{bmatrix}.$$

Therefore, the integral with respect to \mathbf{w} can be written as

$$\mathcal{N}(t|\mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}), \beta^{-1} (1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}))). \quad (3.83)$$

Then, the logarithm of the integrand with respect to β except the terms independent of β can be written as

$$\begin{aligned} & -\frac{1}{2} \ln \beta^{-1} - \frac{\beta}{2(1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}))} (t - \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}))^2 + (a_N - 1) \ln \beta - b_N \beta \\ & = \left(a_N + \frac{1}{2} - 1 \right) \ln \beta - \left(b_N + \frac{(t - \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}))^2}{2(1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}))} \right) \beta. \end{aligned} \quad (3.84)$$

Therefore, the integral with respect to β except the terms independent of t can be written as

$$\left(b_N + \frac{(t - \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}))^2}{2(1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}))} \right)^{-a_N - \frac{1}{2}}. \quad (3.85)$$

Thus,

$$p(t|\mathbf{x}, \mathbf{t}) = \text{St}(t|\mu, \lambda, \nu), \quad (3.86)$$

where

$$\begin{aligned} \mu &= \mathbf{m}_N^\top \boldsymbol{\phi}(\mathbf{x}), \\ \lambda &= \frac{a_N}{b_N} (1 + \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}))^{-1}, \\ \nu &= 2a_N. \end{aligned} \quad (3.87)$$

3.14 (Incomplete)

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}), \end{aligned} \quad (3.88)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. Then, by 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.89)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.90)$$

Let

$$y(\mathbf{x}, \mathbf{w}) = \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}). \quad (3.91)$$

Then,

$$y(\mathbf{x}, \mathbf{m}_N) = \sum_{n=1}^N k(\mathbf{x}, \mathbf{x}_n) t_n, \quad (3.92)$$

where

$$k(\mathbf{x}, \mathbf{x}') = \beta \boldsymbol{\phi}(\mathbf{x})^\top \mathbf{S}_N \boldsymbol{\phi}(\mathbf{x}'). \quad (3.93)$$

Let us suppose that $\boldsymbol{\phi}_j(\mathbf{x})$ are linearly independent, $N > M$ and

$$\phi_0(\mathbf{x}) = 1. \quad (3.94)$$

Then, we can construct a new basis set $\psi_j(\mathbf{x})$ such that

$$\boldsymbol{\Psi}^\top \boldsymbol{\Psi} = \mathbf{I}? \quad (3.95)$$

$$\sum_{n=1}^N \psi_j(\mathbf{x}_n) \psi_k(\mathbf{x}_n) = I_{jk} \quad (3.96)$$

where

$$\mathbf{\Psi} = \begin{bmatrix} \boldsymbol{\psi}(\mathbf{x}_1)^\top \\ \vdots \\ \boldsymbol{\psi}(\mathbf{x}_N)^\top \end{bmatrix}$$

and

$$\psi_0(\mathbf{x}) = 1. \quad (3.97)$$

Under the basis set, if $\alpha = 0$, then

$$\mathbf{S}_N^{-1} = \beta \mathbf{I}, \quad (3.98)$$

so that

$$k(\mathbf{x}, \mathbf{x}') = \boldsymbol{\psi}(\mathbf{x})^\top \boldsymbol{\psi}(\mathbf{x}'). \quad (3.99)$$

Then,

$$\sum_{n=1}^N k(\mathbf{x}, \mathbf{x}_n) = \sum_{n=1}^N \sum_{j=0}^{M-1} \psi_j(\mathbf{x}) \psi_j(\mathbf{x}_n) = 1? \quad (3.100)$$

3.15

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n | \mathbf{w}) &= \mathcal{N}(t_n | \mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w} | \mathbf{0}, \alpha^{-1} \mathbf{I}), \end{aligned} \quad (3.101)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w} | \mathbf{t}) = \mathcal{N}(\mathbf{w} | \mathbf{m}_N, \mathbf{S}_N), \quad (3.102)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.103)$$

By 3.19,

$$\ln p(\mathbf{t}) = \frac{M}{2} \ln \alpha + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi) + \frac{1}{2} \ln(\det \mathbf{S}_N) - E(\mathbf{m}_N), \quad (3.104)$$

where

$$E(\mathbf{m}_N) = \frac{\beta}{2} \|\mathbf{t} - \Phi \mathbf{m}_N\|^2 + \frac{\alpha}{2} \mathbf{m}_N^\top \mathbf{m}_N. \quad (3.105)$$

By 3.22, setting the derivatives of $\ln p(\mathbf{t})$ with respect to α and β to zero gives

$$\begin{aligned} \alpha &= \frac{\gamma}{\mathbf{m}_N^\top \mathbf{m}_N}, \\ \beta &= \frac{N - \gamma}{\|\mathbf{t} - \Phi \mathbf{m}_N\|^2}, \end{aligned} \quad (3.106)$$

where

$$\gamma = \sum_{m=1}^M \frac{\lambda_m}{\alpha + \lambda_m} \quad (3.107)$$

and $\lambda_1, \dots, \lambda_M$ are the eigenvalues of $\beta \Phi^\top \Phi$. If α and β are set as above, then

$$E(\mathbf{m}_N) = \frac{N}{2}. \quad (3.108)$$

3.16

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n | \mathbf{w}) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w} | \mathbf{0}, \alpha^{-1} \mathbf{I}), \end{aligned} \quad (3.109)$$

where \mathbf{w} and ϕ are vectors in M dimensions. By the Bayes' theorem,

$$p(\mathbf{t}, \mathbf{w}) = p(\mathbf{t} | \mathbf{w}) p(\mathbf{w}). \quad (3.110)$$

Integrating both sides with respect to \mathbf{w} gives

$$p(\mathbf{t}) = \int p(\mathbf{t} | \mathbf{w}) p(\mathbf{w}) d\mathbf{w}. \quad (3.111)$$

The logarithm of the integrand of the right hand side except the terms independent of \mathbf{w} can be written as

$$-\frac{\beta}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2 - \frac{\alpha}{2} \mathbf{w}^\top \mathbf{w} = -\frac{1}{2} \begin{bmatrix} \mathbf{w} \\ \mathbf{t} \end{bmatrix}^\top \begin{bmatrix} \alpha \mathbf{I} + \beta \Phi^\top \Phi & -\beta \Phi^\top \\ -\beta \Phi & \beta \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{w} \\ \mathbf{t} \end{bmatrix}.$$

By 2.24,

$$\begin{bmatrix} \alpha \mathbf{I} + \beta \Phi^\top \Phi & -\beta \Phi^\top \\ -\beta \Phi & \beta \mathbf{I} \end{bmatrix}^{-1} = \begin{bmatrix} \alpha^{-1} \mathbf{I} & \alpha^{-1} \Phi^\top \\ \alpha^{-1} \Phi & \alpha^{-1} \Phi \Phi^\top + \beta^{-1} \mathbf{I} \end{bmatrix}.$$

Therefore,

$$p(\mathbf{t}) = \mathcal{N}(\mathbf{t} | \mathbf{0}, \alpha^{-1} \Phi \Phi^\top + \beta^{-1} \mathbf{I}). \quad (3.112)$$

3.17

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n | \mathbf{w}) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w} | \mathbf{0}, \alpha^{-1} \mathbf{I}), \end{aligned} \quad (3.113)$$

where \mathbf{w} and ϕ are vectors in M dimensions. By the Bayes' theorem,

$$p(\mathbf{t}, \mathbf{w}) = p(\mathbf{t} | \mathbf{w}) p(\mathbf{w}). \quad (3.114)$$

Then,

$$p(\mathbf{t}) = \int p(\mathbf{t} | \mathbf{w}) p(\mathbf{w}) d\mathbf{w}. \quad (3.115)$$

The logarithm of the integrand of the right hand side can be written as

$$-\frac{N}{2} \ln(2\pi\beta^{-1}) - \frac{\beta}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2 - \frac{M}{2} \ln(2\pi) - \frac{1}{2} \ln(\det(\alpha^{-1} \mathbf{I})) - \frac{\alpha}{2} \mathbf{w}^\top \mathbf{w}. \quad (3.116)$$

Therefore,

$$p(\mathbf{t}) = \left(\frac{\beta}{2\pi}\right)^{\frac{N}{2}} \left(\frac{\alpha}{2\pi}\right)^{\frac{M}{2}} \int \exp(-E(\mathbf{w})) d\mathbf{w}, \quad (3.117)$$

where

$$E(\mathbf{w}) = \frac{\beta}{2} \|\mathbf{t} - \Phi \mathbf{w}\|^2 + \frac{\alpha}{2} \mathbf{w}^\top \mathbf{w}. \quad (3.118)$$

3.18

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n | \mathbf{w}) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w} | \mathbf{0}, \alpha^{-1} \mathbf{I}), \end{aligned} \quad (3.119)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.120)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.121)$$

By 3.17,

$$p(\mathbf{t}) = \left(\frac{\beta}{2\pi} \right)^{\frac{N}{2}} \left(\frac{\alpha}{2\pi} \right)^{\frac{M}{2}} \int \exp(-E(\mathbf{w})) d\mathbf{w}, \quad (3.122)$$

where

$$E(\mathbf{w}) = \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{w}\|^2 + \frac{\alpha}{2} \mathbf{w}^\top \mathbf{w}. \quad (3.123)$$

The first term of the definition of $E(\mathbf{w})$ can be written as

$$\begin{aligned} & \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N - \boldsymbol{\Phi}(\mathbf{w} - \mathbf{m}_N)\|^2 \\ &= \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N\|^2 - \beta(\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N)^\top \boldsymbol{\Phi}(\mathbf{w} - \mathbf{m}_N) + \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_N)^\top \boldsymbol{\Phi}^\top \boldsymbol{\Phi}(\mathbf{w} - \mathbf{m}_N). \end{aligned} \quad (3.124)$$

Similarly, the second term can be written as

$$\begin{aligned} & \frac{\alpha}{2} (\mathbf{w} - \mathbf{m}_N + \mathbf{m}_N)^\top (\mathbf{w} - \mathbf{m}_N + \mathbf{m}_N) \\ &= \frac{\alpha}{2} (\mathbf{w} - \mathbf{m}_N)^\top (\mathbf{w} - \mathbf{m}_N) + \alpha \mathbf{m}_N^\top (\mathbf{w} - \mathbf{m}_N) + \frac{\alpha}{2} \mathbf{m}_N^\top \mathbf{m}_N. \end{aligned} \quad (3.125)$$

Here,

$$\begin{aligned} & -\beta(\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N)^\top \boldsymbol{\Phi}(\mathbf{w} - \mathbf{m}_N) + \alpha \mathbf{m}_N^\top (\mathbf{w} - \mathbf{m}_N) \\ &= (-\beta \boldsymbol{\Phi}^\top \mathbf{t} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi} \mathbf{m}_N + \alpha \mathbf{m}_N)^\top (\mathbf{w} - \mathbf{m}_N). \end{aligned} \quad (3.126)$$

By the definitions of \mathbf{m}_N and \mathbf{S}_N above, the right hand can be written as

$$(-\beta \boldsymbol{\Phi}^\top \mathbf{t} + \mathbf{S}_N^{-1} \mathbf{m}_N)^\top (\mathbf{w} - \mathbf{m}_N) = 0. \quad (3.127)$$

Therefore,

$$E(\mathbf{w}) = E(\mathbf{m}_N) + \frac{1}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N). \quad (3.128)$$

3.19

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}), \end{aligned} \quad (3.129)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.130)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.131)$$

By 3.17,

$$p(\mathbf{t}) = \left(\frac{\beta}{2\pi}\right)^{\frac{N}{2}} \left(\frac{\alpha}{2\pi}\right)^{\frac{M}{2}} \int \exp(-E(\mathbf{w})) d\mathbf{w}, \quad (3.132)$$

where

$$E(\mathbf{w}) = \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{w}\|^2 + \frac{\alpha}{2} \mathbf{w}^\top \mathbf{w}. \quad (3.133)$$

By 3.18,

$$E(\mathbf{w}) = E(\mathbf{m}_N) + \frac{1}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N). \quad (3.134)$$

Therefore, the integral in the expression above of $p(\mathbf{t})$ can be written as

$$\begin{aligned} & \exp(-E(\mathbf{m}_N)) \int \exp\left(-\frac{1}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N)\right) d\mathbf{w} \\ &= (2\pi)^{\frac{M}{2}} (\det \mathbf{S}_N)^{\frac{1}{2}} \exp(-E(\mathbf{m}_N)). \end{aligned} \quad (3.135)$$

Thus,

$$\ln p(\mathbf{t}) = \frac{M}{2} \ln \alpha + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi) + \frac{1}{2} \ln(\det \mathbf{S}_N) - E(\mathbf{m}_N). \quad (3.136)$$

3.20

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}), \end{aligned} \quad (3.137)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.138)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.139)$$

By 3.19,

$$\ln p(\mathbf{t}) = \frac{M}{2} \ln \alpha + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi) + \frac{1}{2} \ln(\det \mathbf{S}_N) - E(\mathbf{m}_N), \quad (3.140)$$

where

$$E(\mathbf{m}_N) = \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N\|^2 + \frac{\alpha}{2} \mathbf{m}_N^\top \mathbf{m}_N. \quad (3.141)$$

Let $\mathbf{u}_1, \dots, \mathbf{u}_M$ be eigenvectors of $\beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}$ such that

$$\beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi} \mathbf{u}_m = \lambda_m \mathbf{u}_m. \quad (3.142)$$

Then,

$$\mathbf{S}_N^{-1} \mathbf{u}_m = (\alpha + \lambda_m) \mathbf{u}_m, \quad (3.143)$$

so that

$$\det \mathbf{S}_N = \prod_{m=1}^M \frac{1}{\alpha + \lambda_m}. \quad (3.144)$$

Therefore, setting the derivative of $\ln p(\mathbf{t}|\alpha, \beta)$ with respect to α to zero gives

$$0 = \frac{M}{2\alpha} - \frac{1}{2} \sum_{m=1}^M \frac{1}{\alpha + \lambda_m} - \frac{1}{2} \mathbf{m}_N^\top \mathbf{m}_N. \quad (3.145)$$

Multiplying both sides by 2α gives

$$\alpha \mathbf{m}_N^\top \mathbf{m}_N = M - \sum_{m=1}^M \frac{\alpha}{\alpha + \lambda_m}. \quad (3.146)$$

The right hand side can be written as

$$\sum_{m=1}^M \left(1 - \frac{\alpha}{\alpha + \lambda_m} \right) = \sum_{m=1}^M \frac{\lambda_m}{\alpha + \lambda_m}. \quad (3.147)$$

Thus,

$$\alpha = \frac{\gamma}{\mathbf{m}_N^\top \mathbf{m}_N}, \quad (3.148)$$

where

$$\gamma = \sum_{m=1}^M \frac{\lambda_i}{\alpha + \lambda_m}. \quad (3.149)$$

3.21

(a)

Let Σ be a $M \times M$ real symmetric matrix such that

$$\Sigma \mathbf{u}_m = \lambda_m \mathbf{u}_m, \quad (3.150)$$

where $\mathbf{u}_1, \dots, \mathbf{u}_M$ are unit vectors. Let

$$\begin{aligned} \Lambda &= \text{diag}(\lambda_1, \dots, \lambda_M), \\ \mathbf{U} &= [\mathbf{u}_1 \cdots \mathbf{u}_M]. \end{aligned} \quad (3.151)$$

By 2.19,

$$\begin{aligned} \Sigma &= \mathbf{U} \Lambda \mathbf{U}^\top, \\ \mathbf{U}^\top \mathbf{U} &= \mathbf{I}. \end{aligned} \quad (3.152)$$

Therefore,

$$\det \Sigma = \prod_{m=1}^M \lambda_m, \quad (3.153)$$

so that

$$\ln(\det \Sigma) = \sum_{m=1}^M \ln \lambda_i. \quad (3.154)$$

Then,

$$\frac{\partial}{\partial \alpha} \ln(\det \Sigma) = \sum_{m=1}^M \frac{\partial \lambda_m}{\partial \alpha} \frac{1}{\lambda_m}. \quad (3.155)$$

Thus,

$$\frac{\partial}{\partial \alpha} \ln(\det \Sigma) = \text{tr} \left(\Sigma^{-1} \frac{\partial}{\partial \alpha} \Sigma \right). \quad (3.156)$$

(b)

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}), \end{aligned} \quad (3.157)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.158)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.159)$$

By 3.19,

$$\ln p(\mathbf{t}) = \frac{M}{2} \ln \alpha + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi) + \frac{1}{2} \ln(\det \mathbf{S}_N) - E(\mathbf{m}_N), \quad (3.160)$$

where

$$E(\mathbf{m}_N) = \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N\|^2 + \frac{\alpha}{2} \mathbf{m}_N^\top \mathbf{m}_N. \quad (3.161)$$

By 3.21(a),

$$\frac{\partial}{\partial \alpha} \ln(\det \mathbf{S}_N^{-1}) = \text{tr}(\mathbf{S}_N). \quad (3.162)$$

The right hand side can be written as

$$\sum_{m=1}^M \frac{1}{\alpha + \lambda_m}, \quad (3.163)$$

where $\lambda_1, \dots, \lambda_M$ are eigenvalues of $\beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}$. Therefore, setting the derivative of $\ln p(\mathbf{t})$ with respect to α to zero gives

$$0 = \frac{M}{2\alpha} - \frac{1}{2} \sum_{m=1}^M \frac{1}{\alpha + \lambda_m} - \frac{1}{2} \mathbf{m}_N^\top \mathbf{m}_N, \quad (3.164)$$

Thus,

$$\alpha = \frac{\gamma}{\mathbf{m}_N^\top \mathbf{m}_N}, \quad (3.165)$$

where

$$\gamma = \sum_{m=1}^M \frac{\lambda_m}{\alpha + \lambda_m}. \quad (3.166)$$

3.22

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n|\mathbf{w}) &= \mathcal{N}(t_n|\mathbf{w}^\top \boldsymbol{\phi}(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}) &= \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}), \end{aligned} \quad (3.167)$$

where \mathbf{w} and $\boldsymbol{\phi}$ are vectors in M dimensions. By 3.7,

$$p(\mathbf{w}|\mathbf{t}) = \mathcal{N}(\mathbf{w}|\mathbf{m}_N, \mathbf{S}_N), \quad (3.168)$$

where

$$\begin{aligned} \mathbf{m}_N &= \beta \mathbf{S}_N \boldsymbol{\Phi}^\top \mathbf{t}, \\ \mathbf{S}_N^{-1} &= \alpha \mathbf{I} + \beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}. \end{aligned} \quad (3.169)$$

By 3.19,

$$\ln p(\mathbf{t}) = \frac{M}{2} \ln \alpha + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi) + \frac{1}{2} \ln(\det \mathbf{S}_N) - E(\mathbf{m}_N), \quad (3.170)$$

where

$$E(\mathbf{m}_N) = \frac{\beta}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N\|^2 + \frac{\alpha}{2} \mathbf{m}_N^\top \mathbf{m}_N. \quad (3.171)$$

By 3.21(a),

$$\frac{\partial}{\partial \beta} \ln(\det \mathbf{S}_N^{-1}) = \text{tr}(\mathbf{S}_N \boldsymbol{\Phi}^\top \boldsymbol{\Phi}). \quad (3.172)$$

Since

$$\mathbf{S}_N \boldsymbol{\Phi}^\top \boldsymbol{\Phi} = \frac{1}{\beta} (\mathbf{I} - \alpha \mathbf{S}_N), \quad (3.173)$$

the right hand side can be written as

$$\frac{1}{\beta} \left(M - \alpha \sum_{m=1}^M \frac{1}{\alpha + \lambda_m} \right) = \frac{1}{\beta} \sum_{m=1}^M \frac{\lambda_m}{\alpha + \lambda_m}, \quad (3.174)$$

where $\lambda_1, \dots, \lambda_M$ are eigenvalues of $\beta \boldsymbol{\Phi}^\top \boldsymbol{\Phi}$. Therefore, setting the derivative of $\ln p(\mathbf{t})$ with respect to β to zero gives

$$0 = \frac{N}{2\beta} - \frac{1}{2\beta} \sum_{m=1}^M \frac{\lambda_i}{\alpha + \lambda_m} - \frac{1}{2} \|\mathbf{t} - \boldsymbol{\Phi} \mathbf{m}_N\|^2. \quad (3.175)$$

Thus,

$$\beta = \frac{N - \gamma}{\|\mathbf{t} - \Phi \mathbf{m}_N\|^2}, \quad (3.176)$$

where

$$\gamma = \sum_{m=1}^M \frac{\lambda_m}{\alpha + \lambda_m}. \quad (3.177)$$

3.23

Let t_1, \dots, t_N be variables such that

$$\begin{aligned} p(t_n | \mathbf{w}, \beta) &= \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}), \\ p(\mathbf{w}, \beta) &= \mathcal{N}(\mathbf{w} | \mathbf{m}_0, \beta^{-1} \mathbf{S}_0) \text{Gam}(\beta | a_0, b_0), \end{aligned} \quad (3.178)$$

where \mathbf{w} and ϕ are vectors in M dimensions. By marginalisation,

$$p(\mathbf{t}) = \int \int p(\mathbf{t} | \mathbf{w}, \beta) p(\mathbf{w}, \beta) d\mathbf{w} d\beta. \quad (3.179)$$

The right hand side can be written as

$$\int \left(\int \left(\prod_{n=1}^N \mathcal{N}(t_n | \mathbf{w}^\top \phi(\mathbf{x}_n), \beta^{-1}) \right) \mathcal{N}(\mathbf{w} | \mathbf{m}_0, \beta^{-1} \mathbf{S}_0) d\mathbf{w} \right) \text{Gam}(\beta | a_0, b_0) d\beta. \quad (3.180)$$

The logarithm of the integrand with respect to \mathbf{w} can be written as

$$\begin{aligned} & -\frac{N}{2} \ln(2\pi) - \frac{N}{2} \ln \beta^{-1} - \frac{\beta}{2} \sum_{n=1}^N (t_n - \mathbf{w}^\top \phi(\mathbf{x}_n))^2 \\ & -\frac{M}{2} \ln(2\pi) - \frac{1}{2} \ln \det(\beta^{-1} \mathbf{S}_0) - \frac{\beta}{2} (\mathbf{w} - \mathbf{m}_0)^\top \mathbf{S}_0^{-1} (\mathbf{w} - \mathbf{m}_0) \\ & = -\frac{N+M}{2} \ln(2\pi) + \frac{N+M}{2} \ln \beta - \frac{1}{2} \ln(\det \mathbf{S}_0) \\ & \quad - \frac{\beta}{2} \mathbf{w}^\top (\mathbf{S}_0^{-1} + \Phi^\top \Phi) \mathbf{w} + \beta \mathbf{w}^\top (\mathbf{S}_0^{-1} \mathbf{m}_0 + \Phi^\top \mathbf{t}) - \frac{\beta}{2} \|\mathbf{t}\|^2 - \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0. \end{aligned} \quad (3.181)$$

The right hand side can be written as

$$\begin{aligned} & -\frac{N+M}{2} \ln(2\pi) + \frac{N+M}{2} \ln \beta - \frac{1}{2} \ln(\det \mathbf{S}_0) \\ & -\frac{\beta}{2} (\mathbf{w} - \mathbf{m}_N)^\top \mathbf{S}_N^{-1} (\mathbf{w} - \mathbf{m}_N) + \frac{\beta}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N - \frac{\beta}{2} \|\mathbf{t}\|^2 - \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0, \end{aligned} \quad (3.182)$$

where

$$\begin{aligned}\mathbf{m}_N &= \mathbf{S}_N (\mathbf{S}_0^{-1} \mathbf{m}_0 + \Phi^\top \mathbf{t}), \\ \mathbf{S}_N^{-1} &= \mathbf{S}_0^{-1} + \Phi^\top \Phi.\end{aligned}\tag{3.183}$$

Therefore, the logarithm of the integral with respect to \mathbf{w} can be written as

$$\begin{aligned}& -\frac{N}{2} \ln(2\pi) + \frac{N}{2} \ln \beta - \frac{1}{2} \ln(\det \mathbf{S}_0) + \frac{1}{2} \ln(\det \mathbf{S}_N) \\ & + \frac{\beta}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N - \frac{\beta}{2} \|\mathbf{t}\|^2 - \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0.\end{aligned}\tag{3.184}$$

Then, the logarithm of the integrand with respect to β can be written as

$$\begin{aligned}& -\frac{N}{2} \ln(2\pi) + \frac{N}{2} \ln \beta - \frac{1}{2} \ln(\det \mathbf{S}_0) + \frac{1}{2} \ln(\det \mathbf{S}_N) \\ & + \frac{\beta}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N - \frac{\beta}{2} \|\mathbf{t}\|^2 - \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0 \\ & - \ln \Gamma(a_0) + a_0 \ln b_0 + (a_0 - 1) \ln \beta - b_0 \beta \\ & = -\frac{N}{2} \ln(2\pi) - \frac{1}{2} \ln(\det \mathbf{S}_0) + \frac{1}{2} \ln(\det \mathbf{S}_N) \\ & - \ln \Gamma(a_0) + a_0 \ln b_0 + (a_N - 1) \ln \beta - b_N \beta,\end{aligned}\tag{3.185}$$

where

$$\begin{aligned}a_N &= a_0 + \frac{N}{2}, \\ b_N &= b_0 + \frac{\beta}{2} \|\mathbf{t}\|^2 + \frac{\beta}{2} \mathbf{m}_0^\top \mathbf{S}_0^{-1} \mathbf{m}_0 - \frac{\beta}{2} \mathbf{m}_N^\top \mathbf{S}_N^{-1} \mathbf{m}_N.\end{aligned}\tag{3.186}$$

Therefore, the logarithm of the integral with respect to β can be written as

$$-\frac{N}{2} \ln(2\pi) - \frac{1}{2} \ln(\det \mathbf{S}_0) + \frac{1}{2} \ln(\det \mathbf{S}_N) - \ln \Gamma(a_0) + a_0 \ln b_0 + \ln \Gamma(a_N) - a_N \ln b_N.\tag{3.187}$$

Thus,

$$p(\mathbf{t}) = (2\pi)^{-\frac{N}{2}} \left(\frac{\det \mathbf{S}_N}{\det \mathbf{S}_0} \right)^{\frac{1}{2}} \frac{\Gamma(a_N)}{\Gamma(a_0)} \frac{b_0^{a_0}}{b_N^{a_N}}.\tag{3.188}$$

3.24

Refer to 3.12.

4 Linear Models for Classification

4.1

Let x_1, \dots, x_M and y_1, \dots, y_N be two sets of data points. Then, the corresponding convex hulls are defined as the sets of all points \mathbf{x} and \mathbf{y} such that

$$\begin{aligned}\mathbf{x} &= \sum_{m=1}^M \alpha_m \mathbf{x}_m, \\ \mathbf{y} &= \sum_{n=1}^N \beta_n \mathbf{y}_n,\end{aligned}\tag{4.1}$$

where

$$\begin{aligned}\sum_{m=1}^M \alpha_m &= \sum_{n=1}^N \beta_n = 1, \\ \alpha_m &\geq 0, \beta_n \geq 0.\end{aligned}\tag{4.2}$$

Let us assume that $\alpha_1, \dots, \alpha_M$ and β_1, \dots, β_N below are subject to the constraints above.

If the convex hulls intersect, then there exist $\alpha_1, \dots, \alpha_M$ and β_1, \dots, β_N such that

$$\sum_{m=1}^M \alpha_m \mathbf{x}_m = \sum_{n=1}^N \beta_n \mathbf{y}_n.\tag{4.3}$$

Then,

$$\sum_{m=1}^M \alpha_m (\hat{\mathbf{w}}^\top \mathbf{x}_m + w_0) = \hat{\mathbf{w}}^\top \sum_{m=1}^M \alpha_m \mathbf{x}_m + w_0 \sum_m \alpha_m,\tag{4.4}$$

for any $\hat{\mathbf{w}}$ and w_0 . The right hand side can be written as

$$\hat{\mathbf{w}}^\top \sum_{n=1}^N \beta_n \mathbf{y}_n + w_0 \sum_{n=1}^N \beta_n = \sum_{n=1}^N \beta_n (\hat{\mathbf{w}}^\top \mathbf{y}_n + w_0).\tag{4.5}$$

Therefore, there do not exist $\hat{\mathbf{w}}$ and w_0 such that

$$\begin{aligned}\hat{\mathbf{w}}^\top \mathbf{x}_m + w_0 &> 0, \\ \hat{\mathbf{w}}^\top \mathbf{y}_n + w_0 &< 0.\end{aligned}\tag{4.6}$$

Conversely, if there exist $\hat{\mathbf{w}}$ and w_0 such that

$$\begin{aligned}\hat{\mathbf{w}}^\top \mathbf{x}_m + w_0 &> 0, \\ \hat{\mathbf{w}}^\top \mathbf{y}_n + w_0 &< 0,\end{aligned}\tag{4.7}$$

then

$$\begin{aligned}\sum_{m=1}^M \alpha_m (\hat{\mathbf{w}}^\top \mathbf{x}_m + w_0) &> 0, \\ \sum_{n=1}^N \beta_n (\hat{\mathbf{w}}^\top \mathbf{y}_n + w_0) &< 0.\end{aligned}\tag{4.8}$$

The left hand sides can be written as

$$\begin{aligned}\hat{\mathbf{w}}^\top \sum_{m=1}^M \alpha_m \mathbf{x}_m + w_0 \sum_{m=1}^M \alpha_m &= \hat{\mathbf{w}}^\top \sum_{m=1}^M \alpha_m \mathbf{x}_m + w_0, \\ \hat{\mathbf{w}}^\top \sum_{n=1}^N \beta_n \mathbf{y}_n + w_0 \sum_{n=1}^N \beta_n &= \hat{\mathbf{w}}^\top \sum_{n=1}^N \beta_n \mathbf{y}_n + w_0.\end{aligned}\tag{4.9}$$

Therefore, there do not exist $\alpha_1, \dots, \alpha_M$ and β_1, \dots, β_N such that

$$\sum_{m=1}^M \alpha_m \mathbf{x}_m = \sum_{n=1}^N \beta_n \mathbf{y}_n.\tag{4.10}$$

Thus, the convex hulls do not intersect.

4.2 (Incomplete)

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ and $\mathbf{w}_1, \dots, \mathbf{w}_K$ are variables in M dimensions and $\mathbf{t}_1, \dots, \mathbf{t}_N$ are ones in K dimensions. Let

$$E(\tilde{\mathbf{W}}) = \frac{1}{2} \text{tr} \left((\tilde{\mathbf{X}}\tilde{\mathbf{W}} - \mathbf{T})^\top (\tilde{\mathbf{X}}\tilde{\mathbf{W}} - \mathbf{T}) \right),\tag{4.11}$$

where

$$\tilde{\mathbf{X}} = \begin{bmatrix} 1 & \mathbf{x}_1^\top \\ \vdots & \vdots \\ 1 & \mathbf{x}_N^\top \end{bmatrix},$$

$$\tilde{\mathbf{W}} = \begin{bmatrix} w_{10} & \cdots & w_{K0} \\ \mathbf{w}_1 & \cdots & \mathbf{w}_K \end{bmatrix}$$

and

$$\mathbf{T} = \begin{bmatrix} \mathbf{t}_1^\top \\ \vdots \\ \mathbf{t}_N^\top \end{bmatrix}.$$

Setting the derivative with respect to $\tilde{\mathbf{W}}$ to zero gives

$$\mathbf{0} = \tilde{\mathbf{X}}^\top (\tilde{\mathbf{X}} \tilde{\mathbf{W}} - \mathbf{T}). \quad (4.12)$$

Therefore,

$$\underset{\tilde{\mathbf{W}}}{\operatorname{argmin}} E(\tilde{\mathbf{W}}) = \left(\tilde{\mathbf{X}}^\top \tilde{\mathbf{X}} \right)^{-1} \tilde{\mathbf{X}}^\top \mathbf{T}. \quad (4.13)$$

Let $\tilde{\mathbf{W}}^*$ denote the least-square solution above. If

$$\mathbf{a}^\top \mathbf{t}_n + b = 0, \quad (4.14)$$

for $n = 1, \dots, N$, then

$$\mathbf{a}^\top (\tilde{\mathbf{W}}^*)^\top \tilde{\mathbf{x}} + b = 0? \quad (4.15)$$

where $\tilde{\mathbf{x}}$ is a variable in $M + 1$ dimensions whose first element is 1.

$$(\tilde{\mathbf{W}}^*)^\top \tilde{\mathbf{x}} = \mathbf{T}^\top \tilde{\mathbf{X}} \left(\tilde{\mathbf{X}}^\top \tilde{\mathbf{X}} \right)^{-1} \tilde{\mathbf{x}}. \quad (4.16)$$

4.3 (Incomplete)

4.4

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables and let

$$\mathbf{m}_k = \frac{1}{N_k} \sum_{n \in \mathcal{C}_k} \mathbf{x}_n, \quad (4.17)$$

where N_k is the number of \mathbf{x}_n such that n is in \mathcal{C}_k . Setting the derivatives of

$$\mathbf{w}^\top (\mathbf{m}_2 - \mathbf{m}_1) + \lambda (\|\mathbf{w}\|^2 - 1) \quad (4.18)$$

with respect to \mathbf{w} and λ to zero gives

$$\begin{aligned} \mathbf{m}_2 - \mathbf{m}_1 + 2\lambda \mathbf{w} &= \mathbf{0}, \\ \|\mathbf{w}\|^2 - 1 &= 0. \end{aligned} \quad (4.19)$$

Therefore, $\mathbf{w}^\top(\mathbf{m}_2 - \mathbf{m}_1)$ under the constraint

$$\|\mathbf{w}\|^2 = 1 \quad (4.20)$$

is maximised if

$$\mathbf{w} \propto \mathbf{m}_2 - \mathbf{m}_1. \quad (4.21)$$

4.5

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables and let

$$\mathbf{m}_k = \frac{1}{N_k} \sum_{n \in \mathcal{C}_k} \mathbf{x}_n, \quad (4.22)$$

where N_k is the number of \mathbf{x}_n such that n is in \mathcal{C}_k . Let

$$J(\mathbf{w}) = \frac{(m_2 - m_1)^2}{s_1^2 + s_2^2}, \quad (4.23)$$

where

$$\begin{aligned} s_k^2 &= \sum_{n \in \mathcal{C}_k} (y_n - m_k)^2, \\ y_n &= \mathbf{w}^\top \mathbf{x}_n, \\ m_k &= \mathbf{w}^\top \mathbf{m}_k. \end{aligned} \quad (4.24)$$

Then, $J(\mathbf{w})$ can be written as

$$\frac{(\mathbf{w}^\top(\mathbf{m}_2 - \mathbf{m}_1))^2}{\sum_{n \in \mathcal{C}_1} (\mathbf{w}^\top(\mathbf{x}_n - \mathbf{m}_1))^2 + \sum_{n \in \mathcal{C}_2} (\mathbf{w}^\top(\mathbf{x}_n - \mathbf{m}_2))^2} = \frac{\mathbf{w}^\top \mathbf{S}_B \mathbf{w}}{\mathbf{w}^\top \mathbf{S}_W \mathbf{w}}, \quad (4.25)$$

where

$$\begin{aligned} \mathbf{S}_B &= (\mathbf{m}_2 - \mathbf{m}_1)(\mathbf{m}_2 - \mathbf{m}_1)^\top, \\ \mathbf{S}_W &= \sum_{n \in \mathcal{C}_1} (\mathbf{x}_n - \mathbf{m}_1)(\mathbf{x}_n - \mathbf{m}_1)^\top + \sum_{n \in \mathcal{C}_2} (\mathbf{x}_n - \mathbf{m}_2)(\mathbf{x}_n - \mathbf{m}_2)^\top. \end{aligned} \quad (4.26)$$

4.6 (Incomplete)

Let $\mathbf{x}_1, \dots, \mathbf{x}_N$ be variables and let

$$\mathbf{m}_k = \frac{1}{N_k} \sum_{n \in \mathcal{C}_k} \mathbf{x}_n, \quad (4.27)$$

where N_k is the number of \mathbf{x}_n such that n is in \mathcal{C}_k . Let

$$J(\mathbf{w}) = \frac{(m_2 - m_1)^2}{s_1^2 + s_2^2}, \quad (4.28)$$

where

$$\begin{aligned} s_k^2 &= \sum_{n \in \mathcal{C}_k} (y_n - m_k)^2, \\ y_n &= \mathbf{w}^\top \mathbf{x}_n, \\ m_k &= \mathbf{w}^\top \mathbf{m}_k. \end{aligned} \quad (4.29)$$

Then, by 4.5,

$$J(\mathbf{w}) = \frac{\mathbf{w}^\top \mathbf{S}_B \mathbf{w}}{\mathbf{w}^\top \mathbf{S}_W \mathbf{w}}, \quad (4.30)$$

where

$$\begin{aligned} \mathbf{S}_B &= (\mathbf{m}_2 - \mathbf{m}_1)(\mathbf{m}_2 - \mathbf{m}_1)^\top, \\ \mathbf{S}_W &= \sum_{n \in \mathcal{C}_1} (\mathbf{x}_n - \mathbf{m}_1)(\mathbf{x}_n - \mathbf{m}_1)^\top + \sum_{n \in \mathcal{C}_2} (\mathbf{x}_n - \mathbf{m}_2)(\mathbf{x}_n - \mathbf{m}_2)^\top. \end{aligned} \quad (4.31)$$